

Modeling the Historical Disturbance Regimes of NYC

Historical Disturbance Regimes

(Lorimer 1980, Foster et al. 1998, Pan et al. (2011), Lorimer and White 2003)
(Day 1953, Russell 1997, 1997, Krech III 1999,)

Fire

Custom Fuel Models

Custom fuel models are based on the defined set that comes packaged with FARSITE. Hardwood forest types are based on model F9. Where average loads for 1 hr 10 hr and 100 hr fuels were available for northeastern forest types the substituted (Reinhardt and Crookston 2015). The remaining fuel attributes remained the same as the base model. The USFS community types were cross-walked to the NY State Heritage community classification so fuels could be assigned for our landscape.

Fuel Moisture

Region specific initial fuel moistures (Reinhardt and Crookston 2015). All freshwater wetland communities (marshes, shrub-swamps and forested wetlands) were initialized with a wet fuel profile, all other communities with burnable fuel types were initialized with a moist fuel profile.

Size Class	Very Dry	Dry	Moist	Wet
1 hr	5	7	10	19
10 hr	8	9	13	29
100 hr	12	14	17	22
Live woody	89	105	135	140
Liver herbaceous	60	82	116	120

Initial Conditions - Forest Age

Forest communities were initialized with an age drawn from the following normal distribution (Loewenstein et al. 2000, Pan et al. 2011). All no forest types were initialized with a forest age of 0




Figure 1: age class distribution

Initial Conditions - Canopy

Randomized (using uniform distribution) within canopy classes:

- grassland $0 < \text{canopy} < 20$
- shrubland $20 < \text{canopy} < 60$
- forest $60 < \text{canopy}$

Modeling Expected Frequencies

use of Poisson distribution to model expected forest fire frequency (Johnson 2001, Yang et al. 2008)

fire size and frequency literature

effects of fires on temperate forests (Kozlowski 2012) power law(Reed and McKelvey 2002, Stephens 2005, Cui and Perera 2008)

Expected Lightning Frequency

Loope and Anderton (1998) Expected number of lightning and human caused fires are based on areal frequencies from USFS wildfire records in region 9 (U.S. Northeast) between 1940 and 2000 (Stephens 2005). These values were converted from the given units (frequency/400000 ha)/yr to (frequency/km²)/yr.

Region	Lightning		Human	
	Ha burned	No. fires	Ha burned	No. fires
9	11.840	2.170	327.990	39.950

	Lightning		Human	
	Area burned	No. Fires	Area burned	No. Fires
1 hectare	0.0000296	0.000005425	0.000819975	0.000099875
per km ²	0.00296	0.0005425	0.0819975	0.0099875

Critical Rainfall

In the model, fire spread was stopped when it encountered one of the following conditions: (1) a non-flammable type of land cover; (2) boundaries of the region; and (3) when rainfall exceeded a certain critical amount. By assuming that a daily precipitation of 30 mm or more would stop a fire, the R Crit in Eq. (2) was estimated as 0.026 (the proportion of total number of days that has daily precipitation of 30 mm or more) from the historical precipitation data of the Edison weather station. (Li 2000)

Tree Allometry

Description	Equation	Reference
Tree Height	$TH = 44 * \ln(Age) - 93$	(Bean and Sanderson 2008)
DBH	$DBH = (Age - 34.44)/1.18$	(Loewenstein et al. 2000)
Crown Ratio	$CR = 0.4$	(Bean and Sanderson 2008)
Bark Thickness	$BT = vsp * DBH$	(Reinhardt and Crookston 2015)

- Communities to Bark Thickness * bark thickness multipliers for each community, for communities with co-dominate species the average bark thickness was calculated (Reinhardt and Crookston 2015).

community	dominant tree species	vsp scaler
Floodplain forest	avg(sliver maple, sycamore, American elm)	0.032
Red Maple Hardwood Swamp	red maple	0.028
Coastal Plain Atlantic Cedar Swamp	Atlantic cedar	0.025
Pitch pine - scrub oak barrens	avg(pitch pine, oak spp)	0.045
Chestnut oak forest	avg(American chestnut, oak spp)	0.043
Coastal oak beech forest	avg(oak spp, beech)	0.035
Coastal oak hickory forest	avg(oak spp, hickory spp)	0.045
Oak tulip forest	avg(oak spp, yellow-poplar)	0.038
Appalachian oak pine forest	avg(oak spp, pine spp)	0.038
Hemlock northern hardwood forest	hemlock	0.045
Inland Atlantic Cedar Swamp	Atlantic white cedar	0.025
Red maple black gum swamp	avg(red maple, black gum)	0.034
Red maple sweetgum swamp	avg(red maple, sweetgum)	0.032
Maritime holly forest	holly	0.042
Post oak black jack oak barrens	post oak	0.044
Appalachian oak hickory forest	avg(oak spp, hickory spp)	0.045
Beech maple mesic forest	avg(beech, sugar maple)	0.029
Successional maritime hardwoods	other hardwoods	0.044
Successional hardwood forest	other hardwoods	0.044

Fire Mortality Equations

Scorch Height

[1]

$$SH = 3.1817(FL^{1.4503})$$

(Bean and Sanderson 2008)

Crown Kill

[2]

$$CK = 41.961(100(\ln(\frac{SH - CH}{CL}))) - 89.721$$

(Bean and Sanderson 2008)

Percent Mortality

[3]

$$P_m = \frac{1.0}{1.0 + e^{-1.941 + 6.316(1.0 - e^{-BT}) - 0.000535CK^2}}$$

(Bean and Sanderson 2008)

Horticulture

Archaeological Evidence for Gardening

(Kraft 2001, Cantwell and Wall (2001), Benison (1997))

Ethnohistorical

(Ascher 1860, Danckaerts and Sluyter (1867))

crop	yield (kg/ha)	calories/100 g	calories/kg	calories/ha
corn	1720	365	3650	6278000
beans (<i>Phaseolus vulgaris</i>)	110	33	330	36300
squash (Indian squash)	80	16	160	12800

corn-bean-squash ploy-culture yields in Tabasco, Mexico (Gliessman et al. 1998)
pg 224 table 15.3

Caloric density [USDA Basic Reports Agricultural Research Services]

Agricultural Dependency Scenarios

caloric requirements (Speth and Spielmann 1983)

Dependency	calories/person/yr	ha/person	m ² /person
15%	94060.50	0.01	14.87
30%	188121.00	0.03	29.73
60%	376242.00	0.06	59.47
100%	627070.00	0.10	99.11

Beaver

Succession

We used a 4 sere sequence for the freshwater wetland pathway (Allen 1983, Naiman et al. 1988, Johnston and Naiman 1990, *Ecology of Red Maple Swamps in the Glaciated Northeast* 1993, Hay 2010, Logofet et al. 2016). Conversion from non-wetland community to active pond can occur along any perennial streams where the stream gradient is less than 15 degrees. Due to the temporal

scale of our study, forested wetlands are treated as a terminal community in this series (*Ecology of Red Maple Swamps in the Glaciated Northeast* 1993). This rule defines beaver caused disturbance as a unidirectional change in successional trajectory. A non-wetland communities can be converted into a wetland type but this conversion cannot be reversed.

active beaver pond → emergent marsh → shrub swamp → forested wetland

Model Parameters

Parameter	Value		Source
probability of abandonment	0.10		(Logofet et al. 2016)
colony density	0.4	colonies/km ²	(Naiman et al. 1988)
territory (minimum distance)	1000	m	Naiman et al. (1988); Allen (1983)]

Analysis

time since fire (Johnson 2001) fire size frequency hist (Malamud et al. 1998, Reed and McKelvey 2002, Cui and Perera 2008)

References

- Allen, A. W. 1983. Habitat suitability index models: Beaver. Western Energy; Land Use Team, Division of Biological Service, Research; Development, Fish; Wildlife Service, US Department of the Interior.
- Ascher, G. 1860. Henry Hudson the Navigator. Hakluyt Society, London.
- Bean, W. T., and E. W. Sanderson. 2008. Using a spatially explicit ecological model to test scenarios of fire use by Native Americans: An example from the Harlem Plains, New York, NY. *Ecological Modelling* 211:301–308.
- Benison, C. 1997. Horticulture and the Maintenance of Social Complexity in Late Woodland Southeastern New England. *North American Archaeologist* 18:1–17.
- Cantwell, A.-M., and D. d. Wall. 2001. *Unearthing Gotham: The Archaeology of New York City*. Yale University Press, New Haven, CT.
- Cui, W., and A. H. Perera. 2008. What do we know about forest fire size distribution, and why is this knowledge useful for forest management? *International Journal of Wildland Fire* 17:234.

- Danckaerts, J., and P. Sluyter. 1867. *Journal of a Voyage to New York: And a Tour in Several of the American Colonies in 1679-80*. The Long Island Historical Society, Brooklyn.
- Day, G. M. 1953. The Indian as an Ecological Factor in the Northeastern Forest. *Ecology* 34:329–346.
- Ecology of Red Maple Swamps in the Glaciated Northeast: A Community Profile. 1993. U.S. Fish; Wildlife Service.
- Foster, D. R., G. Motzkin, and B. Slater. 1998. Land-Use History as Long-Term Broad-Scale Disturbance: Regional Forest Dynamics in Central New England. *Ecosystems* 1:96–119.
- Gliessman, S. R., E. Engles, and R. Krieger. 1998. *Agroecology: Ecological Processes in Sustainable Agriculture*. CRC Press.
- Hay, K. G. 2010. Succession of Beaver Ponds in Colorado 50 Years After Beaver Removal. *The Journal of Wildlife Management* 74:1732–1736.
- Johnson, E. A. 2001. *Forest Fires: Behavior and Ecological Effects*. Academic Press.
- Johnston, C. A., and R. J. Naiman. 1990. The use of a geographic information system to analyze long-term landscape alteration by beaver. *Landscape Ecology* 4:5–19.
- Kozlowski, T. T. 2012. *Fire and Ecosystems*. Elsevier.
- Kraft, H. C. 2001. *The Lenape-Delaware Indian heritage: 10,000 B.C.- A.D. 2000*. Lenape Books, [S.l.].
- Krech III, S. 1999. *The Ecological Indian*. W.W. Norton & Company, New York.
- Li, C. 2000. Reconstruction of natural fire regimes through ecological modelling. *Ecological Modelling* 134:129–144.
- Loewenstein, E. F., P. S. Johnson, and H. E. Garrett. 2000. Age and diameter structure of a managed uneven-aged oak forest. *Canadian Journal of Forest Research* 30:1060–1070.
- Logofet, D. O., O. I. Evstigneev, A. A. Aleinikov, and A. O. Morozova. 2016. Succession caused by beaver (*Castor fiber* L.) life activity: II. a refined markov model. *Biology Bulletin Reviews* 6:39–56.
- Loope, W. L., and J. B. Anderton. 1998. Human vs. Lightning Ignition of Presettlement Surface Fires in Coastal Pine Forests of the Upper Great Lakes. *The American Midland Naturalist* 140:206–218.
- Lorimer, C. G. 1980. Age Structure and Disturbance History of a Southern Appalachian Virgin Forest. *Ecology* 61:1169–1184.

- Lorimer, C. G., and A. S. White. 2003. Scale and frequency of natural disturbances in the northeastern US: Implications for early successional forest habitats and regional age distributions. *Forest Ecology and Management* 185:41–64.
- Malamud, B. D., G. Morein, and D. L. Turcotte. 1998. Forest fires: An example of self-organized critical behavior. *Science* 281:1840–1842.
- Naiman, R. J., C. A. Johnston, and J. C. Kelley. 1988. Alteration of North American Streams by Beaver. *BioScience* 38:753–762.
- Pan, Y., J. M. Chen, R. Birdsey, K. McCullough, L. He, and F. Deng. 2011. Age structure and disturbance legacy of North American forests.
- Reed, W. J., and K. S. McKelvey. 2002. Power-law behaviour and parametric models for the size-distribution of forest fires. *Ecological Modelling* 150:239–254.
- Reinhardt, E. D., and N. L. Crookston. 2015. The fire and fuels extension to the forest vegetation simulator. U.S. Department of Agriculture, Forest Service.
- Russell, E. W. B. 1997. *People and the land through time: Linking ecology and history*. Yale University Press, New Haven, Conn.
- Speth, J. D., and K. A. Spielmann. 1983. Energy source, protein metabolism, and hunter-gatherer subsistence strategies. *Journal of Anthropological Archaeology* 2:1–31.
- Stephens, S. L. 2005. Forest fire causes and extent on United States Forest Service lands. *International Journal of Wildland Fire* 14:213–222.
- Yang, J., H. S. He, and S. R. Shifley. 2008. Spatial Controls of Occurrence and Spread of Wildfires in the Missouri Ozark Highlands on JSTOR. *Ecological Applications* 18:1212–1225.