

Modeling the Historical Disturbance Regimes of NYC

Historical Disturbance Regimes

scale and frequency (Craig G. Lorimer 1980; Craig G Lorimer and White 2003; Foster, Motzkin, and Slater 1998, Pan et al. (2011))

human influence (Day 1953; E. W. B. Russell 1997; E. W. B. Russell 1997; Krech III 1999, Patterson and Sassaman (1988))

importance of lightning caused forest fires (Loope and Anderton 1998)

Fire

Custom Fuel Models

Hardwood forest types are based on model F9 (Scott and Burgan 2005; Anderson 1982). Where average loads for 1 hr, 10 hr, and 100 hr fuels were available for northeastern forest types the values were substituted into the custom fuel model (Reinhardt and Crookston 2015). The remaining fuel attributes were carried over from the base model. The USFS community types were cross-walked to the NY State Heritage community classification so fuels could be assigned for our landscape.

[cross walk table]

Fuel Moisture

Region specific fuel moistures were used as start conditions in FARSITE (Reinhardt and Crookston 2015). All freshwater wetland communities (marshes, shrub-swamps and forested wetlands) were initialized with a wet fuel profile, all other communities with burnable fuel types were initialized with a moist fuel profile.

Size Class	Very Dry	Dry	Moist	Wet
1 hr	5	7	10	19
10 hr	8	9	13	29
100 hr	12	14	17	22
Live woody	89	105	135	140
Liver herbaceous	60	82	116	120

Initial Conditions - Forest Age

Forest communities were initialized with an age of 65 (Pan et al. 2011; Loewenstein, Johnson, and Garrett 2000). All other community types were initialized with a forest age of 0




Figure 1: age class distribution

Initial Conditions - Canopy

Communities were initialized with their maximum canopy values as defined in the community table.

fire size and frequency literature

effects of fires on temperate forests (Kozlowski 2012) power law(Reed and McKelvey 2002; Cui and Perera 2008; Stephens 2005)

Modeling Expected Frequencies

A Poisson distribution is used to model annual forest fire events (E. A. Johnson 2001; Yang, He, and Shifley 2008). We created distributions for trail fires, garden fires and lightning fires.

Lightning Frequencies

The expected frequency (λ) of lightning caused fires are based on areal frequencies from region 9 (U.S. Northeast) USFS wildfire records between 1940 and 2000 (Stephens 2005). These values were converted from the given units (frequency/400000 ha)/yr to (frequency/km²)/yr.

Region	Lightning		Human	
	Ha burned	No. fires	Ha burned	No. fires
9	11.840	2.170	327.990	39.950

	Lightning		Human	
	Area burned	No. Fires	Area burned	No. Fires
1 hectare	0.0000296	0.000005425	0.000819975	0.000099875
per km ²	0.00296	0.0005425	0.0819975	0.0099875

Human Fire Frequency Scenarios

The extent and effect of human caused fires on the landscape prior to European settlement is debated (Day 1953; E. Russell 1983; Patterson and Sassaman 1988). We have proposed two frequency scenarios:

source	no human fire	Russell (1983)	Day (1953)
trail	0	0.00222	0.01778
garden	0	0.00028	0.00222

Critical Rainfall

In the model, fire spread was stopped when it encountered one of the following conditions: (1) a non-flammable type of land cover; (2) boundaries of the region; and (3) when rainfall exceeded a certain critical amount. By assuming that a daily precipitation of 30 mm or more would stop a fire, the R Crit in Eq. (2) was estimated as 0.026 (the proportion of total number of days that has daily

precipitation of 30 mm or more) from the historical precipitation data of the Edison weather station. (Li 2000)

Tree Allometry

Description	Equation	Reference
Tree Height	$TH = 44 * \ln(Age) - 93$	(Bean and Sanderson 2008)
DBH	$DBH = (Age - 34.44)/1.18$	(Loewenstein, Johnson, and Garrett 2000)
Crown Ratio	$CR = 0.4$	(Bean and Sanderson 2008)
Bark Thickness	$BT = vsp * DBH$	(Reinhardt and Crookston 2015)

Communities to Bark Thickness

Bark thickness multipliers for each community are based on the dominant tree type (Edinger et al. 2014), for communities with co-dominate species we calculated average bark thickness (Reinhardt and Crookston 2015).

community	dominant tree species	vsp scaler
Floodplain forest	avg(sliver maple, sycamore, American elm)	0.032
Red Maple Hardwood Swamp	red maple	0.028
Coastal Plain Atlantic Cedar Swamp	Atlantic cedar	0.025
Pitch pine - scrub oak barrens	avg(pitch pine, oak spp)	0.045
Chestnut oak forest	avg(American chestnut, oak spp)	0.043
Coastal oak beech forest	avg(oak spp, beech)	0.035
Coastal oak hickory forest	avg(oak spp, hickory spp)	0.045
Oak tulip forest	avg(oak spp, yellow-poplar)	0.038
Appalachian oak pine forest	avg(oak spp, pine spp)	0.038
Hemlock northern hardwood forest	hemlock	0.045
Inland Atlantic Cedar Swamp	Atlantic white cedar	0.025
Red maple black gum swamp	avg(red maple, black gum)	0.034
Red maple sweetgum swamp	avg(red maple, sweetgum)	0.032
Maritime holly forest	holly	0.042
Post oak black jack oak barrens	post oak	0.044
Appalachian oak hickory forest	avg(oak spp, hickory spp)	0.045
Beech maple mesic forest	avg(beech, sugar maple)	0.029
Successional maritime hardwoods	other hardwoods	0.044
Successional hardwood forest	other hardwoods	0.044

Fire Mortality Equations

Scorch Height

[1]

$$SH = 3.1817(FL^{1.4503})$$

(Bean and Sanderson 2008)

Crown Kill

[2]

$$CK = 41.961(100(\ln(\frac{SH - CH}{CL})) - 89.721$$

(Bean and Sanderson 2008)

Percent Mortality

[3]

$$P_m = \frac{1.0}{1.0 + e^{-1.941 + 6.316(1.0 - e^{-BT}) - 0.000535CK^2}}$$

(Bean and Sanderson 2008)

Horticulture

Site Selection

Gardens were modeled at historical sites meeting the following two criteria.

1. historical and archealogical records provide evidence for semi-permanent habitation.
2. site point is located within 250 m of freshwater source

Garden loacations are then selected using a suitability model which takes into account proximity to the site center, slope and community type.

Archaeological Evidence for Gardening

(Kraft 2001, Cantwell and Wall (2001), Benison (1997))

Ethnohistorical

(Ascher 1860, Danckaerts and Sluyter (1867))

crop	yield (kg/ha)	calories/100 g	calories/kg	calories/ha
corn	1720	365	3650	6278000
beans (Phaseolus vulgaris)	110	33	330	36300
squash (Indian squash)	80	16	160	12800

corn-bean-squash ploy-culture yields in Tabasco, Mexico (Gliessman, Engles, and Krieger 1998) pg 224 table 15.3

Caloric density [USDA Basic Reports Agricultural Research Services]

Agricultural Dependency Scenarios

caloric requirements (Speth and Spielmann 1983)

Dependency	calories/person/yr	ha/person	m ² /person
15%	94060.50	0.01	14.87
30%	188121.00	0.03	29.73
60%	376242.00	0.06	59.47
100%	627070.00	0.10	99.11

Beaver

Succession

The freshwater wetland succession sequence has four stages(Allen 1983; Hay 2010; Johnston and Naiman 1990; Logofet et al. 2016; Naiman, Johnston, and Kelley 1988; “Ecology of Red Maple Swamps in the Glaciated Northeast: A Community Profile” 1993).

active beaver pond → emergent marsh → shrub swamp → forested wetland

Conversion from non-wetland community to active pond can occur along any perennial streams where the gradient of the stream is less than 15 degrees. Due to the temporal scale of our study (200 yrs), forested wetlands are treated as a terminal community in this series (“Ecology of Red Maple Swamps in the Glaciated Northeast: A Community Profile” 1993). This rule defines beaver caused disturbance as a unidirectional change in successional trajectory. A non-wetland communities can be converted into a wetland type but this conversion cannot be reversed.

Model Parameters

Parameter	Value	Source
abandonment P	0.10	(Logofet et al. 2016)

Parameter	Value	Source
colony density	0.4 colonies/km ²	(Naiman, Johnston, and Kelley 1988)
territory	1000 m	Naiman, Johnston, and Kelley (1988); Allen (1983)]

Analysis

time since fire (E. A. Johnson 2001) fire size frequency hist (Reed and McKelvey 2002; Cui and Perera 2008; Malamud, Morein, and Turcotte 1998)

References

- Allen, Arthur W. 1983. *Habitat Suitability Index Models: Beaver*. Western Energy; Land Use Team, Division of Biological Service, Research; Development, Fish; Wildlife Service, US Department of the Interior. http://breb.ro/Publicatii/hab_suitability_index_models_beaver.pdf.
- Anderson, Hal E. 1982. "Aids to Determining Fuel Models for Estimating Fire Behavior." *The Bark Beetles, Fuels, and Fire Bibliography*, 143. <http://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=1147&context=barkbeetles>.
- Ascher, G.M. 1860. *Henry Hudson the Navigator*. 27. London: Hakluyt Society.
- Bean, William T., and Eric W. Sanderson. 2008. "Using a Spatially Explicit Ecological Model to Test Scenarios of Fire Use by Native Americans: An Example from the Harlem Plains, New York, NY." *Ecological Modelling* 211: 301–8.
- Benison, Chris. 1997. "Horticulture and the Maintenance of Social Complexity in Late Woodland Southeastern New England." *North American Archaeologist* 18 (1): 1–17. doi:10.2190/0KB7-JM4H-PJ8U-VN2L.
- Cantwell, Anne-Marie, and Diana diZerega Wall. 2001. *Unearthing Gotham: The Archaeology of New York City*. New Haven, CT: Yale University Press.
- Cui, Wenbin, and Ajith H. Perera. 2008. "What Do We Know About Forest Fire Size Distribution, and Why Is This Knowledge Useful for Forest Management?" *International Journal of Wildland Fire* 17 (2): 234. doi:10.1071/WF06145.
- Danckaerts, Jasper, and Peter Sluyter. 1867. *Journal of a Voyage to New York: And a Tour in Several of the American Colonies in 1679-80*. Brooklyn: The Long Island Historical Society.
- Day, Gordon M. 1953. "The Indian as an Ecological Factor in the Northeastern Forest." *Ecology* 34 (2): 329–46.
- "Ecology of Red Maple Swamps in the Glaciated Northeast: A Community

- Profile.” 1993. Biological Report 12. U.S. Fish; Wildlife Service. <http://pubs.usgs.gov/unnumbered/70110597/report.pdf>.
- Foster, David R., Glenn Motzkin, and Benjamin Slater. 1998. “Land-Use History as Long-Term Broad-Scale Disturbance: Regional Forest Dynamics in Central New England.” *Ecosystems* 1 (1): 96–119. doi:10.1007/s100219900008.
- Gliessman, Stephen R., Eric Engles, and Robin Krieger. 1998. *Agroecology: Ecological Processes in Sustainable Agriculture*. CRC Press.
- Hay, Keith G. 2010. “Succession of Beaver Ponds in Colorado 50 Years After Beaver Removal.” *The Journal of Wildlife Management* 74 (8): 1732–6. doi:10.2193/2009-055.
- Johnson, Edward A. 2001. *Forest Fires: Behavior and Ecological Effects*. Academic Press.
- Johnston, Carol A., and Robert J. Naiman. 1990. “The Use of a Geographic Information System to Analyze Long-Term Landscape Alteration by Beaver.” *Landscape Ecology* 4 (1): 5–19. doi:10.1007/BF02573947.
- Kozlowski, T. T. 2012. *Fire and Ecosystems*. Elsevier.
- Kraft, Herbert C. 2001. *The Lenape-Delaware Indian Heritage: 10,000 B.C.-A.D. 2000*. [S.l.]: Lenape Books.
- Krech III, Shepard. 1999. *The Ecological Indian*. New York: W.W. Norton & Company.
- Li, Chao. 2000. “Reconstruction of Natural Fire Regimes Through Ecological Modelling.” *Ecological Modelling* 134 (2): 129–44. <http://www.sciencedirect.com/science/article/pii/S0304380000002908>.
- Loewenstein, Edward F., Paul S. Johnson, and Harold E. Garrett. 2000. “Age and Diameter Structure of a Managed Uneven-Aged Oak Forest.” *Canadian Journal of Forest Research* 30 (7): 1060–70. <http://www.nrcresearchpress.com/doi/abs/10.1139/x00-036>.
- Logofet, D. O., O. I. Evstigneev, A. A. Aleinikov, and A. O. Morozova. 2016. “Succession Caused by Beaver (Castor Fiber L.) Life Activity: II. a Refined Markov Model.” *Biology Bulletin Reviews* 6 (1): 39–56. doi:10.1134/S2079086416010047.
- Loope, Walter L., and John B. Anderton. 1998. “Human Vs. Lightning Ignition of Presettlement Surface Fires in Coastal Pine Forests of the Upper Great Lakes.” *The American Midland Naturalist* 140 (2): 206–18. doi:10.1674/0003-0031(1998)140[0206:HVLIOF]2.0.CO;2.
- Lorimer, Craig G, and Alan S White. 2003. “Scale and Frequency of Natural Disturbances in the Northeastern US: Implications for Early Successional Forest Habitats and Regional Age Distributions.” *Forest Ecology and Management, Early-Successional Forests and Shrubland Habitats in the North Eastern*

United States :Critical Habitats dependent on Disturbance, 185 (1–2): 41–64. doi:10.1016/S0378-1127(03)00245-7.

Lorimer, Craig G. 1980. “Age Structure and Disturbance History of a Southern Appalachian Virgin Forest.” *Ecology* 61 (5): 1169–84. doi:10.2307/1936836.

Malamud, Bruce D., Gleb Morein, and Donald L. Turcotte. 1998. “Forest Fires: An Example of Self-Organized Critical Behavior.” *Science* 281 (5384): 1840–2. <http://science.sciencemag.org/content/281/5384/1840.short>.

Naiman, Robert J., Carol A. Johnston, and James C. Kelley. 1988. “Alteration of North American Streams by Beaver.” *BioScience* 38 (11): 753–62. doi:10.2307/1310784.

Pan, Yude, Jing M. Chen, Richard Birdsey, Kevin McCullough, Liming He, and Feng Deng. 2011. “Age Structure and Disturbance Legacy of North American Forests.” <http://www.treesearch.fs.fed.us/pubs/37680>.

Patterson, William A., and Kenneth E. Sassaman. 1988. “Indian Fires in the Prehistory of New England.” In *Holocene Human Ecology in Northeastern North America*, edited by George P. Nicholas, 107–35. New York: Plenum Press.

Reed, William J., and Kevin S. McKelvey. 2002. “Power-Law Behaviour and Parametric Models for the Size-Distribution of Forest Fires.” *Ecological Modelling* 150 (3): 239–54. doi:10.1016/S0304-3800(01)00483-5.

Reinhardt, Elizabeth D., and Nicholas L. Crookston. 2015. “The Fire and Fuels Extension to the Forest Vegetation Simulator.” U.S. Department of Agriculture, Forest Service. <http://citeserx.ist.psu.edu/viewdoc/download?doi=10.1.1.231.6561&rep=rep1&type=pdf>.

Russell, Emily Wyndham Barnett. 1997. *People and the Land Through Time: Linking Ecology and History*. New Haven, Conn.: Yale University Press.

Russell, EWB. 1983. “Indian-Set Fires in the Forests of the Northeastern United States.” *Ecology* 64: 78–88.

Scott, Joe H., and Robert E. ; Burgan. 2005. *Standard Fire Behavior Fuel Models: A Comprehensive Set for Use with Rothermel’s Surface Fire Spread Model*. <http://www.treesearch.fs.fed.us/pubs/9521>.

Speth, John D, and Katherine A Spielmann. 1983. “Energy Source, Protein Metabolism, and Hunter-Gatherer Subsistence Strategies.” *Journal of Anthropological Archaeology* 2 (1): 1–31. doi:10.1016/0278-4165(83)90006-5.

Stephens, Scott L. 2005. “Forest Fire Causes and Extent on United States Forest Service Lands.” *International Journal of Wildland Fire* 14: 213–22. <http://nature.berkeley.edu/stephenslab/wp-content/uploads/2015/04/Stephens-USFS-fire-stats-IJWF-05.pdf>.

Yang, Jian, Hong S. He, and Stephan R. Shifley. 2008. “Spatial Controls of Occurrence and Spread of Wildfires in the Missouri Ozark Highlands on JSTOR.”

Ecological Applications 18 (5): 1212–25. http://www.jstor.org/stable/40062223?seq=1#page_scan_tab_contents.