

# Perfect Epidemics

## Applied Probability Seminar

Department of Statistics, University of Warwick

W S Kendall   S B Connor

Warwick, York


21 January 2025



# Introduction

“Maybe the only significant difference between a really smart simulation and a human being was the noise they made when you punched them.”

([The Long Earth](#), Pratchett & Baxter, 2012)



Handout is on the web: use the QR-code or visit  
[wilfridskendall.github.io/talks/PerfectEpidemics](https://wilfridskendall.github.io/talks/PerfectEpidemics).

This is initial work on using perfect simulation (CFTP) for epidemics.  
WSK acknowledges the support of UK EPSRC grant EP/R022100.

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!
- Key ideas of “*classic CFTP*”:

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!
- Key ideas of “*classic CFTP*”:
  - ▶ extend simulation *backwards* through time not forwards;

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!
- Key ideas of “*classic CFTP*”:
  - ▶ extend simulation *backwards* through time not forwards;
  - ▶ exploit monotonicity (*couple* maximal and minimal processes);

# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!
- Key ideas of “*classic CFTP*”:
  - ▶ extend simulation *backwards* through time not forwards;
  - ▶ exploit monotonicity (*couple* maximal and minimal processes);
  - ▶ seek coalescence.



# 1: Introduction to Perfect Simulation

- Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- The term “perfect simulation” (WSK, 1998) should make you suspicious: perfection is never achieved. This is why the term was chosen!
- Key ideas of “*classic CFTP*”:
  - ▶ extend simulation *backwards* through time not forwards;
  - ▶ exploit monotonicity (*couple* maximal and minimal processes);
  - ▶ seek coalescence.
- Simplest possible example: *random-walk-CFTP*  
(can boost to use Ising model to do simple image reconstruction).

## Classic CFTP for a simple random walk (I)

- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .

## Classic CFTP for a simple random walk (I)

- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**

## Classic CFTP for a simple random walk (I)


- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace  $+1$  jump by staying still, **and**

## Classic CFTP for a simple random walk (I)

- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace  $+1$  jump by staying still, **and**
  - ▶ at state 0 replace  $-1$  jump by staying still.


## Classic CFTP for a simple random walk (I)

- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace  $+1$  jump by staying still, **and**
  - ▶ at state 0 replace  $-1$  jump by staying still.
- Conventional MCMC picks a starting point, then runs the simple random walk for long time till approximate equilibrium.



## Classic CFTP for a simple random walk (I)

- Consider a simple random walk on  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace  $+1$  jump by staying still, **and**
  - ▶ at state 0 replace  $-1$  jump by staying still.
- Conventional MCMC picks a starting point, then runs the simple random walk for long time till approximate equilibrium.
- How long? One way to *estimate* this is to run two (or several?) coupled copies till they meet. If probability of meeting by time  $T$  is high, then deviation of  $X_T$  from equilibrium is statistically small;








## Classic CFTP for a simple random walk (I)

- So now start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.




## Classic CFTP for a simple random walk (I)

- So now start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.
- If not coupled by time 0, than back-off to time  $-2T$  and repeat.




## Classic CFTP for a simple random walk (I)

- So now start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.
- If not coupled by time 0, than back-off to time  $-2T$  and repeat.
- May need to iterate back-off doubling several times.




## Classic CFTP for a simple random walk (I)

- So now start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.
- If not coupled by time 0, than back-off to time  $-2T$  and repeat.
- May need to iterate back-off doubling several times.
- When coupled, top and bottom yield a common value at time 0.



## Classic CFTP for a simple random walk (I)

- So now start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.
- If not coupled by time 0, then back-off to time  $-2T$  and repeat.
- May need to iterate back-off doubling several times.
- When coupled, top and bottom yield a common value at time 0.
- The common value is an exact draw from equilibrium!



## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).



## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that
  - ▶ *Classical CFTP* equivalent to uniform ergodicity (Foss & Tweedie, 1998);

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that
  - ▶ *Classical CFTP* equivalent to uniform ergodicity (Foss & Tweedie, 1998);
  - ▶ *Dominated CFTP* achievable under geometric ergodicity (WSK, 2004);

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that
  - ▶ *Classical CFTP* equivalent to uniform ergodicity (Foss & Tweedie, 1998);
  - ▶ *Dominated CFTP* achievable under geometric ergodicity (WSK, 2004);
  - ▶ Dominated CFTP can work in some **non**-geometrically ergodicity cases (SBC & WSK, 2007a; *nb* corrigendum SBC & WSK, 2007b).

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that
  - ▶ *Classical CFTP* equivalent to uniform ergodicity (Foss & Tweedie, 1998);
  - ▶ *Dominated CFTP* achievable under geometric ergodicity (WSK, 2004);
  - ▶ Dominated CFTP can work in some **non**-geometrically ergodicity cases (SBC & WSK, 2007a; *nb* corrigendum SBC & WSK, 2007b).
- *Dominated CFTP* can be used to carry out perfect simulation for stable point processes (WSK & Møller, 2000);

## 2: A short section on some theory about CFTP

- What about cases where monotonicity fails? or there isn't a sensible “maximal” process? Ideas (WSK, 1998):
  - ▶ cross-couple upper and lower envelope processes;
  - ▶ dominate by amenable “dominating process” (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle* we can show that
  - ▶ *Classical CFTP* equivalent to uniform ergodicity (Foss & Tweedie, 1998);
  - ▶ *Dominated CFTP* achievable under geometric ergodicity (WSK, 2004);
  - ▶ Dominated CFTP can work in some **non**-geometrically ergodicity cases (SBC & WSK, 2007a; *nb* corrigendum SBC & WSK, 2007b).
- *Dominated CFTP* can be used to carry out perfect simulation for stable point processes (WSK & Møller, 2000);
- Detailed expositions: WSK (2005), Huber (2015).  
(Want to implement CFTP in R? see WSK, 2015.)

### 3: Perfect Epidemics: a challenge problem for CFTP

*Many* important inferential questions (Cori & Kucharski, 2024).

Simplest models (versus UK model with  $10^6$  agents!, Fraser & Others, 2023):

**S-I-R deterministic epidemic:** susceptibles  $s$ , infectives  $i$ , removals  $r$   
(constant total population  $s + i + r = n$ ):

$$\begin{aligned}s' &= -\alpha s i, \\ i' &= (\alpha s - \beta) i, \\ r' &= \beta i.\end{aligned}$$

### 3: Perfect Epidemics: a challenge problem for CFTP

Many important inferential questions (Cori & Kucharski, 2024).

Simplest models (versus UK model with  $10^6$  agents!, Fraser & Others, 2023):

**S-I-R deterministic epidemic:** susceptibles  $s$ , infectives  $i$ , removals  $r$   
(constant total population  $s + i + r = n$ ):

$$\begin{aligned}s' &= -\alpha s i, \\ i' &= (\alpha s - \beta) i, \\ r' &= \beta i.\end{aligned}$$

**S-I-R stochastic epidemic:** a Markov chain  $(S, I, R)$  with transitions

**Infection:**  $S \rightarrow S - 1, \quad I \rightarrow I + 1$  at rate  $\alpha S I$ ,

**Removal:**  $I \rightarrow I - 1, \quad R \rightarrow R + 1$  at rate  $\beta I$ .



### 3: Perfect Epidemics: a challenge problem for CFTP

Many important inferential questions (Cori & Kucharski, 2024).

Simplest models (versus UK model with  $10^6$  agents!, Fraser & Others, 2023):

**S-I-R deterministic epidemic:** susceptibles  $s$ , infectives  $i$ , removals  $r$  (constant total population  $s + i + r = n$ ):

$$\begin{aligned}s' &= -\alpha s i, \\ i' &= (\alpha s - \beta) i, \\ r' &= \beta i.\end{aligned}$$

**S-I-R stochastic epidemic:** a Markov chain  $(S, I, R)$  with transitions

**Infection:**  $S \rightarrow S - 1, \quad I \rightarrow I + 1$  at rate  $\alpha S I$ ,

**Removal:**  $I \rightarrow I - 1, \quad R \rightarrow R + 1$  at rate  $\beta I$ .

Both models make an unrealistic assumption: homogeneous mixing.

# The first question asked about a new epidemic

“What is the R-number?”

# The first question asked about a new epidemic

“What is the R-number?”

The R-number is  $\alpha s_0/\beta$ : mean number of new infectives produced per infective at *start* of epidemic with initially  $s_0$  susceptibles.

Whittle (1955)’s threshold theorem: R-number  $\gg 1$  means positive chance of epidemic infecting significant proportion of the population.

# The first question asked about a new epidemic

“What is the R-number?”

The R-number is  $\alpha s_0/\beta$ : mean number of new infectives produced per infective at *start* of epidemic with initially  $s_0$  susceptibles.

Whittle (1955)’s threshold theorem: R-number  $\gg 1$  means positive chance of epidemic infecting significant proportion of the population.

*Wikipedia*: “The British-registered *Diamond Princess* was the first cruise ship to have a major [COVID-19] outbreak on board, with the ship quarantined at Yokohama from 4 February 2020 for about a month. Of 3711 passengers and crew, around 700 people became infected and 9 people died.”

Evidently  $\alpha s_0/\beta \gg 1$  – as was sadly later confirmed, a sorrow for us all.



## 4: Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- ① Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;

## 4: Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- 1 Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- 2 Inference is **really tough**: hard to get information about infection times;

## 4: Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- 1 Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- 2 Inference is **really tough**: hard to get information about infection times;
- 3 It is all **especially tough** in early stages. Answers are most needed when hardly any information is available (a simplified example for a Warwick UG second-year statistics module shows how tough this can be);

## 4: Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- 1 Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- 2 Inference is **really tough**: hard to get information about infection times;
- 3 It is all **especially tough** in early stages. Answers are most needed when hardly any information is available (a simplified example for a Warwick UG second-year statistics module shows how tough this can be);
- 4 Markov chain Monte Carlo (MCMC) can be used (see next slide) but what about burn-in?



## 4: Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- 1 Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- 2 Inference is **really tough**: hard to get information about infection times;
- 3 It is all **especially tough** in early stages. Answers are most needed when hardly any information is available (a simplified example for a Warwick UG second-year statistics module shows how tough this can be);
- 4 Markov chain Monte Carlo (MCMC) can be used (see next slide) but what about burn-in?
- 5 Can we use **perfect simulation**?

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”

(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”

(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- 1 Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O’Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- 1 Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- 2 Visualize  $n$  timelines, along which incidents are scattered:

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- ➊ Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- ➋ Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- ➊ Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- ➋ Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- ➊ Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- ➋ Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.
- ➌ Poisson point processes *of appropriate rates* yield an S-I-R epidemic.

## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- ➊ Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- ➋ Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.
- ➌ Poisson point processes *of appropriate rates* yield an S-I-R epidemic.
- ➍ First step: evolve whole S-I-R trajectory in *algorithmic time* (alter potential infections and removals using immigration-death in discrete algorithmic time).



## 5: An easier question

An absurdly simple variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O'Neill & Roberts, 1999; Gibson & Renshaw, 2001)

An important step on the way: generating an *unconditioned* epidemic.

- ➊ Thus  $n$ ,  $\alpha$ ,  $\beta$  are known, removal times are observed, but unobserved infection times **must be inferred**.
- ➋ Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.
- ➌ Poisson point processes *of appropriate rates* yield an S-I-R epidemic.
- ➍ First step: evolve whole S-I-R trajectory in *algorithmic time* (alter potential infections and removals using immigration-death in discrete algorithmic time).
- ➎ Result: *trajectory-valued chain*, unconditioned S-I-R as equilibrium.

# From incidents to unconditioned epidemic trajectories (1/3)




Figure 1: Light-orange circles denote potential infections (arrows point upwards to targets); light-brown circles denote potential removals.

# From incidents to unconditioned epidemic trajectories (2/3)




Figure 2: Activate (a) *infection* if target on lowest uninfected timeline; (b) *removal* if in infected region, then remove lowest infected (purple disk if different timeline).

# From incidents to unconditioned epidemic trajectories (3/3)




Figure 3: A step in algorithmic time for the unconditioned epidemic simply involves replacing the original incidents by a new set of incidents.

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;



## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;
  - 3 Replace infections ( $I$ s).

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-)*time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;
  - 3 Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .

# Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;
  - 3 Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - 1 For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-)*time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;
  - 3 Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - 1 For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;
  - 2 For  $\tau = (2n+1)T$ , resample timelines (not times) of  $R$ s;

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - ➊ Replace removals ( $R$ s);
  - ➋ Re-sample timelines (though not times) of  $R$ s;
  - ➌ Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - ➊ For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;
  - ➋ For  $\tau = (2n+1)T$ , resample timelines (not times) of  $R$ s;
  - ➌ For  $(2n+1)T < \tau < (2n+2)T$ , update old  $I$ s in  $((2n+2)T - \tau, T)$ .

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - ➊ Replace removals ( $R$ s);
  - ➋ Re-sample timelines (though not times) of  $R$ s;
  - ➌ Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - ➊ For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;
  - ➋ For  $\tau = (2n+1)T$ , resample timelines (not times) of  $R$ s;
  - ➌ For  $(2n+1)T < \tau < (2n+2)T$ , update old  $I$ s in  $((2n+2)T - \tau, T)$ .
- Thus the original update is expressible as a (continuous) composition of updates, each of which satisfies detailed balance in equilibrium.

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-)*time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $R$ s);
  - 2 Re-sample timelines (though not times) of  $R$ s;
  - 3 Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - 1 For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;
  - 2 For  $\tau = (2n+1)T$ , resample timelines (not times) of  $R$ s;
  - 3 For  $(2n+1)T < \tau < (2n+2)T$ , update old  $I$ s in  $((2n+2)T - \tau, T)$ .
- Thus the original update is expressible as a (continuous) composition of updates, each of which satisfies detailed balance in equilibrium.
- The connection “restriction=conditioning” is thereby preserved.

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-) *time-reversible*: so restriction to subset  $S$  of state-space (of *activated* removals occurring precisely at specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - ➊ Replace removals ( $R$ s);
  - ➋ Re-sample timelines (though not times) of  $R$ s;
  - ➌ Replace infections ( $I$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - ➊ For  $2nT < \tau < (2n+1)T$ , update old  $R$ s with times in  $(0, \tau - 2nT)$ ;
  - ➋ For  $\tau = (2n+1)T$ , resample timelines (not times) of  $R$ s;
  - ➌ For  $(2n+1)T < \tau < (2n+2)T$ , update old  $I$ s in  $((2n+2)T - \tau, T)$ .
- Thus the original update is expressible as a (continuous) composition of updates, each of which satisfies detailed balance in equilibrium.
- The connection “restriction=conditioning” is thereby preserved.
- Crucially, step 2 ensures composition action is irreducible over  $S$ !  
(So equilibrium under conditioning is unique.)



# Illustration of technical point (1/8)




Figure 4: No change yet to removals or infections;

# Illustration of technical point (2/8)




Figure 5: Replace first third of removals, infections unchanged;

# Illustration of technical point (3/8)




Figure 6: Replace first two-thirds of removals, infections unchanged;

# Illustration of technical point (4/8)




Figure 7: All removals resampled, infections as yet unchanged;

# Illustration of technical point (5/8)




Figure 8: Re-sample all removal timelines, infections as yet unchanged;

# Illustration of technical point (6/8)




Figure 9: Re-sample last third of infections;

# Illustration of technical point (7/8)




Figure 10: Re-sample last two-thirds of infections;

# Illustration of technical point (8/8)




Figure 11: All infections now re-sampled.



# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).

Consequently:

- ▶ conditioned removals must be able to change timeline (but not time of occurrence);

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

Implications:

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;
- ▶ a conditioned removal timeline can be altered only if it doesn't de-activate;

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;
- ▶ a conditioned removal timeline can be altered only if it doesn't de-activate;
- ▶ an infection cannot be removed if that action loses a conditioned removal;

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;
- ▶ a conditioned removal timeline can be altered only if it doesn't de-activate;
- ▶ an infection cannot be removed if that action loses a conditioned removal;
- ▶ an infection can be introduced only if no new observed removals result.



# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;
  - ▶ a conditioned removal timeline can be altered only if it doesn't de-activate;
  - ▶ an infection cannot be removed if that action loses a conditioned removal;
  - ▶ an infection can be introduced only if no new observed removals result.
- Does this produce a *feasible* and suitably monotonic algorithm?

# Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point).  
Consequently:
  - ▶ conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.

## Implications:

- ▶ a removal can be introduced only if it doesn't activate;
- ▶ a conditioned removal timeline can be altered only if it doesn't de-activate;
- ▶ an infection cannot be removed if that action loses a conditioned removal;
- ▶ an infection can be introduced only if no new observed removals result.
- Does this produce a *feasible* and suitably monotonic algorithm?
- **Housekeeping details** required to establish that monotonicity still works.  
Key notions: *last feasible epidemic* (LFE) and *no-fly zone* (NFZ).

# Initial conditional epidemic




Figure 12: Initial epidemic with conditioned removals indicated using purple circles (and purple disks when different timelines are infected).

# Conditional epidemic update




Figure 13: Epidemic updated under restriction: all conditioned removals remain activated, no new removals are activated. Green infections have been “perpetuated”.

# Last feasible epidemic (LFE)




Figure 14: LFE computed recursively working right-to-left: the slowest sequence of infections deals with all infected timelines in order (includes perpetuated infections).




Figure 15: NFZ computed recursively working right-to-left: it traces the region of timelines that must not be infected if one is not to activate unobserved removals.

## Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).

## Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of **Bailey, 1975**).
- **Assume**



## Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;

## Example


- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;

## Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of **Bailey, 1975**).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).


# Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson \*et al.\*, 2017](#)), we can construct a **perfect simulation GIF** yielding a draw from unobserved pattern of infections.




# Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson \*et al.\*, 2017](#)), we can construct a **perfect simulation GIF** yielding a draw from unobserved pattern of infections.




# Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson \*et al.\*, 2017](#)), we can construct a **perfect simulation GIF** yielding a draw from unobserved pattern of infections.




# Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson \*et al.\*, 2017](#)), we can construct a **perfect simulation GIF** yielding a draw from unobserved pattern of infections.



# Example

- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson \*et al.\*, 2017](#)), we can construct a **perfect simulation GIF** yielding a draw from unobserved pattern of infections.





# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (next steps after **SBC & WSK, 2024**)

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (**next steps after SBC & WSK, 2024**)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (**next steps after SBC & WSK, 2024**)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;
  - ▶ construct steepest ascent algorithm (in effect, variant of Robbins-Monro stochastic optimization) to find *maximum a posteriori* estimates of  $\alpha$  and  $\beta$ ;

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (next steps after **SBC & WSK, 2024**)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;
  - ▶ construct steepest ascent algorithm (in effect, variant of Robbins-Monro stochastic optimization) to find *maximum a posteriori* estimates of  $\alpha$  and  $\beta$ ;
  - ▶ or even, with some computational effort, compute the entire posterior joint density for  $\alpha$  and  $\beta$ !

# So what?

- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (next steps after **SBC & WSK, 2024**)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;
  - ▶ construct steepest ascent algorithm (in effect, variant of Robbins-Monro stochastic optimization) to find *maximum a posteriori* estimates of  $\alpha$  and  $\beta$ ;
  - ▶ or even, with some computational effort, compute the entire posterior joint density for  $\alpha$  and  $\beta$ !
- Finally: generalize to other suitable compartment models?

# Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!





# Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.



# Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.
- *Of course* detailed models resist perfect simulation (but it will be helpful to compare with a simpler model using fewer parameters).



# Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.
- *Of course* detailed models resist perfect simulation (but it will be helpful to compare with a simpler model using fewer parameters).
- Still to be done: statistical estimation of parameters, generalization to other compartment models.



# Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.
- *Of course* detailed models resist perfect simulation (but it will be helpful to compare with a simpler model using fewer parameters).
- Still to be done: statistical estimation of parameters, generalization to other compartment models.
- Thank you for your attention! **QUESTIONS?**



# References I

- Bailey, N.T.J. (1975) *The mathematical theory of infectious diseases and its applications*, 2nd Ed. ed. Griffin.
- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V.B. (2017) Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, **59**, 65–98.
- Cori, A. & Kucharski, A. (2024) Inference of epidemic dynamics in the COVID-19 era and beyond. *Epidemics*, **48**, 100784.
- Donnelly, P. & Kurtz, T.G. (1996) A countable representation of the Fleming-Viot measure-valued diffusion. *The Annals of Probability*, **24**, 698–742.
- Foss, S.G. & Tweedie, R.L. (1998) Perfect simulation and backward coupling. *Stochastic Models*, **14**, 187–203.
- Fraser, C. & Others (2023) OpenABM-Covid19: Agent-based model for modelling the Covid-19 and Contact-Tracing.
- Gibson, G.J. & Renshaw, E. (1998) Estimating parameters in stochastic compartmental models using Markov chain methods. *Mathematical and Medical Biology*, **15**, 19–40.
- Gibson, G.J. & Renshaw, E. (2001) Likelihood estimation for stochastic compartmental models using Markov chain methods. *Statistics and Computing*, **11**, 347–358.
- Huber, M.L. (2015) *Perfect Simulation*. Boca Raton: Chapman; Hall/CRC.
- Liggett, T.M. (1985) *Interacting particle systems*. Springer Verlag.

# References II

- O'Neill, P.D. & Roberts, G.O. (1999) Bayesian Inference for Partially Observed Stochastic Epidemics. *Journal of the Royal Statistical Society Series A: Statistics in Society*, **162**, 121–129.
- Pratchett, T. & Baxter, S. (2012) *The Long Earth*. Doubleday.
- Propp, J.G. & Wilson, D.B. (1996) Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structures and Algorithms*, **9**, 223–252.
- SBC (2020) Omnithermal Perfect Simulation for Multi-server Queues. *ACM Transactions on Modeling and Computer Simulation*, **30**, 1–15.
- SBC & WSK (2007b) Perfect simulation for a class of positive recurrent Markov chains (corrigendum). *Annals of Applied Probability*, **17**, 1808–1810.
- SBC & WSK (2007a) Perfect simulation for a class of positive recurrent Markov chains. *Annals of Applied Probability*, **17**, 781–808.
- SBC & WSK (2015) Perfect simulation of M/G/c queues. *Advances in Applied Probability*, **47**, 1039–1063.
- SBC & WSK (2024) Perfect Epidemics.
- Whittle, P. (1955) The outcome of a stochastic epidemic—a note on Bailey's paper. *Biometrika*, **42**, 116–122.
- WSK (1998) Perfect Simulation for the Area-Interaction Point Process. *Probability towards 2000* (Accardi, L. & Heyde, C.C. eds). Springer-Verlag, pp. 218–234.

# References III

- WSK (2004) Geometric ergodicity and perfect simulation. *Electronic Communications in Probability*, **9**, 140–151.
- WSK (2005) Notes on Perfect Simulation. Singapore: World Scientific, pp. 93–146.
- WSK (2015) Introduction to CFTP using R. *Stochastic geometry, spatial statistics and random fields, Lecture notes in mathematics*. Springer, pp. 405–439.
- WSK & Møller, J. (2000) Perfect simulation using dominating processes on ordered spaces, with application to locally stable point processes. *Advances in Applied Probability*, **32**, 844–865.

## Image information

Image	Attribution	
<i>Terry Pratchett</i> Classic CFTP for a simple random walk	Luigi Novi Result of code written by WSK	<i>CC BY 3.0</i>
<i>Diamond Princess</i> Epidemic CFTP images and animation	Alpsdake Result of code written by WSK	<i>CC BY-SA 4.0</i>

## Previous instances of this talk

Date	Title		Location
19/04/24	Perfect Epidemics	Short Research Talk (12min)	Warwick
15/05/24	McMC and Perfect Simulation	Graduate Seminar, Aristotle Univ. (50min)	Thessaloniki
17/01/25	Perfect Epidemics	Applied Probability Seminar (50min)	Warwick



# A “near-maximal” configuration




Figure 16: A conditional epidemic in which all activated infections occur before time 3.0, also before smallest observed removal time.

# Other technical information

## Software versions

Software used in computations:

<i>Software</i>	<i>Version</i>	<i>Branch</i>	<i>Date of last commit</i>
Quarto	1.6.39	—	
Running under julia	1.11.2	—	
Module EpidemicsCFTP	2.2.488	main	Mon Jan 20 15:38:58 2025
Module EpidemicsUtilities	0.1.2.154	main	Wed Jan 15 13:23:36 2025
This Quarto script	2.2.593	master	Mon Jan 20 17:17:03 2025

## Revision history

These notes were produced from PerfectEpidemics.qmd:

---

Author:	Wilfrid Kendall <a href="mailto:W.S.Kendall@warwick.ac.uk">W.S.Kendall@warwick.ac.uk</a>
Date:	Mon Jan 20 17:17:03 2025 +0000
Summary:	Minor 2.2.593

---