

# Perfect Epidemics

## Seminar at University College Dublin

W S Kendall S B Connor

Warwick, York

20 October 2025



## Introduction

Homage to Dublin  
(Book of Kells, 9th century)



Work on perfect simulation (CFTP) for epidemics, now being written up.  
WSK acknowledges the support of UK EPSRC grant EP/R022100.



Handout is on the web: use the QR-code or visit  
[wilfridskendall.github.io/talks/PerfectEpidemics](https://wilfridskendall.github.io/talks/PerfectEpidemics).



W S Kendall, S B Connor (Warwick, York)

Perfect Epidemics

20 October 2025

1 / 30

W S Kendall, S B Connor (Warwick, York)

Perfect Epidemics

20 October 2025

2 / 30

## Plan of talk

Gregory: Is there any other point to which you would wish to draw my attention?  
Holmes: To the curious incident of the dog in the night-time.  
Gregory: The dog did nothing in the night-time.  
Holmes: That was the curious incident.

(“The Adventure of Silver Blaze”, Sir Arthur Conan Doyle, 1892)

- (a) Introduction to perfect simulation:
- (b) A little theory about CFTP;
- (c) Epidemics and the  $R$ -number;
- (d) “Contact tracing”: inferring infection pattern if removals observed;
- (e) Example with real data.



## Perfect Epidemics

2025-10-09

### Plan of talk

The Sherlock Holmes quote emphasizes how informatin can be supplied from what we *don't* see,  
This itemization summarizes the plan of the talk.  
Content culminates in part (d), describing the use of perfect simulation when inferring infection pattern if one has observed all epidemic removals.

#### Plan of talk

- Gregory: Is there any other point to which you would wish to draw my attention?  
Holmes: To the curious incident of the dog in the night-time.  
Gregory: The dog did nothing in the night-time.  
Holmes: That was the curious incident.
- (“The Adventure of Silver Blaze”, Sir Arthur Conan Doyle, 1892)
- (a) Introduction to perfect simulation;
  - (b) A little theory about CFTP;
  - (c) Epidemics and the  $R$ -number;
  - (d) “Contact tracing”: inferring infection pattern if removals observed;
  - (e) Example with real data.

W S Kendall, S B Connor (Warwick, York)

Perfect Epidemics

20 October 2025

3 / 30


# 1. Introduction to Perfect Simulation

- ① Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- ② The term “perfect simulation” (WSK, 1998) was chosen to encourage you to be suspicious: perfection is never achieved!
- ③ Key ideas of “classic CFTP”:
  - ▶ extend simulation *backwards* through time not forwards;
  - ▶ exploit monotonicity (couple maximal and minimal processes);
  - ▶ seek coalescence.
- ④ Simplest possible example: *random-walk-CFTP* (can boost to use Ising model to do simple image reconstruction).

Statistics

## Classic CFTP for a simple random walk (I)

- ① Consider a simple random walk on  $0:9 = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace the  $+1$  jump by “staying still”, **and**
  - ▶ at state 0 replace the  $-1$  jump by “staying still”.
- ② Conventional MCMC picks a starting point, then runs the simple random walk for long time till approximate equilibrium.



- ③ How long? One way to *estimate* this is to run two (or several?) coupled copies till they meet. If probability of meeting by time  $T$  is high, then deviation of  $X_T$  from equilibrium is statistically small;
- ④ Generally **not true** that location *at* coupling is a draw from equilibrium.

## Perfect Epidemics

2025-10-09

### 1. Introduction to Perfect Simulation

Here is a very brief summary of CFTP/ perfect simulation.

- ① Jim Propp described the discovery of CFTP as like walking down the street and suddenly noticing a 50\$ bill lying on the ground. He also said exact simulation “was in the air”: many related ideas
- ② In particular, “exact simulation” cannot somehow miraculously defeat numerical approximation error :-). And, as with all simulation, its validity depends on correctness of code!
- ③ We will illustrate these key ideas by considering a single very specific and simple example: CFTP for a simple random walk.
- ④ Propp & Wilson (1996) show how to vary random walk CFTP to get exact samples for a **critical** Ising model (Persi Diaconis: “Like seeing the landscape of Mars for the first time”). The Ising model with an external field can be used to model images, hence CFTP can be used for image reconstruction.

#### 1. Introduction to Perfect Simulation

- ① Propp & Wilson (1996) invented exact simulation / Coupling from the Past (CFTP) / perfect simulation;
- ② The term “perfect simulation” (WSK, 1998) was chosen to encourage you to be suspicious: perfection is never achieved!
- ③ Key ideas of “classic CFTP”:
  - ▶ extend simulation *backwards* through time not forwards;
  - ▶ exploit monotonicity (couple maximal and minimal processes);
  - ▶ seek coalescence.
- ④ Simplest possible example: *random-walk-CFTP* (can boost to use Ising model to do simple image reconstruction).

## Perfect Epidemics

2025-10-09

This random process is asymmetric simple random walk on the integers, modified by forbidding any transitions outside the specified state-space 0:9.

- ① The process is irreducible *and* aperiodic, so equilibrium is a truncated Geometric distribution (hint: use reversibility to check this!).
- ② Of course even conventional MCMC is overkill here: simulation directly from the equilibrium distribution is simple and efficient.
- ③ This observation was originally formalized by Aldous. Here we use “synchronous” coupling – variants are possible but not particularly useful.
- ④ In the case of reflecting simple random walk, coupling can only occur at the boundary points 0 or 9! So here the blue and red evolutions can only couple at a boundary; thus self-evidently the value at time of coupling cannot be a draw from the equilibrium!


#### Classic CFTP for a simple random walk (I)

- ① Consider a simple random walk on  $0:9 = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ .
  - ▶  $\mathbb{P}[+1 \text{ jump}] = p \in (0, 1)$ , while  $\mathbb{P}[-1 \text{ jump}] = 1-p$ , **except that**
  - ▶ at state 9 replace the  $+1$  jump by “staying still”, **and**
  - ▶ at state 0 replace the  $-1$  jump by “staying still”.
- ② Conventional MCMC picks a starting point, then runs the simple random walk for long time till approximate equilibrium.
- ③ How long? One way to estimate this is to run two (or several?) coupled copies till they meet. If probability of meeting by time  $T$  is high, then deviation of  $X_T$  from equilibrium is statistically small;
- ④ Generally **not true** that location *at* coupling is a draw from equilibrium.



## Classic CFTP for a simple random walk (I)

- 1 Start at top (9) and bottom (0) at negative time  $-T$ , run to time 0.



- 2 If not coupled by time 0, then back-off to time  $-2T$  and repeat.  
**NB:** re-use randomness!
- 3 May need to iterate back-off doubling several times.
- 4 When coupled, top and bottom yield a common value at time 0.
- 5 The common value (golden thread) is an exact draw from equilibrium!

Statistics

## A little CFTP theory

- 1 What if monotonicity fails? or there isn't a sensible "maximal" process?  
Ideas (WSK, 1998):

- ▶ cross-couple upper and lower envelope processes;
- ▶ dominate by amenable "dominating process" (time-reversible, can draw from equilibrium, can couple target processes below dominating process).

- 2 Theoretical limits: *in principle*

- ▶ Classical CFTP equivalent to uniform ergodicity (Foss & Tweedie, 1998);
- ▶ Dominated CFTP achievable under geometric ergodicity (WSK, 2004);
- ▶ Dominated CFTP can work in some non-geometrically ergodicity cases (SBC & WSK, 2007a; *nb* corrigendum SBC & WSK, 2007b).


- 3 Dominated CFTP delivers perfect simulation for stable point processes (WSK & Møller, 2000);

- 4 Detailed expositions: WSK (2005), Huber (2015).  
(Want to implement CFTP in R? see WSK, 2015.)

Statistics

## Perfect Epidemics

2025-10-09



- 1 Ideally one needs to choose  $T$  neither too small nor too large. But the result is not particularly sensitive to this.
- 2 Very important in this binary back-off that one doesn't use different jump proposals for the same time  $-t$ ! Couple to **re-use randomness**.
- 3 *Binary back-off procedure*, so if initially  $T$  too small then at most four-fold extra work compared to conventional MCMC!
- 4 Re-use of randomness means there is no point in extending the binary backoff beyond coalescence: the common value at time 0 will be the same however far we extend into the past with further back-offs.
- 5 Why is the common value an exact draw from equilibrium? Informally, because one would get the same result however far one backed-off: therefore the draw is effectively a draw from time  $-\infty$ . The golden thread can be viewed as a perfect draw from the last segment of such a simulation (this sort of device is very well-known to ergodic theorists). Remarkably, this can easily be converted into a fully rigorous proof!

## Perfect Epidemics

2025-10-09

### A little CFTP theory

A little CFTP theory

- What if monotonicity fails? or there isn't a sensible "maximal" process? Ideas (WSK, 1998):
  - cross-couple upper and lower envelope processes;
  - dominate by amenable "dominating process" (time-reversible, can draw from equilibrium, can couple target processes below dominating process).
- Theoretical limits: *in principle*
  - Classical CFTP equivalent to uniform ergodicity (Foss & Tweedie, 1998);
  - Dominated CFTP achievable under geometric ergodicity (WSK, 2004);
  - Dominated CFTP can work in some non-geometrically ergodicity cases (SBC & WSK, 2007a; *nb* corrigendum SBC & WSK, 2007b).
- Dominated CFTP delivers perfect simulation for stable point processes (WSK & Møller, 2000);
- Detailed expositions: WSK (2005), Huber (2015).  
(Want to implement CFTP in R? see WSK, 2015.)

- 1 To be computationally effective, WSK (1998) ideas still require a (perhaps very weak) notion of an associated partial order, and also the ability to simulate the dominating process.
- 2 Basic ideas: use the notion of regenerative sets ("small sets"), and (Foster-)Lyapunov arguments. Note that the resulting recipes tend not to be computationally practical: they simply suggest the *possibility* of (possibly computationally infeasible) CFTP. They are intended to frame a challenge: when can one find *practical* methods?
- 3 (Locally finite) point processes are *generally* not accessible *via* Classic CFTP; the "top" pattern would have points *everywhere*. The "in principle" results encourage us to search for CFTP for a wide variety of challenge problems [for example,  $M/G/k$  multiserver queues; see Sigman (2011); SBC & WSK (2015); SBC (2020)].
- 4 This ends the visual introduction to CFTP. People who are interested in practical CFTP may find it useful to work through WSK (2015).

## 2. Perfect Epidemics: a challenge problem for CFTP

### S-I-R deterministic epidemic:

based on susceptibles  $s$ , infectives  $i$ , removals  $r$ :

$$\begin{aligned} s' &= -\alpha s i, \\ i' &= (\alpha s - \beta) i, \\ r' &= \beta i. \end{aligned}$$

Constant total population  $s + i + r = n$ .

**S-I-R stochastic epidemic:** a Markov chain  $(S, I, R)$  with transitions

$$\begin{aligned} \text{Infection: } S &\rightarrow S - 1, \quad I \rightarrow I + 1 & \text{at rate } \alpha S I, \\ \text{Removal: } I &\rightarrow I - 1, \quad R \rightarrow R + 1 & \text{at rate } \beta I. \end{aligned}$$

Both make an **unrealistic assumption**: **homogeneous mixing**.

In contrast, Fraser *et al* (2021) use a **UK model with  $10^6$  agents!**

There are *many* important inferential questions (Cori & Kucharski, 2024).



## The first question asked about a new epidemic

“What is the R-number?”

The R-number is  $\alpha s_0 / \beta$ : mean number of new infectives produced per infective at *start* of epidemic with initially  $s_0$  susceptibles.

Whittle (1955)’s threshold theorem: R-number  $\gg 1$  means strongly positive chance of epidemic infecting significant proportion of the population.

**Wikipedia:** “The British-registered *Diamond Princess* was the first cruise ship to have a major [COVID-19] outbreak on board, with the ship quarantined at Yokohama from 4 February 2020 for about a month. Of 3711 passengers and crew, around 700 people became infected and 9 people died.”

Evidently  $\alpha s_0 / \beta \gg 1$  – as was sadly later confirmed, a sorrow for us all.



## Perfect Epidemics

2025-10-09

### 2. Perfect Epidemics: a challenge problem for CFTP

- Two classical models (one deterministic, one stochastic) from mathematical epidemiology. Even the exceedingly simple case of deterministic S-I-R permits only *partial* closed-form solution.
- The assumption of homogeneous mixing of population greatly simplifies mathematical *and* statistical issues. Without this assumption one risks huge numbers of parameters, very bad news statistically speaking. Much research dealing with more realistic models, for example, UK model of Fraser *et al* (2021) with 1000000 agents.
- The work presented in this talk considers only perfect simulation for homogeneous mixing. Before running, one must learn to walk!
- Cori & Kucharski (2024) provide a broad and stimulating perspective on statistical challenges from a very practical point of view!

2. Perfect Epidemics: a challenge problem for CFTP  
S-I-R deterministic epidemic:  
based on susceptibles  $s$ , infectives  $i$ , removals  $r$ :  
 $s' = -\alpha s i$ ,  
 $i' = (\alpha s - \beta) i$ ,  
 $r' = \beta i$ .  
Constant total population  $s + i + r = n$ .  
S-I-R stochastic epidemic: a Markov chain  $(S, I, R)$  with transitions  
Infection:  $S \rightarrow S - 1, I \rightarrow I + 1$  at rate  $\alpha S I$ ,  
Removal:  $I \rightarrow I - 1, R \rightarrow R + 1$  at rate  $\beta I$ .  
Both make an unrealistic assumption: **homogeneous mixing**.  
In contrast, Fraser *et al* (2021) use a UK model with  $10^6$  agents!  
There are many important inferential questions (Cori & Kucharski, 2024).

## Perfect Epidemics

2025-10-09

### The first question asked about a new epidemic

Whittle (1955) uses a coupling argument that doesn’t mention coupling!

Some recent history

- I am skipping over very influential early work on the *Diamond Princess* incident, which particularly focussed on the incubation period (a time when the subject is infectious, perhaps at time-varying rate, before symptoms appear). This is related to drawing inferences about  $\beta$ .
- See Mizumoto *et al* (2020) for very early analysis of the *Diamond Princess* outbreak.
- Question:** is there a dataset for the *Diamond Princess* outbreak? and how detailed is it? Rocklöv *et al.* (2020) focusses on *R*-number and refers to some public sources of cumulative daily totals.

The first question asked about a new epidemic  
“What is the R-number?”  
The R-number is  $\alpha s_0 / \beta$ : mean number of new infectives produced per infective at start of epidemic with initially  $s_0$  susceptibles.  
Whittle (1955)’s threshold theorem: R-number  $\gg 1$  means strongly positive chance of epidemic infecting significant proportion of the population.  
**Wikipedia:** “The British-registered *Diamond Princess* was the first cruise ship to have a major (coronavirus) outbreak on board, with the ship quarantined at Yokohama from 4 February 2020 for about a month. Of 3711 passengers and crew, around 700 people became infected and 9 people died.”  
Evidently  $\alpha s_0 / \beta \gg 1$  – as was sadly later confirmed, a sorrow for us all.



## Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- 1 Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- 2 Inference is **really tough**: hard to get information about infection times;
- 3 It is all **especially tough** in early stages. Answers are most needed when hardly any information is available (a simplified example for a Warwick UG second-year statistics module shows how tough this can be);
- 4 Markov chain Monte Carlo (MCMC) can be used (see next slide) but what about burn-in?
- 5 Can we use **perfect simulation**?



## Perfect Epidemics

2025-10-09

### Inference on the R-number

- The “R-number” was the subject of continuing interest throughout the COVID-19 pandemic!
- Notice that the “R-number” is as much a social construct as a biological one, and in real situations will interact with behaviour in subtle and complicated ways.
- We will seek to make headway in using perfect simulation to aid inference.
- The PhD thesis of Bensoussane (2025) is a useful source of references.
- We need to fix on a good setting for demanding challenge problems.

#### Inference on the R-number

Important, because the R-number controls severity of epidemic. However:

- Modelling is **tough**. *Either* massive assumptions (homogeneous mixing) *or* very many parameters;
- Inference is **really tough**: hard to get information about infection times;
- It is all **especially tough** in early stages. Answers are most needed when hardly any information is available (a simplified example for a Warwick UG second-year statistics module shows how tough this can be);
- Markov chain Monte Carlo (MCMC) can be used (see next slide) but what about burn-in?
- Can we use **perfect simulation**?

## An easier question

The simplest possible variant of contact tracing:

“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O’Neill & Roberts, 1999; Gibson & Renshaw, 2001)

Important first step: think about generation of an *unconditioned* epidemic.

- 1 Suppose  $n, \alpha, \beta$  are known. Eventually removal times are observed, but unobserved infection times **must be inferred**.
- 2 Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.
- 3 Poisson point processes of *appropriate rates* yield an S-I-R epidemic.
- 4 First step: evolve whole S-I-R trajectory in *algorithmic time* (alter potential infections and removals using immigration-death in discrete algorithmic time).
- 5 Result: *trajectory-valued chain*, unconditioned S-I-R as equilibrium.



## Perfect Epidemics

2025-10-09

### An easier question

Focus on just one aspect of inference:

- 1 Avoid parameter inference for now; consider the large number of nuisance parameters given by (typically unobserved) infection times.
- 2 Timelines: instead of tracking individuals as in the stochastic epidemic model, track the numbers of individuals in various categories.
- 3 Poisson point processes generate incidents, which then collectively influence the S-I-R trajectory.
- 4 I learned the phrase “algorithmic time” from Andrew Stuart. Move *from* Poisson points spread out along timelines *to* discrete immigration-death processes evolving in algorithmic time.
- 5 Exploit the classic connection for reversible Markov processes, between conditioning and restriction of state space.
- 6 Work in *continuous* algorithmic time, obtain a dynamically reversible Markov process, prove restriction=conditioning still works.

#### An easier question

The simplest possible variant of contact tracing:  
“When did the infections occur, supposing we only observe removals?”  
(Gibson & Renshaw, 1998; O’Neill & Roberts, 1999; Gibson & Renshaw, 2001)

Important first step: think about generation of an *unconditioned* epidemic.

- Suppose  $n, \alpha, \beta$  are known. Eventually removal times are observed, but unobserved infection times must be inferred.
- Visualize  $n$  timelines, along which incidents are scattered:
  - ▶ potential removals, activated if timeline is infected;
  - ▶ potential infections, activated if timeline is infected *and* if designated target timeline is lowest uninfected timeline.
- Poisson point processes of *appropriate rates* yield an S-I-R epidemic.
- First step: evolve whole S-I-R trajectory in *algorithmic time* (alter potential infections and removals using immigration-death in discrete algorithmic time).
- Result: *trajectory-valued chain*, unconditioned S-I-R as equilibrium.



## From incidents to unconditioned epidemic trajectories (1/3)




Figure 1: Light-orange circles denote potential infections (arrows point upwards to targets); light-brown circles denote potential removals.


## Perfect Epidemics

2025-10-09

As implied above, incidents comprising a single innovation are produced by Poisson point processes on each of the timelines.

Colour-coding distinguishes infections from deaths. Note that infections need to specify which timeline is going to be infected.

- Rates of Poisson processes generating the (proposed) incidents are chosen to ensure that numbers of actual susceptibles, infectives and removals form a Markov chain with S-I-R epidemic rates.
- For an infection incident to be activated, it must sit on an infected portion of its timeline and must point to the least of the uninfected timelines (calculating at the process time of the incident). This is computationally expensive, (cost is cubic in  $\alpha T$ , not quadratic!) but simplifies monotonicity arguments.
- For a removal incident to be activated, it must sit on an infected portion of its timeline.



## From incidents to unconditioned epidemic trajectories (2/3)





Figure 2: (a) *Infection* activates if target lies on the lowest uninfected timeline; and (b) *removal* activates if on infected timeline; remove lowest infected (purple disk).

## Perfect Epidemics

2025-10-09

- Here is the resulting evolution of an epidemic.
- Note that here we specify the initial number of infectives (here we specify  $I_0 = 1$ ).
- However we can do better: given a prior for  $I_0$ , at the start of each cycle we could introduce an accept-reject move which alters  $I_0$  while respecting detailed balance.
- Options include use of an independence sampler, or of a random-walk Metropolis sampler.






Figure 3: A step in algorithmic time for the unconditioned epidemic simply involves replacing all original incidents by an entirely new set of incidents.

2025-10-09

- An update in algorithmic time, if there is no conditioning, simply involves replacement of one set of incidents by another. (In this illustration there is a radical change, as epidemic dies out at an early stage.)
- But we need to refine this if we are to take account of conditioning: a simple accept/reject procedure will almost always reject proposals of *entire* innovations involving all of both infection and removal incidents (and  $I_0$  if we update this too).
- We need to *localize* the proposal, by considering one change at a time. (Intuitively, a “pixel-by-pixel” analysis working along the time axis.)




Figure 3: A step in algorithmic time for the unconditioned epidemic simply involves replacing all original incidents by an entirely new set of incidents.

## Crucial technical point

- Updates in algorithmic time  $\tau$  are then (algorithmic-)time-reversible: so restriction to a subset  $S$  of state-space (the *activated / conditioned* removals to occur precisely at the specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $\mathcal{R}$ s);
  - 2 Re-sample timelines (though not times) of  $\mathcal{R}$ s;
  - 3 Replace infections ( $\mathcal{I}$ s).
- Re-express using *continuously varying*  $\tau$ . Process time runs over  $[0, T]$ .
  - 1 For  $2nT < \tau < (2n+1)T$ , update old  $\mathcal{R}$ s with times in  $(0, \tau - 2nT)$ ;
  - 2 For  $\tau = (2n+1)T$ , resample timelines (not times) of  $\mathcal{R}$ s;
  - 3 For  $(2n+1)T < \tau < (2n+2)T$ , update old  $\mathcal{I}$ s in  $((2n+2)T - \tau, T)$ .
- Thus the original update is expressible as a (continuous) composition of updates, each of which satisfies detailed balance in equilibrium.
- The connection “restriction=conditioning” still holds.
- Crucially, re-sampling step 2 ensures composite evolution is irreducible over  $S$ ! (So equilibrium under conditioning is unique.)

2025-10-09

### Crucial technical point


It is convenient to express the algorithmic update in *continuous* time: applying a given innovation over a continuous-time cycle of length  $2T$ . The result is no longer reversible, but each individual steps satisfies detailed balance, so the important connection between conditioning and constraint remains valid.

- In the case of discrete algorithmic time, arranging for conditioning would involve horrendous computations. Instead we unroll a “pixel-by-pixel” analysis into continuous algorithmic time.
- Intuitively, we analyze first removals ( $\tau$  runs over algorithmic time range  $(0, T)$ ) and then infections, ( $\tau$  runs over algorithmic time range  $(T, 2T)$ ) each working “pixel-by-pixel” along the process time axis.
- It turns out to be convenient to work *forwards in process time* for removals, then *backwards in process time* for infections – and to re-assign random timelines (but not times) for removals in between these two phases (at algorithmic time  $\tau = T$ ).

**Crucial technical point**

- Updates in algorithmic time  $\tau$  are then (algorithmic-)time-reversible: so restriction to a subset  $S$  of state-space (the *activated / conditioned* removals to occur precisely at the specified set of times) implies a new equilibrium which is the old equilibrium conditioned to lie in  $S$ .
- For later purposes it is convenient to stage the replacement as follows:
  - 1 Replace removals ( $\mathcal{R}$ s);
  - 2 Re-sample timelines (though not times) of  $\mathcal{R}$ s;
  - 3 Replace infections ( $\mathcal{I}$ s).
- Re-express using continuously varying  $\tau$ . Process time runs over  $[0, T]$ .
  - 1 For  $2nT < \tau < (2n+1)T$ , update old  $\mathcal{R}$ s with times in  $(0, \tau - 2nT)$ ;
  - 2 For  $\tau = (2n+1)T$ , resample timelines (not times) of  $\mathcal{R}$ s;
  - 3 For  $(2n+1)T < \tau < (2n+2)T$ , update old  $\mathcal{I}$ s in  $((2n+2)T - \tau, T)$ .
- Thus the original update is expressible as a (continuous) composition of updates, each of which satisfies detailed balance in equilibrium.
- The connection “restriction=conditioning” still holds.
- Crucially, re-sampling step 2 ensures composite evolution is irreducible over  $S$ ! (So equilibrium under conditioning is unique.)

## Free evolution evolving in continuous algorithmic time



GIF MP4

Statistics

## 3. Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point). Consequently:
  - conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals.


### Implications:

- a removal can be introduced only if it doesn't activate;
- a conditioned removal timeline can be altered only if it doesn't de-activate;
- an infection cannot be removed if that action loses a conditioned removal;
- an infection can be introduced only if no new observed removals result.
- Does this produce a *feasible* and suitably *monotonic* algorithm?
- Housekeeping details** used to establish that monotonicity still works: *laziest feasible epidemic* (LFE) and *no-fly zone* (NFZ).

Statistics

## Perfect Epidemics

2025-10-09



We discuss a single step in algorithmic time. **Conditioning not applied here!**

- Start with initial configuration, use immigration-death process of removals to replace old removals. **To avoid premature extinction in demonstration figures, condition on no removals in timeline 1!**
- Because we are not yet conditioning, all old removals have now been removed, and all new removals have been added.
- Once all removals have been processed, resample timelines of all removals. Jitter back and forth here to emphasize this important step.
- Now we resample infections, *but this time working backwards through process time*.
- At the end all removals and infections have been removed and replaced by new ones.

## Perfect Epidemics

2025-10-09

### 3. Conditioning on observed removals

Monotonicity is a key concept here. (Note however that we can still sometimes generate perfect simulation algorithms even when monotonicity does not obtain: we don't need to do this here but should bear it in mind for more general problems in the future.)

- “Restriction=monotonicity **needs careful proof** in the continuum limit: discretization,” convergence stationnaire” (“parking convergence”).
- The heart of the matter lies in establishing (some variant of) monotonicity.
- Additionally, for a complicated algorithm such as this one, it is important to test the underlying logic by implementing the algorithm in computer code (which often reveals the falsity of hidden assumptions!).
- Geometric notions of **LFE** and **NFZ** provide a useful vocabulary for formulating how this works out. They facilitate a rigorous argument for arguing for monotonicity, and are also computationally helpful.

3. Conditioning on observed removals

- The trajectory-valued chain is *dynamically reversible*, in *continuous algorithmic time*.
- Irreducibility is *vital* (otherwise equilibrium depends on starting point). Consequently:
  - conditioned removals must be able to change timeline (but not time of occurrence);
- Forbidding removal of observed removals, and forbidding creation of new activated removals, yields a modified chain whose invariant probability measure conditions on observed pattern of removals. Implications:
  - a removal can be introduced only if it doesn't activate;
  - a conditioned removal timeline can be altered only if it doesn't de-activate;
  - an infection cannot be removed if that action loses a conditioned removal;
  - an infection can be introduced only if no new observed removals result.
- Does this produce a *feasible* and suitably *monotonic* algorithm?
- Housekeeping details used to establish that monotonicity still works: *laziest feasible epidemic* (LFE) and *no-fly zone* (NFZ).



## Initial conditioned epidemic





Figure 4: Initial conditioned epidemic, with conditioned removals indicated using purple circles (and purple disks when non-target timelines are infected).

## Perfect Epidemics

2025-10-09

Here is a different initial epidemic, including *conditioned removals*, indicated by purple circles.

- Times of observed removals are fixed, so removed zone does not change.
- This evolves in algorithmic time as in the unconditioned case (working “pixel-by-pixel”), but forbidding any change that breaks conditioning.
- Dealing with removals amounts to removing all inactivated removals and replacing them with a new set of removals, and then resampling the conditioned removal timelines, accepting only those changes which do not de-activate the conditioned removals.
- Dealing with *infections* requires more care.



## Conditional epidemic update




Figure 5: Epidemic updated under restriction: all conditioned removals remain activated, no new removals are activated. Green infections have been “perpetuated”.

## Perfect Epidemics


2025-10-09

The update proceeds as follows:

1. Accept inactivated removals only if targets avoid infected region.
2. Accept conditioned removals only if targets fall in infected region.
3. Accept new infections only if this doesn't result in infecting an inactivated removal.
4. Retain old infections only if rejection would result in a conditioned removal *not* being infected, in which case retain old infection as *perpetuated* infection.

Options 1 and 2 are directly imposed by the requirement of not changing the set of conditioned removals.

For options 3 and 4 there is a useful correspondence to two recursively-defined geometric structures: the *laziest feasible epidemic* (LFE: *slowest possible epidemic compatible with observed removals*) and the *no-fly zone* (NFZ: *forbid all infections that might infect this zone*).



## Laziest feasible epidemic (LFE)




Figure 6: LFE computed recursively working right-to-left: slowest sequence of infections (including perpetuated infections) generating all observed removals in order.

## Perfect Epidemics


2025-10-09

The **LFE** is a collection of intervals  $[s_j, \infty)$ , ( $j$  indexing timelines). It is recursively defined, working *backwards* in process time. It is computable at process time  $t$  from the *updated* system of incidents over  $[t, \infty)$ . Here  $s_j$  is the latest time of an infection with target in infected region and infectee  $j+1$ :

$$s_j \leq s_{j+1}, \text{ and}$$

$$s_j \leq \inf\{s : s \text{ is the time of a conditioned removal with target } j\}.$$

- In the example the **LFE** boundary differs from the epidemic trajectory, over time intervals  $[1.43, 3.45]$ ,  $[4.71, 5.56]$ .
- At times 0.88, 3.45, 4.08, 5.56 there are perpetuations (green arrows: an old infection has to be retained to ensure conditioning is not broken); two of these perpetuations (0.88, 4.08) form part of the epidemic trajectory. Note a new infection can be in the **LFE** but *not* in the epidemic trajectory.



## No-fly zone (NFZ)




Figure 7: NFZ computed recursively working right-to-left: it traces the region of timelines that must not be infected if one is not to activate unobserved removals.

## Perfect Epidemics


2025-10-09

The **NFZ** is used to decide whether to accept proposals of infections targeting the susceptible region:


- New infections with targets in the infection zone are assessed to see whether their introduction would activate removals, by connecting the infectious zone to a timeline leading to such an activation (a “no-fly” portion of the timeline);
- An infection with target in the susceptible zone may be such that it makes a portion of its timeline *before* its time into “no-fly”.

In this particular case neither of these two mechanisms play a part.

The **NFZ** is evaluated recursively, working backwards in process time, and again is computable at process time  $t$  from the *updated* system of incidents over  $[t, \infty)$ . Inductive argument shows that **NFZ** and **LFE** depend monotonically on the epidemic trajectory. This implies monotonicity of the algorithmic-time evolution of the conditioned epidemic: hence permitting **CFTP**!




## Conditioned evolution evolving in continuous algorithmic time



GIF MP4

## Perfect Epidemics

2025-10-09




### Conditioning applied here!

- Start with initial configuration, use immigration-death process of removals to replace old removals **except** in the infected region.
- All old removals have now been removed, and some new removals have been added.
- Resample timeline of each removal if this does not violate conditioning. Jitter back and forth here.
- Now resample infections, *but this time working backwards through process time*. If addition proposed, add only if pattern of observed removals are unchanged. If deletion proposed, and deletion would change pattern of observed removals, then *perpetuate*.
- We do not discuss the careful argument for monotonicity here.

## 4. Example


- Smallpox outbreak in a closed community of 120 individuals in Abakaliki, Nigeria (much studied! see page 125 of [Bailey, 1975](#)).
- **Assume**
  - ▶ first observed removal is also the first removal: under a plausible improper prior we can then deduce what is the distribution of infectives  $I_0$  at time 0;
  - ▶ *all* removals are recorded;
  - ▶ no further removals after last observed removal (makes life easier).
- Coding in *julia* ([Bezanson et al., 2017](#)), animates (GIF or MP4) a perfect simulation of a draw from unobserved pattern of infections.



## Perfect Epidemics

2025-10-09

### 4. Example



We could vary the assumption concerning the initial and final numbers of infectives while still using essentially the same perfect simulation algorithm! This amounts to presuming some Bayesian prior knowledge.

- Blue is maximal, red is minimal, start at extremal epidemics.
- Suppose we only have a prior for the number of initial infectives?
  - Use independence sampler: draw from (unconditioned) number of initial infectives, accept if this leads to observed removals exactly as conditioned;
  - Or employ a random-walk Metropolis sampler, if prior for initial number of infectives is suitably monotonic;
  - Or use modified independence sampler and notions of optimal scaling.

### Technical note:

The language *julia* ([Bezanson et al., 2017](#)), allows for rapid development by using an expressive type-based syntax (very useful for involved algorithms), and delivers remarkably fast execution using “just in time” compilation techniques.

## So what?

- What about accept-reject methods?
- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for *unknown*  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on *different*  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (next steps after SBC & WSK, 2024)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;
  - ▶ construct steepest ascent algorithm (in effect, variant of Robbins-Monro stochastic optimization) to find *maximum a posteriori* estimates of  $\alpha$  and  $\beta$ ;
  - ▶ or even, with some computational effort, compute the entire posterior joint density for  $\alpha$  and  $\beta$ !
- Finally: can we generalize to other suitable compartment models?

Warwick  
Statistics

W S Kendall, S B Connor (Warwick, York)

Perfect Epidemics

20 October 2025

24 / 30

## Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.
- *Of course* detailed models resist perfect simulation (but it will be helpful to compare with a simpler model using fewer parameters).
- Experiments suggest CFTP out-competes non-naïve accept-reject.
- Still to be done: seek faster CFTP; statistical estimation of parameters, generalization to other compartment models.
- Thank you for your attention! **QUESTIONS?**



Warwick  
Statistics

W S Kendall, S B Connor (Warwick, York)

Perfect Epidemics

20 October 2025

25 / 30

## Perfect Epidemics

2025-10-09

### So what?

Accept-reject approach for non-trivial problems requires *much* more computation.

Perfect simulation supplies a *well-behaved* stochastic integration mechanism, using repeated MCMC samples in a Monte Carlo calculation. Rao-Blackwellization available (interpolate between conditioned removals using nonlinear growth process bridges!).

Other compartment models presenting harder challenges:

- ① Split population into interacting sub-populations;
- ② (For purposes of insight even if impractical.) Treat each individual as a sub-population on its own;
- ③ Allow parameters to change at specific times of day or week;
- ④ Allow for some activated removals to be unobserved!
- ⑤ Current recipe then becomes non-monotonic! Perhaps cross-over arguments might help?

So what?

- What about accept-reject methods?
- Why this emphasis on unobserved infections given fixed  $\alpha$  and  $\beta$ , when we need inference on R-number  $\alpha n/\beta$  for unknown  $\alpha$  and  $\beta$ ?
- Good question. But a re-weighting argument allows us to get (unbiased) estimates based on different  $\alpha$  and  $\beta$ . The perfect simulation provides exact simulation-based computation to integrate out pattern of unobserved infections.
- So (next steps after SBC & WSK, 2024)
  - ▶ estimate likelihood test statistic for specified  $\alpha$  and  $\beta$ ;
  - ▶ construct steepest ascent algorithm (in effect, variant of Robbins-Monro stochastic optimization) to find maximum a posteriori estimates of  $\alpha$  and  $\beta$ ;
  - ▶ or even, with some computational effort, compute the entire posterior joint density for  $\alpha$  and  $\beta$ !
- Finally: can we generalize to other suitable compartment models?

## Perfect Epidemics

2025-10-09

### Conclusion

There is much still to be done!

Warwick talk (January 2025): Paul Jenkins asked whether this relates to the “look-down” argument in population genetics (Donnelly & Kurtz, 1996)? Provisional answer, after rereading Donnelly & Kurtz (1996):

the “look-down” argument is related because it also involves a particle model as discussed in Liggett (1985). I can’t see a closer connection.

Conclusion

- If MCMC burn-in is a concern, try to build a perfect simulation!
- CFTP works even for significantly complex and relevant models of real-life phenomena.
- *Of course* detailed models resist perfect simulation (but it will be helpful to compare with a simpler model using fewer parameters).
- Experiments suggest CFTP out-competes non-naïve accept-reject.
- Still to be done: seek faster CFTP; statistical estimation of parameters, generalization to other compartment models.
- Thank you for your attention! **QUESTIONS?**



References I

Bailey, N.T.J. (1975) *The mathematical theory of infectious diseases and its applications*, 2nd Ed. ed. Griffin.

Bensoussane, H. (2025) Bayesian Individual-level Epidemic Models : Accounting for Missing Data and Utilising Covariate Information (PhD No. January).

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V.B. (2017) Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, **59**, 65–98.

Cori, A. & Kucharski, A. (2024) Inference of epidemic dynamics in the COVID-19 era and beyond. *Epidemics*, **48**, 100784.

Donnelly, P. & Kurtz, T.G. (1996) A countable representation of the Fleming-Viot measure-valued diffusion. *The Annals of Probability*, **24**, 698–742.

Foss, S.G. & Tweedie, R.L. (1998) Perfect simulation and backward coupling. *Stochastic Models*, **14**, 187–203.

Fraser, C. (2021) OpenABM-Covid19: Agent-based model for modelling the Covid-19 and Contact-Tracing.

Gibson, G.J. & Renshaw, E. (1998) Estimating parameters in stochastic compartmental models using Markov chain methods. *Mathematical and Medical Biology*, **15**, 19–40.

Gibson, G.J. & Renshaw, E. (2001) Likelihood estimation for stochastic compartmental models using Markov chain methods. *Statistics and Computing*, **11**, 347–358.



References III

Sigman, K. (2011) Exact simulation of the stationary distribution of the FIFO M/G/c queue. *Journal of Applied Probability*, **48**, 209–213.

Whittle, P. (1955) The outcome of a stochastic epidemic—a note on Bailey’s paper. *Biometrika*, **42**, 116–122.

WSK (1998) Perfect Simulation for the Area-Interaction Point Process. *Probability towards 2000* (Accardi, L. & Heyde, C.C. eds). Springer-Verlag, pp. 218–234.

WSK (2004) Geometric ergodicity and perfect simulation. *Electronic Communications in Probability*, **9**, 140–151.

WSK (2005) Notes on Perfect Simulation. Singapore: World Scientific, pp. 93–146.

WSK (2015) Introduction to CFTP using R. *Stochastic geometry, spatial statistics and random fields, Lecture notes in mathematics*. Springer, pp. 405–439.

WSK & Møller, J. (2000) Perfect simulation using dominating processes on ordered spaces, with application to locally stable point processes. *Advances in Applied Probability*, **32**, 844–865.



References II

Huber, M.L. (2015) *Perfect Simulation*. Boca Raton: Chapman; Hall/CRC.

Liggett, T.M. (1985) *Interacting particle systems*. Springer Verlag.

O’Neill, P.D. & Roberts, G.O. (1999) Bayesian Inference for Partially Observed Stochastic Epidemics. *Journal of the Royal Statistical Society Series A: Statistics in Society*, **162**, 121–129.

Propp, J.G. & Wilson, D.B. (1996) Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structures and Algorithms*, **9**, 223–252.

Rocklöv, J., Sjödin, H., & Wilder-Smith, A. (2020) COVID-19 outbreak on the Diamond Princess cruise ship: estimating the epidemic potential and effectiveness of public health countermeasures. *Journal of Travel Medicine*, **27**, 7 pp.

SBC (2020) Omnithermal Perfect Simulation for Multi-server Queues. *ACM Transactions on Modeling and Computer Simulation*, **30**, 1–15.

SBC & WSK (2007b) Perfect simulation for a class of positive recurrent Markov chains (corrigendum). *Annals of Applied Probability*, **17**, 1808–1810.

SBC & WSK (2007a) Perfect simulation for a class of positive recurrent Markov chains. *Annals of Applied Probability*, **17**, 781–808.

SBC & WSK (2015) Perfect simulation of M/G/c queues. *Advances in Applied Probability*, **47**, 1039–1063.

SBC & WSK (2024) Perfect Epidemics.



Image information

Image	Attribution	
Book of Kells	Huber Gerhard	CC BY 4.0
Classic CFTP for a simple random walk	Result of code written by WSK	
Diamond Princess	Alpsdake	CC BY-SA 4.0
Epidemic CFTP images and animation	Result of code written by WSK	

Previous instances of this talk

Date	Title		Location	
19/04/24	Perfect Epidemics	Short Research Talk	12mn	Warwick
15/05/24	McMC+Perfect Simulation	Graduate Seminar, Aristotle Univ.	50mn	Thessaloniki
17/01/25	Perfect Epidemics	Applied Probability Seminar	50mn	Warwick
27/06/25	Perfect Epidemics	UK Research Network Stochastics	45mn	Liverpool





# Other technical information

## Software used in computations

Software	Version	Branch	Last commit
quarto	1.6.39	—	
Running under julia	1.11.5	—	
EpidemicsCFTP	2.2.532	develop	Tue Jul 8 17:13:42 2025 +0100
EpidemicsUtilities	0.1.2.176	main	Fri Sep 26 15:11:24 2025 +0100
This quarto script	0.2.2.719	2025-10-09-Dublin-preparation	Thu Oct 9 10:38:07 2025 +0100

## Project information

<b>Version:</b>	0.2.2.721 (2025-10-09-Dublin-preparation)
<b>Author:</b>	Wilfrid Kendall <w.s.kendall@warwick.ac.uk>
<b>Date:</b>	Thu Oct 9 13:04:22 2025 +0100

**Comment:**  
Fixed Fraser et al reference

