

# Greenland Sea Ice Floe Segmentation and Tracking Via the Ice Floe Tracker Algorithm

The Ice Floe Tracker (IFT) algorithm automatically identifies sea ice floes in marginal ice zones from optical satellite imagery, then uses a feature-matching approach to track individual ice floe rotation and displacement. The algorithm is described in Lopez-Acosta et al. (2019) and briefly summarized below. It was developed by Rosalinda Lopez-Acosta during her PhD work at University of California-Riverside under the guidance of Monica M. Wilhelmus (Lopez-Acosta, 2021). This dataset contains IFT results for nearly 20 years of satellite imagery from the Moderate Resolution Imaging Spectroradiometer (MODIS) instrument, aboard NASA's *Aqua* and *Terra* satellites. Analysis of this dataset is included in Lopez-Acosta (2021) and in Watkins et al. (2023).

This repository contains code to translate the original output, in proprietary MATLAB format, into cross-platform compatible GeoTiff and CSV formats. MATLAB data and script files are included as well as Python scripts to read and format the MATLAB data. In addition to the original implementation of IFT, we introduce a post-processing routine for quality control based on a logistic regression classifier. We also address differences in the image resolution in the initial processing for the 2020 data so that the final dataset is self-consistent. Access to additional datasets, in the form of true and false color MODIS imagery and National Snow and Sea Ice Data Center (NSIDC) Climate Data Record of Sea Ice Concentration (Meier et al., 2021), is required to run the scripts.

Questions? Contact: Daniel Watkins ([daniel\\_watkins@brown.edu](mailto:daniel_watkins@brown.edu)).

## The Ice Floe Tracker algorithm

Description of the IFT, with a flowchart.

## Setup

### Installing required software

The python environment used to run the scripts here can be recreated using the `ift_env.yml` file. After installing miniconda, open a terminal, navigate to the project folders, and run `conda env create -f ift_env.yml`

You'll also need to adjust the paths at the start of each script to point to the location where the imagery is stored.

### Downloading MODIS imagery via IFT Pipeline

The MODIS dataset is large, even when subsetting to the study area, and is therefore not included in this repository. To download the data, we use the Ice Floe Tracker Pipeline. The file `scripts/00_setup_ft_table.py` generates the set of CSV files in the folder `data/modis_download_spec_files`. To download the MODIS imagery on the Oscar HPC system at Brown, after installing the Ice Floe Tracker Pipeline, modify the Cylc graph in `flow_template_hpc.j2` to read:

```
R1 = global_setup => mkpaths<param_set> => pullfetchimage & pulljuliaimage => fetchdata<param_set>
& soit<param_set>
```

Copy the specification files to the `config` folder. Each year is run separately. For year 2019, as an example, load the IFT environment and run the python command

```
python workflow/scripts/flow_generator.py \
--csvfile "./config/fram_strait_spec_tables/location_specs_2019.csv" \
--template "flow_template_hpc.j2" \
--template_dir "./config/cylc_hpc" \
--crs "epsg3413" \
```

```
--minfloearea 100 \
--maxfloearea 90000
```

Then run the IFT pipeline in cylc via

```
cylc install -n fram_strait_images ./config/cylc_hpc && \
cylc validate fram_strait_images && \
cylc play fram_strait_images && \
cylc tui fram_strait_images
```

## Sea ice concentration data

We use the sea ice concentration Climate Data Record (Meier et al., 2021). After downloading the 2003-2020 CDR data, change the `sic_loc` parameter in the script `03_extract_shape_properties` to point to this location.

## Processing framework




Figure 1: Flowchart describing processing pathway. Blue arrows are inputs, red arrows are outputs. Light purple blocks indicate temporary files, and dark purple blocks are archived files.

The first step is to parse information in the `.mat` files and align the data into CSV files. The MATLAB code produces files with floe properties and floe positions for (a) all candidate floe segments and (b) for all floes that were matched to a subsequent image. The file `01_parse_ft_data.py` extracts the floe properties and positions from the MATLAB output. Along with the values originally in the `props.mat` file, it adds a `floe_label` so that tracked floes can be assembled into trajectories. The files `time_data.csv` were manually created using a variety of sources including saved diagnostic images and output from the SOIT python function. It maps the index in the `FLOE_LIBRARY` and `props.mat` to time stamps and specific satellites.

Next, we pull shapes from the `FLOE_LIBRARY.mat` data objects and place them into GeoTiffs. The MATLAB file structure efficiently holds the sparse dataset of labeled floe shapes. However it is not easily visualized or shared, and it is not self-describing. The script `02_extract_shapes.py` reads the data in the `FLOE_LIBRARY` and in the floe property tables, then creates a GeoTiff sharing dimensions and coordinate reference

system with the reference image `NE_Greenland.2017100.terra.250m.tif`. The file produced is an unfiltered segmented image where the labels of each floe correspond to the index in the `FLOE_LIBRARY`. A tracked floe will have different label numbers in each image.

Floe properties were initially calculated in MATLAB and are saved by the `01_parse_ft_data.py` script. There are differences in the algorithms used by scikit image region properties function and the identically named function in MATLAB. For future compatibility with the IFT Julia version, which uses scikit image, we recalculate region properties and add these to the floe property tables. Calculations are carried out in the script `03_extract_shape_properties.py`. This step also allows us to get consistent bounding boxes and row/col centroid data for the shapes. Using the shapes extracted in the previous step, and the truecolor and falsecolor images, we get the mean intensity for each color channel within each floe. The results are saved. This data is used for filtering true and false positives from the floe property tables.

The IFT segmentation step produces a set of candidate ice floes for matching. For estimates of the floe size distribution, ideally all detected floe shapes can be used (rather than only tracked floes). Tracking floes filters out candidate segments corresponding to bright patches in clouds, ice filaments, clumps of ice floes below the image resolution, and other similar objects due to the tendency of these objects to deform strongly between images. Buckley et al. (2023) used floe circularity, a function of the floe perimeter and area, to filter out false positives. However, the floe circularity is, in general, a necessary but not sufficient criterion. Many false positives also have similar circularity properties as real floes. We leverage the information from the truecolor and falsecolor images to perform logistic regression classification. We use tracked floes to form a set of true positives for the training dataset, and apply information on sea ice concentration and geometry to form a set of false positives. The logistic regression methodology is described in detail in a later section. The script `04_logistic_regression_classifier.py` trains, tests, and applies the classification method, and saves the results to the archive with the format `ift_raw_floe_properties_YYYY.csv`, where `YYYY` is the year.

After fitting and applying the logistic regression function, we apply a decision rule where all the floes marked as true positive and false positive using the manual criteria retain those labels, and the remaining (majority) of objects are automatically assigned to the two categories. The function `05_filter_geotiffs.py` reads the raw classified floe properties tables from the previous step and the raw segmented images, then removes all segments classified as false positives. These images are saved into the archive in the folder `labeled_clean`.

Finally, we calculate daily estimates of floe position, rotation, and displacement using the script `06_interpolate_trajectories.py`. We read all the property tables, extract the tracked floes only, and use linear interpolation to regrid the trajectories to a 24 h regular grid. Velocity is calculated in stereographic coordinates using forward differences then rotated into traditional north/south and east/west components. Rotation rates in the raw data represent the rotation of each object between overpasses of individual satellites, e.g. if a floe was observed by *Aqua*, then the rotation is calculated only if it is observed again by *Aqua* the next day. Let  $\theta_s(t_0, t_1)$  be the rotation observed in an ice floe by satellite  $s$  between times  $t_0$  and  $t_1$ . Set  $\Delta t = t_1 - t_0$ . Then the daily rotation rate  $\zeta$  is decided as follows: 1. If  $\theta_{Aqua}$  is defined and  $\theta_{Terra}$  is not, set  $\zeta = \theta_{Aqua}/\Delta t$ . 2. If  $\theta_{Terra}$  is defined and  $\theta_{Aqua}$  is not, set  $\zeta = \theta_{Terra}/\Delta t$ . 3. If both are defined, and  $|\theta_{Aqua} - \theta_{Terra}| < 30^\circ$ , calculate the average  $\zeta$  between the two estimates. 4. Otherwise, no estimated rotation is returned.

## Final dataset structure

TBD: File tree for the Digital Repository

TBD: Number of images, file types

TBD: Data structure for the floe property tables

Column	Description	Units
area	perimeter	major_axis
minor_axis	orientation	
x_pixel	Column coordinate in the original image	y_pixel
Row coordinate in the original image	convex_area	Area of the best-fit convex polygon
solidity	Ratio of area to convex area	Unitless
bbox1	bbox2	bbox3
bbox4	orig_idx	Index of row in floe property table in Matlab files
NA	satellite	Name of satellite
NA	floe_id	Unique label assigned to tracked floes
YYYY_NNNNN	datetime	Time of satellite overpass of the image centroid
YYYY-mm-dd HH:MM	x_stere	X-position of the

floe centroid in NSIDC N. Polar Stereographic|meters| |y\_stere|Y-position of the floe centroid in NSIDC N. Polar Stereographic|meters| |longitude|Longitude of the floe centroid|Decimal Degrees| |latitude|Latitude of the floe centroid|Decimal Degrees| |nsidc\_sic|Sea ice concentration of nearest grid cell from NSIDC CDR|Fraction| |label|Integer object label in the segmented image| NA |

## Cleaning the dataset using the logistic regression classifier

The logistic regression model maps a set of variables to a value from 0 to 1, interpreted as a probability of belonging to a class. To train and evaluate the model, we first need label a set of floes as true positives and false positives. We rely on the floe tracker to select true positive floes. We filter the tracked floes to only include those that traveled a total distance of at least a pixel, had average speeds greater than 0.01 m/s and less than 1.5 m/s, and were in a region with sea ice in the NSIDC sea ice concentration dataset. Next, we identify false positives in the floes with 0 sea ice concentration from NSIDC, anomalous floe length scale relative to the sea ice concentration, and with either circularity less than 0.2 or solidity less than 0.4.

We use the scikit-learn logistic regression cross validation function to fit the model. We use 10-fold cross validation. Data is split so that 2/3 of the random sample is used for training and 1/3 for testing. Model metrics:

F1 score: 0.907

Recall: 0.893

Precision: 0.922

## References

1. Buckley, E., Cañuelas, Timmermans, M.-L., and Wilhelmus, M. M. (2023), “Seasonal Evolution of the Sea Ice Floe Size Distribution from Two Decades of MODIS Data,” EGU sphere (preprint), <https://doi.org/10.5194/egusphere-2024-89>
2. Lopez-Acosta, R., Schodlok, M. P., and Wilhelmus, M. M. (2019). “Ice Floe Tracker: An algorithm to automatically retrieve Lagrangian trajectories via feature matching from moderate-resolution visual imagery”, *Remote Sensing of Environment*, 234, 111406, pp. 1-15. DOI:10.1016/j.rse.2019.111406
3. Lopez-Acosta, R. (2021), “Sea Ice Drift in Arctic Marginal Ice Zones Derived from Optical Satellite Imagery”, Doctoral dissertation, University of California Riverside. 162 pages.
4. Manucharyan, G., Lopez-Acosta, R., and Wilhelmus, M. M. (2022), “Spinning ice floes reveal intensification of mesoscale eddies in the western Arctic Ocean”, *Scientific Reports*, 12, 7070, pp. 1-13
5. Meier, W., Fetterer, F., Windnagel, A.K., and Stewart, S. (2021). “NOAA / NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 4”. National Snow and Ice Data Center. Accessed Aug. 17, 2023. URL: <https://doi.org/10.7265/efmz-2t65>
6. Pedregosa et al. (2011), “Scikit-learn: Machine Learning in Python”, *Journal of Machine Learning Research*, 12, pp. 2825-2830, 2011.
7. Van der Walt, S., Schönberger, Johannes L, Nunez-Iglesias, J., Boulogne, Francois, Warner, J. D., Yager, N., ..., Yu, T. (2014). scikit-image: image processing in Python. *PeerJ*, 2, e453.
8. Watkins, D. M., Bliss, A. C., Hutchings, J. K., Wilhelmus, M. M. (2023), “Evidence of abrupt transitions between sea ice dynamical regimes in the East Greenland marginal ice zone”, *Geophysical Research Letters*, 50, e2023GL103558, pp. 1-10

## Contributors

- Daniel Watkins
- Rosalinda Lopez-Acosta
- Monica Martinez Wilhelmus
- Minki Kim
- Ashfaq Ahmed

- Ellen Buckley
- Simon Hatcher