

Problem Set 2 solutions

Question 1: Relativity and Particle Decays

Learning objectives

In this question you will:

- Review relativistic expressions relevant for determining the mass and lifetime of a particle from its decay products
- gain experience in using python to analyze data provided in a text file

1a.

The decays of particles with lifetimes longer than ~ 0.5 ps can be observed in high resolution particle detectors. BY measuring many such decays, properties such as the decaying particle's mass and lifetime can be determined.

The file `decayData.dat` contains a set of simulated observations of particle decays that come from one specific species of hadron, which we designate particle X . The X is observed through its decay $X \rightarrow p\pi^-$ where the p is a proton. All the X particles are produced at the origin ($x = 0, y = 0, z = 0$) but they have with a range of momenta. The position of the decay and the momentum of the proton and π^- are measured.

The following code reads this data file and puts the data into a form that can be easily used in python:



```

In [4]: 1 import math
        2 import numpy as np
        3 import matplotlib.pyplot as plt
        4 from scipy.optimize import curve_fit
        5
        6 # Parse the input file.
        7 file = "decayData.dat"
        8
        9 #Each row corresponds to one event. The columns are:
       10 # x-position of the decay vertex in cm
       11 # y-position of the decay vertex in cm
       12 # z-position of the decay vertex in cm
       13 # species of first particle (always a proton)
       14 # p1x: x-momentum of the proton produced in the decay in GeV
       15 # p1y: y-momentum of the proton produced in the decay in GeV
       16 # p1z: z-momentum of the proton produced in the decay in GeV
       17 # species of second particle (always a pi^-)
       18 # p2x: x-momentum of the pi^- produced in the decay in GeV
       19 # p2y: y-momentum of the pi^- produced in the decay in GeV
       20 # p2z: z-momentum of the pi^- produced in the decay in GeV
       21
       22
       23 inMeta = False
       24 vx = []
       25 vy = []
       26 vz = []
       27 p1x = []
       28 p1y = []
       29 p1z = []
       30 p2x = []
       31 p2y = []
       32 p2z = []
       33
       34
       35 inMeta = True
       36 for line in open(file, "r"):
       37     line = line.strip()
       38     info = line.split(",")
       39     if inMeta and ("<metadata>" in info[0]):
       40         inMeta = True
       41     elif inMeta and ("</metadata>" in info[0]):
       42         inMeta = False
       43     elif not inMeta:
       44         vx.append(float(info[0]))
       45         vy.append(float(info[1]))
       46         vz.append(float(info[2]))
       47         p1x.append(float(info[4]))
       48         p1y.append(float(info[5]))
       49         p1z.append(float(info[6]))
       50         p2x.append(float(info[8]))
       51         p2y.append(float(info[9]))
       52         p2z.append(float(info[10]))
       53 massPiInGeV = 0.13957
       54 massProtonInGeV = 0.93827

```

Verify that all these events correspond to the decay of a particle of a specific species by making a


histogram of the invariant mass of the decays.

In []: 1 PUT ANSWER HERE

Solution:

```
In [17]: 1 particleMass = []
2 E1 = []
3 E2 = []
4 for i in range(len(p1x)):
5     E1.append(math.sqrt(massProtonInGeV**2 + p1x[i]**2 + p1y[i]**2 + p1z[i]**2))
6     E2.append(math.sqrt(massPiInGeV**2 + p2x[i]**2 + p2y[i]**2 + p2z[i]**2))
7     particleMass.append(math.sqrt((E1[i]+E2[i])**2 - (p1x[i]+p2x[i])**2 - (p1y[i]+p2y[i])**2 - (p1z[i]+p2z[i])**2))
8
9 theMass = sum(particleMass)/len(particleMass)
10 print("The Average Mass =", theMass)
11
12 import matplotlib.pyplot as plt
13 plt.hist(particleMass, bins=100, range=(1.00, 1.2), facecolor='green')
14 # Label the x and y axes and add a title
15 plt.xlabel('Mass(pPi) [GeV]')
16 plt.ylabel('Number of Events')
17 plt.title("Measured Invariant Mass")
18 plt.show()
```

The Average Mass = 1.1156460781296806



All the decays have an invariant mass of 1.1155 GeV indicating that they are all consistent with coming from the same species of particle. From the particle data book, you can deduce that the particle is a Λ^0

1b.

Using these data, determine the lifetime of the X particle. What evidence do you have that the X has a decay distribution consistent with a single species with one lifetime?

In []: 1 PUT ANSWER HERE

Solution:

```
In [44]: 1 # The data file gives the position of the decay vertex. We need to turn thi
2 # using the relativistic boost factor gamms
3 properDistance = []
4 for i in range(len(p1x)):
5     dist = math.sqrt(vx[i]**2+vy[i]**2+vz[i]**2)
6     # Relativistic gamma = E/m
7     gamma = (E1[i]+E2[i])/theMass
8     properDistance.append(dist/gamma)
9
10 # If the decay is exponential, the Lifetime (here measured as  $c\tau$  in uni
11 # proper distance
12 ctau = sum(properTime)/len(properTime)
13 print("Lifetime in cm (ctau) = ",ctau)
```

Lifetime in cm (ctau) = 7.985848216543455


Note: This is consistent with the expected lifetime for a Λ^0

```

In [43]: 1 # There are several ways to demonstrate that the decay is consistent with a
          2 #
          3 # First method: plot the decay rate using a log scale for the y-axis. An ex
          4 # Make a histogram of the proper distance
          5 plt.hist(properDistance,bins=15,range=(0,40.0),facecolor='green')
          6 plt.yscale('log')
          7 # Label the x and y axes and add a title
          8 plt.xlabel('Proper Decay Distance [cm]')
          9 plt.ylabel('Number of Events')
         10 plt.show()
         11
         12 print("The distribution is a straight line on a log plot, so it is exponenti

```

Lifetime in cm (ctau) = 7.985848216543455




The distribution is a straight line on a log plot, so it is exponential

```

In [49]: 1 # Second method: Fit the decay distribution to an exponential
2 # Make a histogram of the proper distance
3 (entries,bins,patches) = plt.hist(properDistance,bins=15,range=(0,40.0),face
4 # Label the x and y axes and add a title
5 plt.xlabel('Proper Decay Distance [cm]')
6 plt.ylabel('Number of Events')
7
8 #
9 # We now will fit the histogrammed data to an exponential
10 def expo_model(x,a,b):
11     return a*np.exp(-x/b)
12
13 # Warning: The sqrt of N errors only works if the number of entries in the
14 bincenters = np.array([0.5 * (bins[i] + bins[i+1]) for i in range(len(bins)-
15 binErrors = np.array([max(math.sqrt(entries[i]),1.6) for i in range(len(bins
16 from scipy.optimize import curve_fit
17 popt, pcov = curve_fit(expo_model, xdata=bincenters, ydata=entries, p0=[entr
18
19 # Add the fit function as a curve
20 Y = np.zeros(len(bins))
21 for j in range(len(bins)):
22     Y[j] = expo_model(bins[j],popt[0],popt[1])
23 plt.plot(bins,Y, linewidth=2)
24 plt.show()
25 print("Average proper distance of decays = ",sum(properTime)/len(properTime)
26 print("Fitted lifetime =",popt[1])

```



Average proper distance of decays = 7.985848216543455

Fitted lifetime = 7.8916533553760955

The average proper distance is consistent with what we would expect for a Λ^0 decay. Also, the fitted lifetime is consistent with what we get from a simple average, indicating that a single exponential is pretty good. WARNING: You should NOT use \sqrt{N} as the error bar unless N is large. For this problem, you should really use a binned likelihood fit instead.

Question 2: π^0 Decay

Learning objectives

In this question you will:

- Review basic concepts of Special Relativity and Lorentz Boosts
- Apply these concepts to the case of $\pi^0 \rightarrow \gamma\gamma$ decay
- Learn techniques needed to simulate the decay of an ensemble of π^0 s with non-zero momentum

Adapted from Perkins 4th Edition Problem 1.4

In this problem you will derive an expression for the distribution of photon energies produced in the decays of π^0 s that are moving with fixed momentum. Then, you will learn how to create a simulated sample of such π^0 decays. Note: In this problem, we will use natural units where $\hbar = c = 1$.

2a.

A particle beam consists of π^0 's all with energy E_{lab} and all traveling in the $+z$ direction. Find an expression for the energy of the photons produced from the π^0 decays as a function of m_π , E_{lab} and θ^* (the angle of emission of the photon with respect to the z -axis in the pion rest frame). Using this expression, show that the lab energy spectrum of the photons is flat, extending from $E_{lab} (1 + \beta)/2$ to $E_{lab} (1 - \beta)/2$, where β is the velocity of the π^0 in the lab frame.

Solution:

In the rest frame of the π^0 , the two photons are back-to-back and each have energy $E_1 = E_2 = \frac{m_\pi}{2}$. We can therefore write the momenta of the photons in the π^0 rest frame as:

$$\begin{aligned}\vec{p}_1 &= \frac{m_\pi}{2} [\sin \theta^* \cos \phi^* \hat{x} + \sin \theta^* \sin \phi^* \hat{y} + \cos \theta^* \hat{z}], \\ \vec{p}_2 &= -\vec{p}_1.\end{aligned}$$

We now want to boost to the lab frame. The boost is along the z -direction, so the x and y components of the momentum are not changed by the transform. The energy and z component are transformed:

$$\begin{aligned}E' &= \gamma (E - \beta p_z) \\ p'_x &= p_x \\ p'_y &= p_y \\ p'_z &= \gamma (p_z - \beta E)\end{aligned}$$

Relativity tells us energy of a π^0 in the beam is $E_{lab} = \gamma m_\pi$. The energy of the two photons in the lab frame are therefore:

$$E_1 = \frac{E_{lab}}{m_\pi} \left(\frac{m_\pi}{2} - \beta \frac{m_\pi}{2} \cos \theta^* \right) = \frac{E_{lab}}{2} (1 - \beta \cos \theta^*)$$

$$E_2 = \frac{E_{lab}}{m_\pi} \left(\frac{m_\pi}{2} + \beta \frac{m_\pi}{2} \cos \theta^* \right) = \frac{E_{lab}}{2} (1 + \beta \cos \theta^*)$$

But since the decay of the π^0 is isotropic in its rest frame, the $\cos \theta^*$ distribution is flat between -1 and 1. Therefore from the expression above the distribution of the energy of each of the photons is flat and extends from $E_{lab} (1 + \beta)/2$ to $E_{lab} (1 - \beta)/2$.

2b.

Find an expression for the disparity D (the ratio of the energy of the higher energy photon to the energy of the lower energy) and show that in the relativistic limit $\beta \approx 1$, $D > 3$ in half the decays and $D > 7$ in one quarter

Solution:

Using the expressions for the energy derived above and the definition of D given above:

$$D = \frac{E_{lab} (1 + \beta \cos \theta^*)}{E_{lab} (1 - \beta \cos \theta^*)} = \frac{1 + \beta \cos \theta^*}{1 - \beta \cos \theta^*}$$

where $0 \leq \cos \theta^* \leq 1$ to ensure that the higher energy photon is in the numerator.

We can find the value of $\cos \theta^*$ that corresponds to a given value D of disparity:

$$D = \frac{1 + \beta \cos \theta^*}{1 - \beta \cos \theta^*}$$

$$D - D\beta \cos \theta^* = 1 + \beta \cos \theta^*$$

$$D - 1 = \beta (D + 1) \cos \theta^*$$

$$\cos \theta^* = \frac{D - 1}{\beta (D + 1)} \approx \frac{D - 1}{D + 1}$$

Thus $D = 3$ corresponds to $\cos \theta^* = \frac{2}{4} = \frac{1}{2}$ and $D = 7$ corresponds to $\cos \theta^* = \frac{6}{8} = \frac{3}{4}$.

Since $\cos \theta^*$ is distributed uniformly, we can find the fraction of events with $D > 3$ is $1 - \frac{1}{2} = \frac{1}{2}$ and the fraction of events with $D > 7$ is $1 - \frac{3}{4} = \frac{1}{4}$ in the relativistic limit.

2c.

It is often useful for physicists to simulate experimental data. Such simulations allow us generate an ensemble of events corresponding to a given physical process and to study them. Generated events can be passed through a simulated detector that has imperfections (finite resolution, missing channels, incomplete angular coverage, etc) and the effect of such imperfections on our measurements can be assessed. This problem is our first example of creating such simulated data. Our simulation will be quite simple but the concepts developed here will be used through the semester.

Assume we have a beam of 10000 π^0 all with energy 5 GeV. Simulate the decay of these pions and plot (histogram) the following distributions:

- The energies of the photons produced in the π^0 decay
- The disparity of the decays
- The angles θ between the momenta of the photons and that of the π^0 in the lab frame.

Hints:

- For each decay, first simulate the decay in the pion center of mass and then Lorentz boost to the lab frame
- Since in the rest frame of the pion, the decay is isotropic, the distribution of $\cos \theta^*$ is uniformly distributed. If for each event you pull a random number uniformly distributed between 0 and 1 and set $\cos \theta^*$ for that event equal to the random number, the decays will have the right distribution.
- In principle, you could find the ϕ^* angle for each decay by pulling a second uniformly distributed random number, but for this problem you will not need the x and y components of the photon momentum separately so you don't need to do this.

Solution:

```

In [1]: 1 import numpy as np
2 import matplotlib.pyplot as plt
3 %matplotlib inline
4
5 # set pi^0 mass; use GeV for everything (remember h=c=1)
6 mass = 0.1349770
7
8 def piZeroDecay(piEnergy, nevents):
9     """Compute the lab-frame energies and angles of pi^0 decay photons.
10     Randomly picks cos(Theta*) for the pi^0 decays and then boosts the
11     photons from the pi^0 rest frame to the lab frame.
12
13     Parameters
14     =====
15     piEnergy : float
16         the energy of the pi^0 in the lab frame in GeV
17
18     nevents : int
19         number of decays to simulate
20
21     Returns
22     =====
23     Elab1 : array
24         1-d array of length nevents representing higher-energy photon lab
25         frame energies
26
27     Elab2 : array
28         1-d array of length nevents representing lower-energy photon lab
29         frame energies
30
31     ThetaLab1 : array
32         1-d array of length nevents representing angles (in radians) between
33         higher-energy photon and pi^0 momenta
34
35     ThetaLab2 : array
36         1-d array of length nevents representing angles (in radians) between
37         lower-energy photon and pi^0 momenta
38     """
39
40     # Calculate Lorentz variables gamma and beta
41     gamma = piEnergy/mass
42     beta = (1-gamma**-2)**.5
43
44     # Throw random number to pick Cos(Theta*)
45     # Note: We only bother to throw 0<cos(Theta*)<1 since it makes it easier
46     cosThetastar = np.random.rand(nevents)
47     # Calculate photon energies in lab frame
48     # Note: we have defined things so that photon 1 has the higher energy
49     Elab1 = 0.5*piEnergy*(1+beta*cosThetastar)
50     Elab2 = 0.5*piEnergy*(1-beta*cosThetastar)
51
52     # calculate theta in the Lab frame for each photon. To do this note th
53     # are not affected by the boost. Define pT=sqrt(px^2+py^2)=mass/2 sin(T
54     pT = 0.5 * mass * np.sin(np.arccos(cosThetastar))
55     ThetaLab1 = np.arcsin(pT/Elab1)
56     ThetaLab2 = np.arcsin(pT/Elab2)

```

```

57
58     return Elab1, Elab2, ThetaLab1, ThetaLab2

```


In [2]:

```

1 requestedEnergy = 5 # energy in GeV
2 nevents = 10000 # number of events to simulate
3
4 E1, E2, ThetaLab1, ThetaLab2 = piZeroDecay(requestedEnergy, nevents)
5 gammaEnergy = np.concatenate((E1,E2)) #array of photon energies
6 disparity = E1/E2
7
8 # Fill the histogram. The hist function returns entries per bin, bin bounda
9 # the objects representing the plotted patches
10 n, bins, patches = plt.hist(gammaEnergy,bins=10,range=(0.0,requestedEnergy),f
11
12 # The code below draws error bars with size sqrt(number of events)
13 bin_centers = 0.5*(bins[1:] + bins[:-1])
14 plt.errorbar(bin_centers,n,xerr=None,yerr=n**0.5,fmt='o',ecolor='black',mark
15
16 # Label the x and y axes and add a title
17 plt.xlabel('Photon Energy [GeV]')
18 plt.ylabel('Number of Photons')
19 name = "Photon Energy in Lab Frame for "+str(requestedEnergy)+r" GeV $\pi^0$
20 plt.title(name)


```

Out[2]: Text(0.5, 1.0, 'Photon Energy in Lab Frame for 5 GeV π^0 Decays')



```
In [3]: 1 # Fill the histogram. Note: np.clip() makes the last bin an overflow bin
2 maxRange=50
3 plt.hist(np.clip(disparity,1.0,maxRange),bins=100,range=(1.0,maxRange),facec
4 # Label the x and y axes and add a title
5 plt.xlabel('disparity')
6 plt.ylabel('Number of Events')
7 name = "Disparity for "+str(requestedEnergy)+" GeV  $\pi^0$  Decays"
8 plt.title(name)
```

Out[3]: Text(0.5, 1.0, 'Disparity for 5 GeV π^0 Decays')




```

In [4]: 1 # Fill the histogram. Note: np.clip() makes the last bin an overflow bin
        2
        3 # Convert radians to milli-radians for easier plotting
        4 plt.hist(ThetaLab1*1e3,bins=100,range=(0,300),alpha=0.8,label="high-energy p
        5 plt.hist(ThetaLab2*1e3,bins=100,range=(0,300),alpha=0.8,label="low-energy ph
        6
        7 # Label the x and y axes and add a title
        8 plt.xlabel(r'\theta$ [milli-radian]')
        9 plt.ylabel('Number of Events')
       10 name = r"\theta$ for "+str(requestedEnergy)+r" GeV $\pi^0$ Decays"
       11 plt.title(name)
       12 plt.legend(frameon=False)

```

Out[4]: <matplotlib.legend.Legend at 0x11406f7d0>



2d.

Modify your simulation so that instead of having a fixed energy beam, the π^0 energy is uniformly distributed between 1 and 20 GeV. Make a 2D histogram of the opening angle between the two photons (measured in milli-radians) as a function of the π^0 energy.


Solution:

```

In [5]: 1 bounds = 1,20 #range of pion energies
        2 # This gives a flat distribution of energy between energy bounds
        3 energies = bounds[0] + (bounds[1]-bounds[0])*np.random.rand(nevents)
        4 E1, E2, ThetaLab1, ThetaLab2 = piZeroDecay(energies,nevents)
        5
        6 #since the photons must have opposite phi, the angle between them is the sum
        7 openingAngles = (ThetaLab1+ThetaLab2)*1e3
        8
        9 plt.scatter(energies,openingAngles,s=2) # s controls the size of the points
       10 plt.xlabel('Pion Energy [GeV]')
       11 plt.ylabel('Opening Angle [milli-radian]')
       12 plt.ylim([0,200])
       13 name = r"Opening Angle for %.1f to %.1f GeV  $\pi^0$  Decays"%bounds
       14 plt.title(name)

```

Out[5]: Text(0.5, 1.0, 'Opening Angle for 1.0 to 20.0 GeV π^0 Decays')

**2e.**

In the ATLAS detector, photons are identified in the electromagnetic calorimeter by looking for a narrow energy cluster. Assume that two photons will be *merged* into a single cluster if their opening angles differ by more than 75 milli-radians. Using your scatter plot above, estimate the maximum energy π^0 for which the decay photons can be cleanly separated. (Note: the ATLAS detector is more complicated than the description presented in this problem, having different granularities in θ and ϕ directions. Moreover, the experiment can *identify* π^0 at higher energies than suggested here by looking at the width of the merged energy deposit from the two clusters.)

Solution:

Looking at the plot above, ATLAS can on cleanly separate photons from pions with energy about 3.5 GeV

Question 3: Mandelstam Variables

Type *Markdown* and LaTeX: α^2

3a.

Show that s is the square of the center-of-mass energy of the system 1 + 2

In the center of mass frame $\vec{p}_1 = -\vec{p}_2$. So:

$$\begin{aligned} s &\equiv (p_1 + p_2)^2 \\ &= (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 \\ &= (E_1 + E_2)^2 \end{aligned}$$

Thus, s is the square of the center of mass energy

3b.

Show that

$$s + t + u = m_1^2 + m_2^2 + m_3^2 + m_4^2$$

$$\begin{aligned} s + t + u &= (p_1 + p_2)^2 + (p_1 - p_3)^2 + (p_1 - p_4)^2 \\ &= p_1^2 + p_2^2 + 2p_1 \cdot p_2 + p_1^2 + p_3^2 - 2p_1 \cdot p_3 + p_1^2 + p_4^2 - 2p_1 \cdot p_4 \\ &= 3p_1^2 + p_2^2 + p_3^2 + p_4^2 + 2p_1 \cdot (p_2 - p_3 - p_4) \\ &= 3p_1^2 + p_2^2 + p_3^2 + p_4^2 - 2p_1^2 \\ &= p_1^2 + p_2^2 + p_3^2 + p_4^2 \\ &= m_1^2 + m_2^2 + m_3^2 + m_4^2 \end{aligned}$$

Question 4: β -decay and the uncertainty principle

$$\begin{aligned} \Delta x \Delta p &\geq \hbar \\ \Delta p &\geq \frac{\hbar}{\Delta x} \\ \Delta pc &\geq \frac{\hbar c}{\Delta x} \\ &\geq \frac{197 \text{ MeV fm}}{1 \text{ fm}} \\ &\geq 197 \text{ MeV} \end{aligned}$$

The endpoint of the tritium β -decay spectrum is only about 18 keV, much smaller than the value obtained from the uncertainty principle. Thus, the electron cannot be bound in the nucleus and must be produced in a decay

Question 5: Kinematics in 2-body particle decays

$$\begin{aligned}
 m_a^2 &= E_a^2 - \vec{p}_a^2 \\
 &= (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 \\
 &= E_1^2 + E_2^2 + 2E_1 E_2 - (p_1^2 + p_2^2 + 2\vec{p}_1 \cdot \vec{p}_2) \\
 &= E_1^2 - \vec{p}_1^2 + E_2^2 - \vec{p}_2^2 + 2E_1 E_2 \left(1 - \frac{|\vec{p}_1||\vec{p}_2|}{E_1 E_2} \cos \theta \right) \\
 &= E_1^2 - \vec{p}_1^2 + E_2^2 - \vec{p}_2^2 + 2E_1 E_2 (1 - \beta_1 \beta_2 \cos \theta)
 \end{aligned}$$

In []:

1