

# From Rainbows to Resurgence: Asymptotics of the Airy Function


Will Hoffer

University of California, Riverside

*math@willhoffer.com*

December 14, 2020

# Supernumerary, or Spurious, Rainbow



# Historical Preface

- In 1838, G. B. Airy found a theoretical model for the illumination, involving (the square of) the integral:

$$W(m) = \int_0^{\infty} \cos\left(\frac{\pi}{2}(w^3 - mw)\right) dw$$

# Historical Preface

- In 1838, G. B. Airy found a theoretical model for the illumination, involving (the square of) the integral:

$$W(m) = \int_0^{\infty} \cos\left(\frac{\pi}{2}(w^3 - mw)\right) dw$$

- Airy could only calculate two zeroes before the process became intractable

# Historical Preface

- In 1838, G. B. Airy found a theoretical model for the illumination, involving (the square of) the integral:

$$W(m) = \int_0^{\infty} \cos\left(\frac{\pi}{2}(w^3 - mw)\right) dw$$

- Airy could only calculate two zeroes before the process became intractable
- In 1841, W. H. Miller observed/measured 30 dark bands for the primary bow

# Historical Preface

- In 1838, G. B. Airy found a theoretical model for the illumination, involving (the square of) the integral:

$$W(m) = \int_0^{\infty} \cos\left(\frac{\pi}{2}(w^3 - mw)\right) dw$$

- Airy could only calculate two zeroes before the process became intractable
- In 1841, W. H. Miller observed/measured 30 dark bands for the primary bow
- In 1850, Stokes employed another method, managing to calculate 50 zeroes!

# Stokes' Method I

After some rescaling/changing variables, Stokes worked with:

$$u = e^{-i\frac{\pi}{6}} \int_0^\infty e^{-(z^3 - pz)} dx = U - iU'$$

# Stokes' Method I

After some rescaling/changing variables, Stokes worked with:

$$u = e^{-i\frac{\pi}{6}} \int_0^\infty e^{-(z^3 - pz)} dx = U - iU'$$

$U$  satisfies the Airy equation:

$$\frac{d^2U}{dn^2} + \frac{n}{3}U = 0$$

# Stokes' Method I

After some rescaling/changing variables, Stokes worked with:

$$u = e^{-i\frac{\pi}{6}} \int_0^\infty e^{-(z^3 - pz)} dx = U - iU'$$

$U$  satisfies the Airy equation:

$$\frac{d^2U}{dn^2} + \frac{n}{3}U = 0$$

Airy's method was to expand the exponential in  $pz$ .

# Stokes' Method I

After some rescaling/changing variables, Stokes worked with:

$$u = e^{-i\frac{\pi}{6}} \int_0^\infty e^{-(z^3 - pz)} dx = U - iU'$$

$U$  satisfies the Airy equation:

$$\frac{d^2U}{dn^2} + \frac{n}{3}U = 0$$

Airy's method was to expand the exponential in  $pz$ .

As noted by Stokes, though, this is “not convenient” when  $n$  becomes large.

## Stokes' Method II

Stokes had the idea to write:<sup>1</sup>

$$U = e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( An^\alpha + Bn^\beta + Cn^\gamma + \dots \right)$$

---

<sup>1</sup>Based on formulae M. Cauchy had found for Fresnel's integrals.

## Stokes' Method II

Stokes had the idea to write:<sup>1</sup>

$$U = e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( An^\alpha + Bn^\beta + Cn^\gamma + \dots \right)$$

From the equation, he deduced that:

$$U = An^{-\frac{1}{4}}e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( 1 - \frac{1 \cdot 5 \cdot i}{16\sqrt{3n^2}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2} \left( \frac{i}{16\sqrt{3n^2}} \right)^2 + \dots \right)$$

and similarly for negative values  $n' = -n$ .

---

<sup>1</sup>Based on formulae M. Cauchy had found for Fresnel's integrals.

## Stokes' Method II

Stokes had the idea to write:<sup>1</sup>

$$U = e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( An^\alpha + Bn^\beta + Cn^\gamma + \dots \right)$$

From the equation, he deduced that:

$$U = An^{-\frac{1}{4}}e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( 1 - \frac{1 \cdot 5 \cdot i}{16\sqrt{3n^2}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2} \left( \frac{i}{16\sqrt{3n^2}} \right)^2 + \dots \right)$$

and similarly for negative values  $n' = -n$ .

Just one small problem...

---

<sup>1</sup>Based on formulae M. Cauchy had found for Fresnel's integrals.

## Stokes' Method II

Stokes had the idea to write:<sup>1</sup>

$$U = e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( An^\alpha + Bn^\beta + Cn^\gamma + \dots \right)$$

From the equation, he deduced that:

$$U = An^{-\frac{1}{4}}e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left( 1 - \frac{1 \cdot 5 \cdot i}{16\sqrt{3n^2}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2} \left( \frac{i}{16\sqrt{3n^2}} \right)^2 + \dots \right)$$

and similarly for negative values  $n' = -n$ .

Just one small problem... the series diverges.

---

<sup>1</sup>Based on formulae M. Cauchy had found for Fresnel's integrals. A set of small, light-blue navigation icons typically used in Beamer presentations for navigating between slides and sections.

## Stokes' Method III

Here's what Stokes' original series looks like:

$$U = An^{-\frac{1}{4}} e^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left\{ 1 - \frac{1 \cdot 5}{1} \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2} \left( \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} \right)^2 \right. \\ \left. - \frac{1 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17}{1 \cdot 2 \cdot 3} \left( \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} \right)^3 + \dots \right\} \dots \quad (14)$$

# Stokes' Method III

Here's what Stokes' original series looks like:

$$U = An^{-\frac{1}{4}} \epsilon^{\frac{2}{3}\sqrt{-\frac{n^3}{3}}} \left\{ 1 - \frac{1 \cdot 5}{1} \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2} \left( \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} \right)^2 \right. \\ \left. - \frac{1 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17}{1 \cdot 2 \cdot 3} \left( \frac{\sqrt{-1}}{16 \sqrt{(3n^3)}} \right)^3 + \dots \right\} \dots \quad (14)$$

Secondly, suppose  $n$  negative, and equal to  $-n'$ . Then, writing  $-n'$  for  $n$  in (14), and changing the arbitrary constant, and the sign of the radical, we get

$$U = Cn'^{-\frac{1}{4}} \epsilon^{-\frac{2}{3}\sqrt{\frac{n'^3}{3}}} \left\{ 1 - \frac{1 \cdot 5}{1 \cdot 16 (3n^3)^{\frac{1}{2}}} + \frac{1 \cdot 5 \cdot 7 \cdot 11}{1 \cdot 2 \cdot 16 \cdot 3n^3} - \dots \right\} \dots \quad (17)$$

# Stokes' Method IV

G. G. Stokes

When  $n$  or  $n'$  is at all large, the series [for  $U$ ] are at first rapidly convergent, but they are ultimately in all cases hypergeometrically divergent. Notwithstanding this divergence, we may employ the series in numerical calculation, provided we do not take in the divergent terms.

# Stokes' Method IV

## G. G. Stokes

When  $n$  or  $n'$  is at all large, the series [for  $U$ ] are at first rapidly convergent, but they are ultimately in all cases hypergeometrically divergent. Notwithstanding this divergence, we may employ the series in numerical calculation, provided we do not take in the divergent terms.

Stokes was able to calculate 50 zeroes, encountering “only” 7-figure logarithms instead of 10-figure ones.

# Stokes' Method IV

## G. G. Stokes

When  $n$  or  $n'$  is at all large, the series [for  $U$ ] are at first rapidly convergent, but they are ultimately in all cases hypergeometrically divergent. Notwithstanding this divergence, we may employ the series in numerical calculation, provided we do not take in the divergent terms.

Stokes was able to calculate 50 zeroes, encountering “only” 7-figure logarithms instead of 10-figure ones.

This is exactly what in modern days we would deem optimal (or least) truncation of an asymptotic expansion.

## Stokes Method V

The expansion for  $U$  is different for negative  $n$  (what we now call the first example of the Stokes phenomenon.)

## Stokes Method V

The expansion for  $U$  is different for negative  $n$  (what we now call the first example of the Stokes phenomenon.)

This expansion was less tractable:

G. G. Stokes

This mode of determining the constant [in front of the negative  $U$  expansion] is anything but satisfactory. I have endeavored in vain to deduce the leading term in  $U$  for  $n$  negative from the integral itself...

## Stokes Method V

The expansion for  $U$  is different for negative  $n$  (what we now call the first example of the Stokes phenomenon.)

This expansion was less tractable:


G. G. Stokes

This mode of determining the constant [in front of the negative  $U$  expansion] is anything but satisfactory. I have endeavored in vain to deduce the leading term in  $U$  for  $n$  negative from the integral itself...

Today, we do know how to deduce the behavior for all  $n$  from the integral itself!



# The Airy Function (Real Inputs)

$$\text{Ai}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(xt + \frac{t^3}{3})} dt$$



# The Airy Function (Real Inputs)

$$\text{Ai}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(xt + \frac{t^3}{3})} dt$$



# Method of Steepest Descent

Defining the Airy function for complex  $z$  takes some care; the integral is highly oscillatory.

# Method of Steepest Descent

Defining the Airy function for complex  $z$  takes some care; the integral is highly oscillatory.

The method of steepest descent for an integral of the form:

$$\int_C f(t) e^{xg(t)} dt, \quad x \gg 1$$

involves deforming the integration contour to pass along the direction of steepest descent (i.e. parallel to  $-\nabla u$ , where  $g = u + iv$ ) to pass by saddle points (viz. near where the integral is maximal, or at least less rapid oscillations cause less cancellation.)

# Method of Steepest Descent

Defining the Airy function for complex  $z$  takes some care; the integral is highly oscillatory.

The method of steepest descent for an integral of the form:

$$\int_C f(t) e^{xg(t)} dt, \quad x \gg 1$$

involves deforming the integration contour to pass along the direction of steepest descent (i.e. parallel to  $-\nabla u$ , where  $g = u + iv$ ) to pass by saddle points (viz. near where the integral is maximal, or at least less rapid oscillations cause less cancellation.)

(In particular, this enables one to use Laplace's method to estimate the integral.)

# Airy Function, Deformed Contour

For the Airy integral, these curves occur when the cubic term is negative. In particular, this happens if the phase of  $t$  is  $\pm \frac{i\pi}{3}$ , or  $t$  is negative.

# Airy Function, Deformed Contour

For the Airy integral, these curves occur when the cubic term is negative. In particular, this happens if the phase of  $t$  is  $\pm \frac{i\pi}{3}$ , or  $t$  is negative.

Thus, we write:

$$\text{Ai}(z) = \frac{1}{2\pi i} \int_{\gamma} e^{\frac{t^3}{3} - zt} dt, \quad \text{Im}(\gamma) = (\infty e^{-\frac{\pi i}{3}}, 0] \cup [0, \infty e^{\frac{\pi i}{3}})$$

# Airy Function, Deformed Contour

For the Airy integral, these curves occur when the cubic term is negative. In particular, this happens if the phase of  $t$  is  $\pm \frac{i\pi}{3}$ , or  $t$  is negative.

Thus, we write:

$$\begin{aligned}\text{Ai}(z) &= \frac{1}{2\pi i} \int_{\gamma} e^{\frac{t^3}{3} - zt} dt, & \text{Im}(\gamma) &= (\infty e^{-\frac{\pi i}{3}}, 0] \cup [0, \infty e^{\frac{\pi i}{3}}) \\ &= \frac{1}{2\pi i} \int_{\infty e^{-\frac{i\pi}{3}}}^{\infty e^{\frac{i\pi}{3}}} e^{\frac{t^3}{3} - zt} dt\end{aligned}$$

# Airy Function Asymptotics

Asymptotic Expansions:

$$\text{Ai}(z) \sim \frac{1}{2\sqrt{\pi}} z^{-\frac{1}{4}} e^{-\zeta} \sum_{n=0}^{\infty} (-1)^n \frac{c_n}{\zeta^n} \quad |\arg(z)| < \pi$$

# Airy Function Asymptotics

Asymptotic Expansions:

$$\text{Ai}(z) \sim \frac{1}{2\sqrt{\pi}} z^{-\frac{1}{4}} e^{-\zeta} \sum_{n=0}^{\infty} (-1)^n \frac{c_n}{\zeta^n} \quad |\arg(z)| < \pi$$

$$\begin{aligned} \text{Ai}(-z) \sim & \frac{1}{\sqrt{\pi}} z^{-\frac{1}{4}} \left( \sin(\zeta + \frac{\pi}{4}) \sum_{n=0}^{\infty} (-1)^n \frac{c_{2n}}{\zeta^{2n}} \right. \\ & \left. - \cos(\zeta + \frac{\pi}{4}) \sum_{n=0}^{\infty} (-1)^n \frac{c_{2n+1}}{\zeta^{2n+1}} \right) \quad |\arg(z)| < \frac{2}{3}\pi \end{aligned}$$

# Airy Function Asymptotics

Asymptotic Expansions:

$$\text{Ai}(z) \sim \frac{1}{2\sqrt{\pi}} z^{-\frac{1}{4}} e^{-\zeta} \sum_{n=0}^{\infty} (-1)^n \frac{c_n}{\zeta^n} \quad |\arg(z)| < \pi$$

$$\begin{aligned} \text{Ai}(-z) \sim & \frac{1}{\sqrt{\pi}} z^{-\frac{1}{4}} \left( \sin(\zeta + \frac{\pi}{4}) \sum_{n=0}^{\infty} (-1)^n \frac{c_{2n}}{\zeta^{2n}} \right. \\ & \left. - \cos(\zeta + \frac{\pi}{4}) \sum_{n=0}^{\infty} (-1)^n \frac{c_{2n+1}}{\zeta^{2n+1}} \right) \quad |\arg(z)| < \frac{2}{3}\pi \end{aligned}$$

Notation:


$$\zeta = \frac{2}{3} z^{\frac{3}{2}}, \quad c_0 = 1, \quad c_n = \frac{\Gamma(3n + \frac{1}{2})}{54^n \Gamma(n+1) \Gamma(n + \frac{1}{2})}$$

# Asymptotics on the Real Line

$$\text{Ai}(x) \sim \begin{cases} \frac{1}{2\sqrt{\pi}} x^{-\frac{1}{4}} e^{-\frac{2}{3}x^{\frac{3}{2}}} & x > 0 \\ \frac{1}{\sqrt{\pi}} (-x)^{-\frac{1}{4}} \sin\left(\frac{2}{3}(-x)^{\frac{3}{2}} + \frac{\pi}{4}\right) & x < 0 \end{cases}$$

# Asymptotics on the Real Line

$$\text{Ai}(x) \sim \begin{cases} \frac{1}{2\sqrt{\pi}} x^{-\frac{1}{4}} e^{-\frac{2}{3}x^{\frac{3}{2}}} & x > 0 \\ \frac{1}{\sqrt{\pi}} (-x)^{-\frac{1}{4}} \sin\left(\frac{2}{3}(-x)^{\frac{3}{2}} + \frac{\pi}{4}\right) & x < 0 \end{cases}$$




# Asymptotics in the Complex Plane

Complex plots of the approximations and where they agree.

$$\text{Ai}(z)$$

(Entire)




# Asymptotics in the Complex Plane

Complex plots of the approximations and where they agree.


$$\text{Ai}(z)$$

(Entire)



$$\frac{z^{-\frac{1}{4}}}{2\sqrt{\pi}} e^{-\frac{2}{3}z^{\frac{3}{2}}}$$

$$|\arg(z)| < \pi$$





# Asymptotics in the Complex Plane

Complex plots of the approximations and where they agree.

$$\frac{(-z)^{-\frac{1}{4}}}{\sqrt{\pi}} \sin \left( \frac{2}{3}(-z)^{\frac{3}{2}} + \frac{\pi}{4} \right) \quad \text{Ai}(z)$$


$$| \arg(-z) | < \frac{2\pi}{3}$$



(Entire)

$$\frac{z^{-\frac{1}{4}}}{2\sqrt{\pi}} e^{-\frac{2}{3}z^{\frac{3}{2}}}$$

$$| \arg(z) | < \pi$$



# Resumming the Asymptotic Series

The Airy function is governed by the asymptotic expansion:

$$\varphi_{\text{Ai}}(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^n} = \sum_{n=0}^{\infty} \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} \frac{1}{z^n}$$

# Resumming the Asymptotic Series

The Airy function is governed by the asymptotic expansion:

$$\varphi_{\text{Ai}}(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^n} = \sum_{n=0}^{\infty} \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} \frac{1}{z^n}$$

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

# Resumming the Asymptotic Series

The Airy function is governed by the asymptotic expansion:

$$\varphi_{\text{Ai}}(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^n} = \sum_{n=0}^{\infty} \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} \frac{1}{z^n}$$

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

(This is the same formula as before, but with a change of variables to make growth akin to  $\Gamma(n)$  more manifest.)

$$a_n = \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} = \left(-\frac{2}{3}\right)^{-n} \frac{\Gamma(3n + \frac{1}{2})}{54^n\Gamma(n+1)\Gamma(n + \frac{1}{2})}$$

# Resumming the Asymptotic Series

The Airy function is governed by the asymptotic expansion:

$$\varphi_{\text{Ai}}(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^n} = \sum_{n=0}^{\infty} \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} \frac{1}{z^n}$$

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

More remarks:

- $\varphi_{\text{Ai}}$  is factorially divergent (of Gevrey-class one.)

# Resumming the Asymptotic Series

The Airy function is governed by the asymptotic expansion:


$$\varphi_{\text{Ai}}(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^n} = \sum_{n=0}^{\infty} \left(-\frac{3}{4}\right)^n \frac{\Gamma(n + \frac{1}{6})\Gamma(n + \frac{5}{6})}{2\pi\Gamma(n+1)} \frac{1}{z^n}$$

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$


More remarks:

- $\varphi_{\text{Ai}}$  is factorially divergent (of Gevrey-class one.)
- $z = k^{\frac{3}{2}}$  is a natural change of variables for ensuing resummation.

# Interlude: Borel Summation Schematic



# Borel Summation Example



# Airy Series: Borel Summation

- The minor of  $\varphi_{\text{Ai}}$  is its (formal) Borel transform, forgetting the constant term:

$$\tilde{\varphi}_{\text{Ai}} := \mathcal{B}[\varphi_{\text{Ai}}] = \sum_{n=1}^{\infty} a_n \frac{\zeta^{n-1}}{(n-1)!}$$

# Airy Series: Borel Summation

- The minor of  $\varphi_{\text{Ai}}$  is its (formal) Borel transform, forgetting the constant term:

$$\tilde{\varphi}_{\text{Ai}} := \mathcal{B}[\varphi_{\text{Ai}}] = \sum_{n=1}^{\infty} a_n \frac{\zeta^{n-1}}{(n-1)!}$$

- $\tilde{\varphi}_{\text{Ai}}$  extends analytically to the universal cover of  $\mathbb{C} \setminus \{0, -\frac{4}{3}\}$

# Airy Series: Borel Summation

- The minor of  $\varphi_{\text{Ai}}$  is its (formal) Borel transform, forgetting the constant term:

$$\tilde{\varphi}_{\text{Ai}} := \mathcal{B}[\varphi_{\text{Ai}}] = \sum_{n=1}^{\infty} a_n \frac{\zeta^{n-1}}{(n-1)!}$$

- $\tilde{\varphi}_{\text{Ai}}$  extends analytically to the universal cover of  $\mathbb{C} \setminus \{0, -\frac{4}{3}\}$
- For any direction  $\theta$  not along the negative real axis, the following converges for  $\text{Re}(ze^{i\theta}) > 0$ :

$$S_{\theta}\varphi_{\text{Ai}}(z) := a_0 + \mathcal{L}_{\theta}\mathcal{B}[\varphi_{\text{Ai}}](z) = a_0 + \int_0^{\infty e^{i\theta}} \tilde{\varphi}_{\text{Ai}}(\zeta) e^{-z\zeta} d\zeta$$

# A Borel Resummed Expansion

Where before:

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

We now have:

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} S_0 \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

# A Borel Resummed Expansion

Where before:

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

We now have:

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} S_0 \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

This resummation is valid for  $|\arg(k)| < \frac{\pi}{3}$ ,  $|k| > 0$ .

# A Borel Resummed Expansion

Where before:

$$\text{Ai}(k) \sim \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

We now have:

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} e^{-\frac{2}{3}k^{\frac{3}{2}}} S_0 \varphi_{\text{Ai}}(k^{\frac{3}{2}})$$

This resummation is valid for  $|\arg(k)| < \frac{\pi}{3}$ ,  $|k| > 0$ .

One can rotate the direction of summation for new regions of validity.

# Contours near the Singularity at $-\frac{4}{3}$

Rotation of summation is fine up until one encounters the singularity on the negative real line.

# Contours near the Singularity at $-\frac{4}{3}$

Rotation of summation is fine up until one encounters the singularity on the negative real line.




FIGURE 2. Right and left Borel-resummation.

One can compare right and left-resummations, since

$$(4) \quad S_{-\pi^-} \varphi_{Ai}(z) = S_{-\pi^+} \varphi_{Ai}(z) + \int_{\gamma} \widetilde{\varphi_{Ai}}(\zeta) e^{-z\zeta} d\zeta$$

# “Alien” Calculus & Behavior across the Singularity

The Hankel contour  $\gamma$  can be expressed using the so-called alien derivative:

$$\int_{\gamma} \tilde{\varphi}_{\text{Ai}}(\zeta) e^{-z\zeta} d\zeta = e^{+\frac{4}{3}z} S_{-\pi} \left( \Delta_{-\frac{4}{3}}^z \varphi_{\text{Ai}} \right) (z)$$

# “Alien” Calculus & Behavior across the Singularity

The Hankel contour  $\gamma$  can be expressed using the so-called alien derivative:

$$\int_{\gamma} \tilde{\varphi}_{\text{Ai}}(\zeta) e^{-z\zeta} d\zeta = e^{+\frac{4}{3}z} S_{-\pi} \left( \Delta_{-\frac{4}{3}}^z \varphi_{\text{Ai}} \right) (z)$$

In this case,

$$\Delta_{-\frac{4}{3}}^z \varphi_{\text{Ai}} = -i\varphi_{\text{Bi}}, \quad \varphi_{\text{Bi}}(z) := \sum_{n=0}^{\infty} (-1)^n \frac{a_n}{z^n}$$

# “Alien” Calculus & Behavior across the Singularity

The Hankel contour  $\gamma$  can be expressed using the so-called alien derivative:

$$\int_{\gamma} \tilde{\varphi}_{\text{Ai}}(\zeta) e^{-z\zeta} d\zeta = e^{+\frac{4}{3}z} S_{-\pi} \left( \Delta_{-\frac{4}{3}}^z \varphi_{\text{Ai}} \right) (z)$$

In this case,

$$\Delta_{-\frac{4}{3}}^z \varphi_{\text{Ai}} = -i\varphi_{\text{Bi}}, \quad \varphi_{\text{Bi}}(z) := \sum_{n=0}^{\infty} (-1)^n \frac{a_n}{z^n}$$

$\varphi_{\text{Bi}}$  is also Gevrey-1 and its minor  $\tilde{\varphi}_{\text{Bi}}$  extends analytically to the universal cover of  $\mathbb{C} \setminus \{0, +\frac{4}{3}\}$ .

# Airy Function on the Negative Real Line

Deducing the behavior  $\text{Ai}$  for negative real inputs.

# Airy Function on the Negative Real Line

Deducing the behavior  $\text{Ai}$  for negative real inputs.

Airy expansion when  $|\arg(k) - \pi| < \frac{\pi}{3}$ ,  $z = k^{\frac{3}{2}}$ :

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} \left( e^{-\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Ai}}(z) + ie^{+\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Bi}}(z) \right)$$

# Airy Function on the Negative Real Line

Deducing the behavior  $\text{Ai}$  for negative real inputs.

Airy expansion when  $|\arg(k) - \pi| < \frac{\pi}{3}$ ,  $z = k^{\frac{3}{2}}$ :

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} \left( e^{-\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Ai}}(z) + ie^{+\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Bi}}(z) \right)$$

Note the new exponential term that appeared.

# Airy Function on the Negative Real Line

Deducing the behavior  $\text{Ai}$  for negative real inputs.

Airy expansion when  $|\arg(k) - \pi| < \frac{\pi}{3}$ ,  $z = k^{\frac{3}{2}}$ :

$$\text{Ai}(k) = \frac{1}{2\sqrt{\pi}} k^{-\frac{1}{4}} \left( e^{-\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Ai}}(z) + ie^{+\frac{2}{3}z} S_{-\frac{3\pi}{2}} \varphi_{\text{Bi}}(z) \right)$$

Note the new exponential term that appeared.

Once can rewrite the LHS as the resummed version of the second expansion we saw previously.

## Zeroes “Resurge” from the Original Expansion

- This expression can be employed to calculate the zeroes on the Airy function (viz. the Delabaere reference.)

## Zeroes “Resurge” from the Original Expansion

- This expression can be employed to calculate the zeroes on the Airy function (viz. the Delabaere reference.)
- This procedure follows the original calculations of Stokes, but now by manipulating the full series itself. The mathematical justification for the resummation comes from resurgent analysis.

## Zeroes “Resurge” from the Original Expansion

- This expression can be employed to calculate the zeroes on the Airy function (viz. the Delabaere reference.)
- This procedure follows the original calculations of Stokes, but now by manipulating the full series itself. The mathematical justification for the resummation comes from resurgent analysis.
- In particular, the behavior on the negative real line is manifestly contained in the expansion on the positive real line— an example of resurgence.

# Historical References

- G. B. Airy, “On the Intensity of Light in the neighbourhood of a Caustic,” Trans. Cambridge Phil. Soc. Vol. 6, Pt. 3, 397-402 (1838)
- W. H. Miller “On Spurious Rainbows,” Trans. Cambridge Phil. Soc. Vol. 7, Pt. 3, 277-286 (1841)
- G. G. Stokes, “On the numerical Calculation of a Class of Definite Integrals and Infinite Series,” Trans. Cambridge Phil. Soc. Vol. 9 Pt. 2, 166-187 (1850)
- G. G. Stokes, “On the discontinuity of arbitrary constants that appear as multipliers of semi-convergent series (A letter to the Editor),” Acta Math. Stockholm 26, 393-397 (1902)

# Journal References

- A. B. Olde Daalhuis, S. J. Chapman, J. R. King, J. R. Ockendon, and R. H. Tew, “Stokes phenomenon and matched asymptotic expansions,” SIAM J. Appl. Math, Vol. 55, No. 6, 1469-1483 (1995)
- E. Delabaere, “Effective Resummation Methods for an Implicit Resurgent Function,” arXiv:math-ph/0602026.
- M. V. Berry and C. J. Howls, “Hyperasymptotics for integrals with saddles,” Proc. R. Soc. Lond. A 434, 657-675 (1991)
- O. Costin and M. Huang, “Behavior of Lacunary Series at the Natural Boundary,” Advances in Mathematics 222(4):1370-1404, DOI: 10.1016/j.aim.2009.06.011

# Book References

- Abramowitz, M. and Stegun, I. A., eds. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, "Chapter 10," Applied Mathematics Series, 55, (Ninth reprint, 1983). Washington D.C.; New York: United States Department of Commerce, National Bureau of Standards; Dover Publications. p. 446-447
- O. Costin, *Asymptotics and Borel Summability*, Chapman & Hall/CRC Press, 2009.

# Images

- Mika-Pekka Markkanen, Supernumerary Rainbows [Photograph], 23 May 2010, Wikimedia Commons. License: CC BY-SA 4.0.
- Les Cowley, Bow from 0.75mm diameter drops illuminated by a distant point source [Simulated Image, Cropped], n.d., Atmospheric Optics, attopics.co.uk. Copyright of Les Cowley.
- Geek3, Airy Ai Asymptotic [SVG, converted to PNG], 7 Feb. 2015, Wikimedia Commons. License: CC BY 3.0.