STA 610 LAB 3

William Tirone

2023-09-06

EDA

First, we notice that there are some null values that present issues with fitting the model, so we remove them. In adddition, after plotting a density of the net income, we see a very large range of values, so I chose to scale them. We can't take the log because there are some negative net income values.

```
hc2014 = hc2014 >
    drop_na() |>
    mutate(scaled_income = scale(netincome))
  agg = hc2014 |>
    group_by(control) |>
    summarise(total_beds = sum(numbeds)) |>
    arrange(desc(total_beds))
  agg[0:10,]
# A tibble: 10 x 2
  control
                          total_beds
  <fct>
                                <dbl>
1 Nonprofit-other
                              383324
2 Corporation
                              166482
3 Nonprofit-church
                              104345
4 Gvmt-State
                               46498
5 Gvmt-County
                                31339
6 Gvmt-Hospital District
                               27978
7 Gvmt-Other
                                12480
8 Partnership
                                10317
```


```
9 Gvmt-City-County 9225
10 Gvmt-City 7935
```

We see a very sharp spike at 0, but also some pretty large outliers.

```
ggplot(hc2014, aes(scaled_income)) +
  geom_density(aes(y=..density..),color="black",linetype="dashed") +
  theme(legend.position="none") +
  geom_density(alpha=.25, fill="navy") +
  labs(title="Scaled Income of Hospitals in the U.S.",y="net profit") +
  theme_classic()
```


Warning: The dot-dot notation (`..density..`) was deprecated in ggplot2 3.4.0. i Please use `after_stat(density)` instead.

Scaled Income of Hospitals in the U.S.




Below, it looks like with a random sample of states, we have a pretty significant spread among them. New York has some hospitals with pretty large negative scaled profits and Texas has some with very positive values.

```
set.seed(1000)
sample_state <- sample(unique(hc2014$state),15,replace=F)</pre>
```



And considering control, there's large variability in the Nonprofit-church ownership. It's interesting that Nonprofit-other has the largest number of beds by far, but they have much less variability.



Modeling

Our final model will be:

$$y_{ij} = \mu_0 + \alpha_j + \beta_1 * \text{num beds} + \beta_2 * \text{control} + \epsilon_{ij}$$

We will fit 2 models and compare them, a standard linear regression and a model with a random effect.

Neither of the random slope models we tried to fit in lab converged or would run. In particular:

```
model3 = lmer(netincome ~ numbeds + control + (1 + control | state), data=hc2014)
model3 = lmer(netincome ~ numbeds + control + (numbeds + control | state),
data=hc2014)
```

```
model1 = lm(scaled_income ~ numbeds + control, data=hc2014) # standard
model2 = lmer(netincome ~ numbeds + control + (1 | state), data=hc2014) # random effect
lrtest(model2, model1)
```

Warning in modelUpdate(objects[[i - 1]], objects[[i]]): original model was of class "lmerMod", updated model is of class "lm"

Likelihood ratio test

```
Model 1: netincome ~ numbeds + control + (1 | state)

Model 2: scaled_income ~ numbeds + control

#Df LogLik Df Chisq Pr(>Chisq)

1 16 -121645

2 15 -8615 -1 226060 < 2.2e-16 ***
---

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Because we observe a p-value < 0.05, we will reject the null hypothesis that both models fit the data equally well. Thus, we conclude that we want to keep the model with the random effect rather than the simple linear model.

Now, we can look at the fixed effects coefficients with their respective standard errors:

```
coef(summary(model2))
```


	Estimate	Std. Error	t value
(Intercept)	7259876.9	4003254.696	1.8134936
numbeds	101315.4	7044.451	14.3822952
controlNonprofit-other	-13184254.7	4214050.326	-3.1286420
controlIndividual	-10538126.1	24209263.186	-0.4352931
controlCorporation	-11435963.7	4496205.262	-2.5434701
controlPartnership	-6748597.7	7848296.520	-0.8598806
controlOther Proprietary	-8348467.1	9678452.271	-0.8625829
controlGvmt-Federal	-86469599.4	14860200.757	-5.8188715
controlGvmt-City-County	-14558699.7	10110371.438	-1.4399767
controlGvmt-County	-16813418.3	5935319.241	-2.8327741
controlGvmt-State	-37104410.6	7075197.400	-5.2442933
controlGvmt-Hospital District	-13399278.6	6249812.076	-2.1439490
controlGvmt-City	-18180789.2	11714552.152	-1.5519833
controlGvmt-Other	-1627110.7	11497265.451	-0.1415215

And we can view the random intercept and each intercept's respective uncertainty (that is, if I'm understanding this plot correctly).

```
dotplot(ranef(model2, condVar=TRUE))
```

\$state

state



Model Assessment

As a quick check, our residuals don't show any obvious patterns.

plot(model2)

