

Methods 6

Will Tirone

Q1)

Here, we'll just multiply the successive probabilities of being selected together.

$$\pi_{ij} = \frac{nx_i}{t_x} \frac{m_i x_{ij}}{x_i} \frac{q_{ij}}{x_{ij}} = \frac{nmq_{ij}}{t_x}$$

Q2)

First, we have to create unique values for the ppsu's, we'll do this by multiplying by 1000 and adding the ppsu's.

```
ncvs = ncvs |>
  mutate(ppsu_unique = pstrat * 1000 + ppsu)

ncvs_survey = svydesign(~ppsu_unique, strata=~pstrat, weight=~pweight, data=ncvs)
```

a)

Here, I checked the codebook and `numinc` is the number of incidents per person, so we just use the usual code. The total is 15,850,360.

```
svytotal(~numinc, ncvs_survey)
```

	total	SE
numinc	15850360	492124

b)

Same thing here, just using `medexp` for medical expenses. The total is \$86,285,440

```
svytotal(~medexp, ncvs_survey)
```

	total	SE
medexp	86285440	28354406

c)

And a similar thing here but with robbery. The mean is fairly low, which probably makes sense. About 1 in 610 people commits a robbery.

```
svymean(~robbery, ncvs_survey)
```

	mean	SE
robbery	0.00164	2e-04