

HW4

Introml

Tianyunyang Wang

N14855366 tw1719

1. Suggest possible response variables and predictors for the following classification problems. For each problem, indicate how many classes there are. There is no single correct answer.

- (a) Given an audio sample, to detect the gender of the voice.
- (b) A electronic writing pad records motion of a stylus and it is desired to determine which letter or number was written. Assume a segmentation algorithm is already run which indicates very reliably the beginning and end time of the writing of each character.

(a)

Class: Male or Female

Variables: mean frequency; peak frequency; spectral entropy; standard deviation of frequency.

(b)

Class: 26 uppercase letters, 26 lowercase letters and 10 numbers

Variables: the trace of the motion and pressure.

2. Suppose that a logistic regression model for a binary class label $y = 0, 1$ is given by

$$P(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-z}}, \quad z = \beta_0 + \beta_1 x_1 + \beta_2 x_2,$$

where $\beta = [1, 2, 3]^T$. Describe the following sets:

- (a) The set of \mathbf{x} such that $P(y = 1|\mathbf{x}) > P(y = 0|\mathbf{x})$.
- (b) The set of \mathbf{x} such that $P(y = 1|\mathbf{x}) > 0.8$.
- (c) The set of x_1 such that $P(y = 1|\mathbf{x}) > 0.8$ and $x_2 = 0.5$.

(a)

$$P(y = 1 | x) = \frac{1}{1+e^{-z}}$$

$$P(y = 0 | x) = 1 - \frac{1}{1+e^{-z}}$$

$$P(y = 1 | x) > P(y = 0 | x)$$

$$\frac{1}{1+e^{-z}} > 1 - \frac{1}{1+e^{-z}}$$

$$e^{-z} < 1$$

$$\therefore z = \beta_0 + \beta_1 x_1 + \beta_2 x_2 > 0$$

(b)

$$P(y = 1 \mid x) > 0.8$$

$$\frac{1}{1+e^{-z}} > 0.8$$

$$z > \ln(4)$$

(c)

$$\text{when } z = 1 + 2x_1 + 3x_2 > \ln(4) \text{ and } x_2 = 0.5$$

$$\therefore x_1 > \frac{\ln(4)-2.5}{3}$$

3. A data scientist is hired by a political candidate to predict who will donate money. The data scientist decides to use two predictors for each possible donor:

- x_1 = the income of the person(in thousands of dollars), and
- x_2 = the number of websites with similar political views as the candidate the person follow on Facebook.

To train the model, the scientist tries to solicit donations from a randomly selected subset of people and records who donates or not. She obtains the following data:


Income (thousands \$), x_{i1}	30	50	70	80	100
Num websites, x_{i2}	0	1	1	2	1
Donate (1=yes or 0=no), y_i	0	1	0	1	1

(a)

```

In [2]: import numpy as np
import matplotlib
import matplotlib.pyplot as plt
%matplotlib inline
X1=np.array([30,50,70,80,100])
X2=np.array([0,1,1,2,1])
y=np.array([0,1,0,1,1])
le=len(y)
for i in range(le):
    if y[i]==0:
        plt.scatter(X1[i],X2[i],c='r')
    if y[i]==1:
        plt.scatter(X1[i],X2[i],c='b')

```



(b)

One of the linear classifiers could be w [0,1] and $b=-0.5$

(c)

$$P(y_i = 1 | x_i) = \frac{1}{1+e^{-z_i}}$$

$$P(y_i = 0 | x_i) = 1 - \frac{1}{1+e^{-z_i}}$$

$$P(y_i | x_i) = \frac{1}{1+e^{-u_i}}$$

where $u_i = z_i$ if $y_i = 1$

and $u_i = -z_i$ if $y_i = 0$

Through calculation, we reach the misclassified point when $i=3$.

(d)

Since $\alpha > 0$, \hat{y} will not change.

But the likelihoods will change under the demand of whether $0 < \alpha < 1$ or $\alpha > 1$

$$P(y_i = 1 | x_i) = \frac{1}{1+e^{-\alpha z_i}}$$

$$P(y_i = 0 | x_i) = \frac{1}{1+e^{\alpha_i}}$$

4. Suppose we collect data for a group of students in a machine learning class with variables X_1 = hours studied, X_2 = undergrad GPA, and Y = receive an A. We fit a logistic regression and produce estimated coefficient, $\beta_0 = -6$, $\beta_1 = 0.05$, $\beta_2 = 1$.

- (a) Estimate the probability that a student who studies for 40 h and has an undergrad GPA of 3.5 gets an A in the class.
 (b) How many hours would the student in part (a) need to study to have a 50 % chance of getting an A in the class?

(a)

$$z = \beta_0 + \beta_1 x_1 + \beta_2 x_2$$

$$z = -6 - 0.05x_1 + x_2 = -0.5 < 0$$

$$\therefore P(y = 1 | x) = \frac{1}{1+e^{-z}} = 0.378$$

(b)

$$P(y = 1 | x) = \frac{1}{1+e^{-z}} = 0.5$$

$$z = 0, x_1 = 50$$

The student needs 50 hours.

5. The loss function for logistic regression is the binary cross entropy defined as

$$J(\beta) = \sum_{i=1}^N \ln(1 + e^{z_i}) - y_i z_i$$

where $z_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}$ for two features \mathbf{x}_1 and \mathbf{x}_2 .

- (a) What are the partial derivatives of z_i with respect to β_0 , β_1 , and β_2 .
 (b) Compute the partial derivatives of $J(\beta)$ with respect to β_0 , β_1 , and β_2 . You should use the chain rule of differentiation.
 (c) Can you find the close form expressions for the optimal parameters $\hat{\beta}_0$, $\hat{\beta}_1$, and $\hat{\beta}_2$ by putting the derivatives of $J(\beta)$ to 0? What methods can be used to optimize the loss function $J(\beta)$?

(a)

$$\frac{\alpha z_i}{\alpha \beta_0} = 1$$

$$\frac{\alpha z_i}{\alpha \beta_1} = x_{1i}$$

$$\frac{\alpha z_i}{\alpha \beta_2} = x_{2i}$$

(b)

$$J(\beta) = \sum_{i=1}^N \ln(1 + e^{z_i}) - y_i z_i$$

$$\frac{\alpha J}{\alpha z_i} = \sum_{i=1}^n \frac{e^{z_i}}{1+e^{z_i}} - y_i$$

$$\frac{\alpha J}{\alpha z_i} = \sum_{i=1}^n \frac{e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}}{1 + e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}} - y_i$$

$$\frac{\alpha J}{\alpha \beta_0} = \frac{\alpha J}{\alpha z_i} \frac{\alpha z_i}{\alpha \beta_0} = \sum_{i=1}^n \frac{e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}}{1 + e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}} - y_i$$

$$\frac{\alpha J}{\alpha \beta_1} = \frac{\alpha J}{\alpha z_i} \frac{\alpha z_i}{\alpha \beta_1} = x_{1i} \left(\sum_{i=1}^n \frac{e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}}{1 + e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}} - y_i \right)$$

$$\frac{\alpha J}{\alpha \beta_2} = \frac{\alpha J}{\alpha z_i} \frac{\alpha z_i}{\alpha \beta_2} = x_{2i} \left(\sum_{i=1}^n \frac{e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}}{1 + e^{\beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}}} - y_i \right)$$

(c)

We cannot find the close form expressions for the optimal parameters by putting the derivatives of $J(\beta) = 0$.

Instead, we could use numerical method to optimize the loss function $J(\beta)$