

# Detecting the impact of land cover change on observed rainfall.

Chun X. Liang<sup>1</sup>, Floris F. van Ogtrop<sup>1</sup>, and R. Willem Vervoort<sup>1</sup>

<sup>1</sup>Sydney Institute of Agriculture, The University of Sydney, NSW 2006

Corresponding author:

R. Willem Vervoort<sup>1</sup>

Email address: [willem.vervoort@sydney.edu.au](mailto:willem.vervoort@sydney.edu.au)

## ABSTRACT

Analysis of observational data to identify relationships between rainfall and land cover change are difficult due to multiple environmental factors that cannot be strictly controlled. In this study we present a methodology to investigate the relationship using statistical methods on data from best available sources at two sites in Australia. Gridded data of rainfall and tree cover were used as spatially corresponding local conditions. Large scale effects were represented by climatic indicators, such as SOI and IOD. Regression analysis and step trend tests were used to assess the effect of abrupt land surface intervention. At a Queensland site, significant tree cover change between 2002 - 2005 did not result in strong statistically significant precipitation changes. On the other hand, results from a bushfire affected NSW/VIC region suggests significant changes in the rainfall. This indicates the method works better when a abrupt change in the data can be clearly identified. The results from the step trend test implied a positive relationship between the tree cover and the rainfall at 0.1 significance level in both locations in data up to 2009. However, high rainfall variability and possible regrowth meant that no significant changes were observed in longer time series to 2015.

## INTRODUCTION

Land use and land cover changes can lead to changes in the local climate. Empirical and modelling studies have found that cloud types and rainfall are correlated to large scale vegetation cover changes, such as deforestation in the Amazon and in the Sahel (Chagnon and Bras 2005; Pinto et al. 2009; Wang et al. 2009; Mei and Wang 2010; Kucharski, Zeng, and Kalnay 2013; Pitman and Lorenz 2016) and afforestation in south Israel (Otterman et al. 1990; Ben-Gai et al. 1998). Using airborne measurement in Western Australia, Junkermann et al. (2009) showed a significantly higher level of aerosols over an agricultural area compared to an adjacent natural vegetation. They suggested that a modification of aerosol concentrations due to deforestation could have contributed to a reduction of local rainfall, as more, but smaller rain droplets were observed. Nair et al. (2011) reported from the Bunny Fence Experiment in Western Australia that local land use change altered the synoptic west coast trough dynamics and surface roughness, and this resulted in an observed rainfall decrease. Maximum temperatures were also found to be sensitive to land cover change in eastern Australia (McAlpine et al. 2007).

Overall the number of empirical studies analyzing changes to rainfall due to land cover change from observational data is limited. Most of the studies mentioned previously were either model simulations, or comparisons of modelled data with observations. This is because there are some fundamental experimental difficulties in both space (where does evaporated water reappear as rainfall?) and in time (how much time does it take for land cover change effects to appear or disappear?). In addition, in many areas across the globe, rainfall variability is related to a complex set of interactions, of which land use change might only be a minor component.

Locally, there are two main sources that generate rainfall: moisture from advective atmospheric transport; and local evapotranspiration (Eltahir and Bras 1996; Bosilovich and Chern 2006; Dirmeyer, Brubaker, and DelSole 2009; Gimeno et al. 2010). The local evapotranspiration component is the component considered to be affected by land use change (Eltahir and Bras 1996). According to Trenberth (1999), the contribution of advective moisture partially depends on the availability of external moisture and

47 atmospheric transport. On the longer time scale, such as monthly and annually, large scale atmospheric  
48 dynamics are affected by large scale climate drivers. For example, many studies have reported significant  
49 relationships between rainfall in large parts of Australia and the El Niño-Southern Oscillation (ENSO)  
50 (Verdon et al. 2004; Risbey et al. 2009; Speer, Leslie, and Fierro 2011). In contrast, local ET is determined  
51 by local land surface characteristics, which influence local scale atmospheric dynamics and hence the  
52 amount of rainfall, including contribution from both main sources.

53 Although climate drivers demonstrate some capability to predict Australian rainfall, there is still a  
54 large amount of unexplained variance. Westra and Sharma (2010) pointed out that models based on global  
55 sea surface temperature anomalies can only predict up to 14.7% of annual precipitation variance. Some of  
56 the remaining variance could be due to land surface processes as suggested in studies predicting local  
57 rainfall (e.g. Ma et al. 2011; Zeng et al. 2012; Pitman and Lorenz 2016; Saha, Dirmeyer, and Chase 2016).  
58 However, most are based on modelling experiments and few empirical observational studies have been  
59 reported. However, Pitman et al. (2004) found a good match between observations and simulated rainfall  
60 changes in southwest Western Australia, forced by land cover change. Timbal and Arblaster (2006)  
61 were able to reproduce the rainfall decline in south west Australia by including land cover influence. In  
62 addition, local land use change might not be a primary, but is likely to be a secondary cause of rainfall  
63 change (Nicholls 2006).

64 Therefore, the aim of this study is to use a statistical approach on rainfall data at regional scales  
65 to investigate the cause and effect relationship between land cover change and local rainfall, which is  
66 demonstrated in many modelling studies. More specifically, we hypothesize that a step change on the  
67 land cover on the surface will cause a step change in the rainfall. To demonstrate this we study changes  
68 in observed rainfall over time at a Queensland and NSW/Victoria location where there are possible step  
69 changes in land cover change due to land clearing and bush fires. The methodology uses statistical  
70 approaches to identify changes in rainfall, which are subsequently associated with land cover change  
71 through spatial comparison.


72 In this paper, after this section (the introduction), section 2 covers the case study areas and the observed  
73 land use change. Section 3 describes the data used in the study in more detail. Section 4 details the  
74 statistical methods and the underlying assumptions related to the modelling approach, Section 5 gives the  
75 results, which are further discussed in section 6 and finally section 7 offers the conclusions.

## 76 1 STUDY REGIONS AND TREE COVER CHANGE

77 In Australia, significant tree cover change has mainly occurred in the north east and south east of the  
78 continent, as well as in the southwest of Western Australia. According to the National Dynamic Land  
79 Cover Dataset (DLCD) (Lymburner et al. 2010), most of these areas have experienced decreases in the  
80 Enhanced Vegetation Index (EVI) post 2000 derived from satellite data. As an index for vegetation  
81 greenness, the decreasing values indicate lower biomass over time in the tree cover regions. The possible  
82 EVI reduction might be due to land clearing, bush fires or drought.

83 Two regions were selected where significant tree cover change since 2000 was reported. The first  
84 region is located in south central Queensland partly covering the north of the Murray Darling Basin  
85 (MDB) (site 1 in Figure @ref(fig:selreg)). High rates of land clearing have been reported in this region  
86 during the early 2000s (Department of Natural Resources and Water 2007). The second study region  
87 is located at the border of New South Wales and Victoria, and includes the Snowy Mountain ranges  
88 (site 2 in Figure @ref(fig:selreg)). Severe bush fires occurred in this area in early 2003 (see Figure  
89 @ref(fig:figure2bushfire)). The 2003 bush fires were the largest and the worst in the last 60 years (The  
90 State Government of Victoria 2011). Two thirds of Kosciuszko national park was heavily burned and  
91 regrowth was reported to be slow due to drought and cold conditions (ABC News 2003), and the type of  
92 species in this region. However, in the longer term, after an early high transpiration period a recovery of  
93 pre-fire evapotranspiration would be expected (Kuczera 1987). For the purpose of this study, significant  
94 tree cover loss has happened in both study areas in the last decade, either permanently or temporarily.

95 The two regions have different characteristics. The QLD region is partially grassland and subtropical,  
96 while the NSW/VIC region is mainly within the temperate zone, under the Köppen classification. Accord-  
97 ing to Australian Bureau of Meteorology (BoM), the NSW/VIC region receives 1000 - 2000 mm rainfall  
98 annually, which is more than double the annual rainfall in the QLD region. Evapotranspiration is similar  
99 in both regions. Marine moisture and orographic effects are likely to be the main contributors to rainfall  
100 in the southeast mountain areas of the NSW/VIC region.




**Figure 1.** Selected study regions are highlighted by red rectangles in the main map (the red rectangle in the insert indicates the location of the main map). The types of tree cover in 2008 from the DLCD product is shown at the background. In site 1 (the QLD region), the tree cover is mostly sparse. In site 2 (the NSW/VIC region), many areas have open or close forest where the tree cover is denser.

The land use and land cover characteristics in the two regions are also different. In the Queensland region, the tree cover is sparse over most of the area. The MODIS satellite tree cover data (discussed in more detail in section 3) shows that tree cover in this region is generally below 20% of total ground area. Grazing is the main activity in this region, with over 90% of land used by the grazing industry (ABARES 2010). Our starting assumption is that the main cause of the EVI decline over large part of the region is due to land clearing. Tree cover has been cleared at a massive scale over the last decade, especially during 2002 - 2004. The reports from the Queensland Statewide Land Cover and Trees Study (SLATS) (e.g. Department of Natural Resources and Mines 2005; Department of Science, Information Technology and Innovation 2017) were used to investigate the time and location of the land clearing in the QLD region.


The Kosciuszko national park is within the NSW/VIC region. Here tree cover is denser with open or even closed forest (the tree cover distribution is bimodal at 10 - 20% and 60 - 70%). The dominant species in the alpine area are Snow Gum and large stand species such as Alpine Ash and Mountain Gum in the sub-alpine area. These trees can reach a great height but they take long time to grow. For example, Alpine Ash would need about 20 years to mature. Although land clearing is not a major issue in this region, it is vulnerable to fires and drought. The MODIS burned area product, MCD45A1 (Roy, Lewis, and Justice 2002; Roy et al. 2005; Roy et al. 2008), was used to locate bush fires areas in the NSW/VIC region, with a grid resolution of 500 m. MCD45A1 provides monthly burning information on all pixels, which helps to pinpoint an abrupt event.

Due to the difference in nature of the land cover change in the two regions, the post-change vegetation status is hypothesized to be different as well (see Figure @ref(fig:figure3-tc-simple)).

The overall hypothesis is that the effect of 2003 - 2004 land clearings in the QLD region and the 2003 bush fires in the NSW/VIC region cause a step change in the local rainfall. The actual tree cover change during this time at the pixel level was derived from the 15-year MODIS data (discussed below). The difference of tree cover before and after the land disturbance was tested using a Student's t-test. As the length of the tree cover data is shorter than available rainfall data, earlier land clearing in the QLD region cannot be identified spatially, hence they are excluded from the analysis.



**Figure 2.** Location of bushfires occurring in January 2003, in and around the NSW/VIC study region, as shown by the red pixels. The map shows the large area in the Kosciuszko national park that has been burned. Some locations in the southwest of the Australian Capital Territory (ACT) have also experienced intensive bushfires.



**Figure 3.** The expected evolution of the land surface after trees have been removed in (a) the QLD region and (b) the NSW/VIC region.

### 1.1 Data

Several land surface data sets were used in this study. The main one was the MOD44B product Global Vegetation Continuous Field data set (version 5). This data set provides estimates of percent tree cover (percentage of ground surface covered by trees) at a grid resolution of 250 m (Townshend et al. 2011), which is finer than the earlier mentioned burned product MCD45A1. The data set is available on an annual time interval for the study period of 2000 - 2015. The tree cover data was produced from 16-day Terra MODIS Land Surface Reflectance data and Land Surface Temperature (Townshend et al. 2011). The National Dynamic Land Cover Dataset (DLCD) (Lymburner et al. 2010) from the Australian Collaborative Land Use Mapping Program (ACLUMP) was used to verify the trend of vegetation cover change calculated from the previous data set. This data set, developed by Geoscience Australia and Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES), is the first nationally consistent and thematically comprehensive land cover reference for Australia. The DLCD is based on the 16-day Enhanced Vegetation Index (EVI), again from the MODIS satellite, between April 2000 and April 2015. It also has a grid resolution of 250 m. The data set provides information on the final land cover types (as in 2015) and estimated trend of EVI statistics (annual mean, maximum and minimum).

The SILO Rainfall product data for Australia was used (Jeffrey et al. 2001) (available online at <https://silo.longpaddock.qld.gov.au/>). The data has been projected onto a national 0.05° by 0.05° grid (approximately 5 km by 5 km). This gridded data set was generated from station observations using spline interpolation and kriging (Jeffrey et al. 2001). The data has been compared to other gridded products and observed data and is generally of high quality (Tozer, Kiem, and Verdon-Kidd 2009; Tozer, Kiem, and Verdon-Kidd 2012). The data is available on a daily and monthly basis from 1889 to current. Here a subset of 36 years (1979 - 2015) was used. The study was conducted on monthly data, as a land cover change effect on annual rainfall might be negligible, but can often be significant in particular months or seasons (e.g. Otterman et al. 1990; Gaertner et al. 2001; Semazzi and Song 2001; Oleson et al. 2004; Deo et al. 2009).

Large scale climate drivers are represented by various climatic indices. The Southern Oscillation Index (SOI) is generally regarded as a good predictor of Australian rainfall (Risbey et al. 2009; Chowdhury and Beecham 2010; Westra and Sharma 2010), but its skill is weaker in some parts of Australia. For example the Southern Annular Mode (SAM) is found to be more important than ENSO in south Western Australia (Meneghini, Simmonds, and Smith 2007). The testing of the suitability of each index for the regions of

157 interest is described in a later section. The following climate indices were used as candidate predictors for  
158 local rainfall.

- 159 • Southern Oscillation Index (SOI). The Troup version of the monthly SOI series used in this  
160 study was obtained from BoM (available online at [http://www.bom.gov.au/climate/  
161 current/soihtml.shtml](http://www.bom.gov.au/climate/current/soihtml.shtml)).
- 162
- 163 • Eastern, East Central and Central Tropical Pacific Sea Surface Temperatures (NINO 3, NINO  
164 3.4 and NINO 4). Monthly SST anomalies are available from IRI/LDEO data library and the  
165 extended NINO data set is used (available online at [http://iridl.ldeo.columbia.edu/  
166 SOURCES/.Indices/.nino/.EXTENDED/](http://iridl.ldeo.columbia.edu/SOURCES/.Indices/.nino/.EXTENDED/)).
- 167
- 168 • Pacific Decadal Oscillation (PDO). The Pacific Decadal Oscillation is the leading principal  
169 component of monthly SST anomaly in the North Pacific Ocean.. The monthly PDO series was  
170 provided by JISAO (Joint Institute for the Study of the Atmosphere and Ocean, University of  
171 Washington) (available online at <http://jisao.washington.edu/pdo/PDO.latest>).
- 172
- 173 • Indian Ocean Dipole (IOD). The Indian Ocean dipole is commonly measured by the difference  
174 between SST anomaly in the western (50 - 70°E and 10°S-10°N) and eastern (90 - 110°E and 0  
175 - 10°S) equatorial India Ocean (Saji et al. 1999). Monthly IOD was obtained from JAMSTEC  
176 (the Japan Agency for Marine-Earth Science and Technology) (available online at [http://www.  
177 jamstec.go.jp/frcgc/research/d1/iod/DATA/dmi.monthly.txt](http://www.jamstec.go.jp/frcgc/research/d1/iod/DATA/dmi.monthly.txt)).

## 178 **1.2 Statistical method**


179 As an initial analysis a simple boxplot and t-test is used to analyse whether there is a significant change in  
180 tree cover in time, and before and after the suspected change in the regions.

181 To assess the actual causal relationship between the tree cover and the rainfall a flexible regression  
182 model is applied (discussed in detail below). A step change is not directly obvious in the time series of the  
183 rainfall anomalies (Figure @ref(fig:ts-mean)) for both regions, even though the data is deseasonalised and  
184 detrended. In this study we apply different statistical methods to analyse the effect of tree cover change  
185 on rainfall. Both methods make use of a regression model to remove year-on-year variability in rainfall to  
186 strengthen the tree cover change signal.

187 In the first method, the tree cover change is implemented as a factor variable in the regression model,  
188 and the significance of this variable is tested. In the second method, a rank sum test (step trend test), is  
189 applied to the regression model residuals after effects of other major factors were removed. This assumes  
190 that after removal of all climate, long term linear trend and seasonal variation, the vegetation cover change  
191 is the only factor explaining the non-random pattern in the rainfall residuals.


### 192 **1.2.1 Regression model**

193 As highlighted in the introduction, the Australian climate is influenced by sea surface temperatures in  
194 the tropical Pacific and Indian Oceans, as well as pressure systems in the Southern Ocean (BoM 2012b).  
195 Risbey et al. (2009) compared five large-scale drivers, including ENSO (measured by SOI and the  
196 Tropical Pacific Sea surface temperatures (SSTs)), IOD, SAM, MJO (Madden-Julian oscillation) and  
197 blocking, in relation to Australian rainfall variability. The MJO is a large scale eastward-propagating  
198 wave-like disturbance located around equatorial latitudes (Risbey et al. 2009). They identified SOI as  
199 the most important index among all climate indices tested for broad parts of Australia (including QLD  
200 and NSW/VIC) in almost any season. In this study, four climate indices were selected from the main  
201 climatic indicators (see section Data) and used as the explanatory variables in the model for each study  
202 region. A further complicating factor is the influence of the “millenium drought” over the study period  
203 and in particular the change to wet conditions in 2010 - 2011 (Dijk and Viney 2013). Therefore, the  
204 spatially averaged monthly rainfall in the Murray Darling Basin (MDB, downloaded from the Bureau of  
205 Meteorology) was used to explain the year-on-year variation in the rainfall in the regions. As climate  
206 alone are unlikely able to fully explain the temporal variation in rainfall in the regions, including the  
207 drought (Dijk and Viney 2013; Westra and Sharma 2010), this variable was included. Since both regions  
208 at least partly overlap with the MDB, the average rainfall across was assumed to be a useful explaining  
209 variable.



**Figure 4.** The deseasonalised and detrended rainfall over the 30 years period in (a) the QLD region and (b) the NSW/VIC region. The vertical red lines indicate the year of 2003, in which the studied land cover changes occurred. A change in the time series data is not obvious before and after the land cover changes.






**Figure 5.** Cross-correlation of six climate indices and rainfall in QLD study region. For the PDO analysis, 108-year rainfall data (1900 - 2008) are used. Otherwise, 36-year rainfall data are used. The correlation with NINO 3 is not shown as it is very similar to but weaker than for NINO 3.4.

Correlations between rainfall and each climate index were analysed. Rainfall in each study region was first deseasonalised and detrended using the seasonal decomposition function `ds` in the package `deseasonalise` in R (R Core Team, 2018). Using detrended data gives a better indication of the underlying correlation compared to the correlation between trends in the data (Smith and Timbal 2012). The cross-correlations between the deseasonalised and detrended rainfall and the climatic indices were tested using the Pearson's product moment correlation method, assuming the relationships are linear. Although the optimal technique for exploring the correlation with each index could be different as described in Risbey et al. (2009), the Pearson's method was applied to all indices for consistency. Because the PDO describes the multi-decadal SST with lower frequency (MacDonald and Case 2005; Zanchettin et al. 2008; Kamruzzaman, Beecham, and Metcalfe 2011), instead of 37-year rainfall data, a longer period (115 years, from 1900 to 2015) was used to estimate the correlation with PDO, up to lag 24. For the other indices, the 37-year data was used.

Based on the correlation between the climatic indices and rainfall in the regions (as shown in Figure @ref(fig:cor-rain-qld) and Figure @ref(fig:cor-rain-nsw)), it can be concluded that:

- In QLD, the correlation between rainfall and SOI at zero time lags is the strongest across all indices, outweighing the other ENSO indicators. IOD and PDO have a weak influence in QLD.
- In NSW/VIC, again the SOI has the strongest correlation with rainfall, followed by the IOD. For both, the strongest correlations occur at the zero time lags.





**Figure 6.** Cross-correlation of six climate indices and rainfall in NSW/VIC study region.

- in both cases PDO had the weakest correlations, and this factor was not further considered as a predictor.
- in general, for the better correlated indices, strongest correlations occurred at zero time lags.

The above findings are consistent with previous studies. Although some indices are serially correlated with rainfall up to several months, the lag zero events have the greatest correlation coefficients. Using multiple climatic index series was generally found most useful in rainfall prediction (e.g. Risbey et al. 2009; Kamruzzaman, Beecham, and Metcalfe 2011).

Rainfall in Australia shows strong seasonal patterns (Holper 2011; Australian Bureau of Statistics 2012). For example, the north part of the country is summer rainfall dominant with a dry winter, while most of the southern part has a winter rainfall regime. This character is driven by the movement of subtropical high pressure systems which dominate the Australian climate (BoM 2012a). These characteristic summer and winter rainfall patterns mean that the seasonal component of rainfall has a periodic pattern which should be included in the model.

Long term trends in the regional rainfall in some parts of Australia are significant (Hughes 2003; Gallant, Hennessy, and Risbey 2007; Chowdhury and Beecham 2010). In the northern and eastern parts of the continent, increasing rainfall is reported over the last century (Hughes 2003). The presence of such long term trends may be confused with the outcome of a step change in rainfall. As a result a linear trend term was implemented in the model to remove any long term effect.

We assumed all the factors are additive smooth components in determining rainfall following Kamruzzaman, Beecham, and Metcalfe (2011). In this case, the rainfall model is a generalised additive model (GAM) (Hastie and Tibshirani 1986) with a log link function  $\eta(\cdot)$  and assuming the residuals are gamma distributed (see Figure @ref(fig:hist-rain)). This means all predictors are modelled as smooth functions, in this case using the shrinkage version of the cubic regression splines (Wood 2011). All splines were limited to 3 knots in flexibility (Wood 2011), to reduce the risk of overfitting.

$$g(E(\mathbf{R}_r)) = \beta_0 + s_1(\mathbf{MDB}_{\text{monthlyRain}} + s_2(\mathbf{SOI}) + s_3(\mathbf{IOD}) + s_4(\mathbf{Nino3.4}) + s_5(\mathbf{Nino4}) + s_6(\mathbf{Season}) + \beta_1 \mathbf{Trend} + \varepsilon_r \quad (\#eq : model) \quad (1)$$


The bold letters represent the time series vectors. The region is indicated by  $r$ , while  $\beta_u$  ( $u=0, 1$ ) are the fitted coefficients in the model.  $s_v$  ( $v=1, 2, 3, \dots$ ) are the smooth penalized cubic regression spline functions on the climatic indices and the season.  $\mathbf{MDB}_{\{\text{monthlyRain}\}}$  is the spatially averaged monthly rainfall across the Murray Darling Basin. Apart from dropping PDO as a predictor, all other climate indices were included, allowing the model to select the appropriate predictors.

A possible linear long term trend in the rainfall data is modelled by  $\mathbf{Trend} = 1, 2, 3, \dots, n$ , where  $n$  is the total number of months in the time series.  $\mathbf{Season}$  is the seasonal component. The climatic terms are also modelled with smooth functions. The effect of large scale drivers on Australian rainfall is more likely to be seasonal (Murphy and Timbal 2008; Schepen, Wang, and Robertson 2012), and the smooth spline function is more flexible in reproducing the variability in impacts of the climatic indices.

### 1.3 Tree cover change as factor variable

One of the main difficulties in empirical observation studies related to the effect of land cover change on rainfall is the lack of continuous monitoring of land surface variables. Moreover, no specific variable can possibly be defined that can clearly represent the land surface process. Given the lack of a full picture of the land surface process, a factor variable was used in the regression model to represent the abrupt land surface change (see Equation @ref(eq:model)). The change could be a result of either land clearing or bush fires as long as it is permanent or takes a long time to recover. As indicated we approached the problem in two different models ways.

In the first method, the tree cover change was used as a predictor in the regression model, represented by a factor variable  $\mathbf{LC}$ . The significance of the coefficient of  $\mathbf{LC}$ , denoted as  $\beta'_5$  in Equation @ref(eq:model),



**Figure 7.** Distribution of monthly rainfall in (a) QLD and (b) NSW/VIC. Using a Kolmogorov-Smirnov test with shape = 1 and 2.4 for QLD and NSW/VIC respectively, rainfall in both regions can be modelled as a gamma distribution.

272 can be determined by a ratio test.

$$LC = \begin{cases} \text{Trees} \\ \text{Removed} \end{cases} \quad (\#eq : dummyC) \quad (2)$$

273 Therefore in both regions, land cover is “trees” for the period before land cover change and “removed”  
 274 for the period after the change. Here we simply assumed that vegetation cover change has occurred on  
 275 every pixel. The remaining term  $\varepsilon_r$  is the amount of rainfall that is attributed to other unspecified factors  
 276 and random errors. Hence the regression model becomes

$$g(E(\mathbf{R}_r)) = \beta'_0 + s'_1(\mathbf{MDB}_{\text{monthlyRain}} + s'_2(\mathbf{SOI}) + s'_3(\mathbf{IOD}) + s'_4(\mathbf{Nino3.4}) + s'_5(\mathbf{Nino4}) + s'_6\mathbf{Season} + \beta'_1\mathbf{Trend} + \beta'_2\mathbf{LC} + \varepsilon'_r \quad (\#eq : model2) \quad (3)$$

277 One of the difficulties is to point an exact time to the changes in the vegetation cover in the two regions.  
 278 In the QLD region, no exact time can be assigned to the land clearing. According to the SLATS reports,  
 279 the most substantial clearing occurred between 2003 - 2004 . However, the information on the change in  
 280 type of land cover during the time period is missing. Therefore, four scenarios were initially tested in the  
 281 analysis (see **supplementary material**). In the NSW/VIC region, severe bush fires were reported in early  
 282 January 2003. Hence the “tree” cover state was up to December 2002 then it was changed to “removed”  
 283 state from January 2003. As a starting date, the regression model was run from 1979 for both regions.

#### 284 1.4 Step trend test

285 To support the regression analysis, a step trend test was used to detect changes in rainfall as a result  
 286 of vegetation cover change. This nonparametric statistical test was modified from the Mann-Whitney  
 287 Rank-Sum test by (Hirsch and Gilroy 1985) and can identify a step change in data which is cross correlated.  
 288 The gridded rainfall dataset used in this study has a high spatial correlation between neighbouring pixels  
 289 due to the underlying interpolation method. The advantages of using the Rank-Sum test therefore are:  
 290 (1) it does not depend on assumptions of the data distribution; (2) it is not restricted to datasets with no  
 291 missing data; (3) it is robust and not as easily influenced by outliers and negative numbers (Hirsch and  
 292 Gilroy 1985).

293 The rainfall residuals from the regression model in Equation @ref(eq:model2) were used for the step  
 294 trend test. According to (Hirsch and Gilroy 1985), to detect a step change, using deseasonalised and  
 295 detrended data is important. Furthermore, as rainfall can only be partially attributed to local sources and  
 296 conditions, other effects are introduced by large scale dynamics and changes in other climatic factors.  
 297 The assumption is that the regression model should remove these effects, and additionally deseasonalise  
 298 and detrend the rainfall data. As a result, the local landuse effects are amplified in the variation of the  
 299 remaining residuals. The test, described in the following sub section, subsequently associates trends in  
 300 the rainfall residuals with tree cover changes.

##### 301 1.4.1 Mann-Whitney rank-sum statistic

302 As indicated, the step trend test is a modified version of the Mann-Whitney rank-sum statistic (Hirsch and  
 303 Gilroy 1985). As a nonparametric rank-based test, the Mann-Whitney test does not use the exact values of  
 304 rainfall but depends on the ranks of the data. For each month, rainfall residuals of each year were ranked  
 305 in an ascending order. The ranking of January rainfall in a sample pixel k in QLD is illustrated below:

**Table 1.** Example of ranking rainfall residuals

Year	Rainfall residuals	Rank $R'_{1k}$
1998	-0.3	6
1999	-60.9	2
2000	-16.1	4
2001	-71.7	1

Year	Rainfall residuals	Rank $R'_{1k}$
2002	111.1	7
2005	-7.2	5
2006	-60.5	3

Therefore, the smallest or most negative value has rank 1 and the largest value has the maximum rank.

The before and after period in the data formed two groups of samples. The split point of the two periods was based on the timing of the vegetation cover changes. In the QLD region, changes occurred anytime during 2003 and 2004. In contrast to the previous method, the time period covering the land cover change was excluded, as the nonparametric test allows missing data. Hirsch and Gilroy (1985) also pointed out that the power of the test is higher if the data of the change period is ignored. Hence 2003 and 2004 were excluded from the analysis. As a result, the after-change period was 2005 - 2015 for the Queensland location.

In the case of NSW/VIC, the bushfires broke out in early January 2003. The change was within a relatively short period of the year. Therefore the after change period in this region still started in January 2003. Following Hirsch and Gilroy (1985), the before period was set to five years (1998 - 2002) in both regions. The length of the after period is difficult for the NSW/VIC as the regrowth would at some point have impacted the local effects.

There is a further complication in the data. The year 2009 was the final year of the millenium drought (Dijk and Viney 2013), followed by two very wet years (also visible in Figure @ref(fig:ts-mean) in 2010 and 2011. As highlighted earlier, this means the the length of after period could influence the analysis by the inclusion of more or fewer years. as a result the period up to (and including) 2009 was compared to the full period up to 2015. Further analysis up to 2010 is included in the supplementary material.

The rank of rainfall in month  $j$  year  $i$  in pixel  $k$  is denoted as  $R'_{ijk}$ . The sum of ranks of rainfall in month  $j$  in pixel  $k$  before the known intervention is:

$$W_{jk} = \sum_{i=1}^{n_1} R'_{ijk} \cdot (\#eq : W_j) \quad (4)$$

$n_1$  is the number of years before the land cover change. The expected value of  $W_{jk}$  is

$$\mu_w = n_1(n_1 + n_2 + 1)/2 \quad (5)$$

$n_2$  is the number of years after the change. Hence the expected value of the rank sum before the intervention is the same for all months and all pixels. The sum of ranks for the whole time period is fixed, as  $(n_1 + n_2)(n_1 + n_2 + 1)/2$ . In this study, since there are only two groups (before and after), knowing the rank-sum of one group is the same as knowing the rank-sum of the other group. If the rainfall data is temporally and spatially independent, the variance of  $W_{jk}$  is

$$\sigma_w^2 = n_1 \cdot n_2(n_1 + n_2 + 1)/m \quad (6)$$

where  $m$  is the number of months which is 12 in the case of a full year.

#### 1.4.2 Step trend test

Instead of completing the Mann-Whitney U-test, Hirsch and Gilroy (1985) applied a standard normal Z test to the rank-sum statistics. As highlighted, this modified test accounts for serial and cross correlation in the data. In the case here, the deseasonalised and detrended data shows little autocorrelation in the time series but possesses strong cross correlation between neighbouring pixels, i.e.  $R > 0.99$ .

The sum of  $W_{jk}$  for a block of  $ns$  pixels over the whole year,  $\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}$ , has mean

$$E\left(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}\right) = 12 \cdot ns \cdot \mu_w \quad (7)$$

339 and variance

$$Var(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}) = \sum_{j=1}^{12} \sum_{k=1}^{ns} \sum_{h=1}^{ns} C(W_{jk}, W_{jh}). \quad (8)$$

340  $C(W_{jk}, W_{jh})$  is the covariance of the W statistics between pixel k and pixel h in month j. When  $k = h$ ,  
 341  $C(W_{jk}, W_{jh}) = \sigma_w^2$ . When  $k \neq h$ ,

$$C(W_{jk}, W_{jh}) = \sigma_w^2 r(R_k, R_h) \quad (9)$$

342 where  $r(R_k, R_h)$  is the product moment correlation coefficient of the concurrent ranks in pixel k and  
 343 h. Here  $r$  is calculated on the full time series in each pixel. In the analysis, the test was applied to a  
 344 square block of four pixels each time. As argued by Hirsch and Gilroy (1985),  $ns = 4$  is the most optimal  
 345 solution to balance the cost and the gain in the test power.

346 The statistic of the step trend test is then defined as

$$Z' = \frac{\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk} - 12 \cdot ns \cdot \mu_w}{\sqrt{Var(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk})}}. (\#eq : Z) \quad (10)$$

347 The above statistic is written for a 12 month period. By changing the value 12, it can also be used to  
 348 test seasonal rainfall change or for other customized periods.

349 The null hypothesis ( $H_0$ ) in this study is that there was no change in rainfall due to land surface  
 350 intervention. The results of the step trend test can be interpreted according to the sign of the Z' score (see  
 351 Table @ref(tab:Zscore) (Chapter 23, P887 Hipel and McLeod 1994)). Z' is normally distributed similar to  
 352 the standard normal statistics Z. Hence it can be compared to a standard normal distribution to determine  
 353 the p value.


**Table 2.** The interpretation of Z' score in the step trend test

$Z' > 0$ & rainfall decreases after change
$Z' < 0$ & rainfall increases after change
$Z' = 0$ & rainfall does not change

## 354 2 RESULTS

### 355 2.1 Tree cover change

356 The pixels, where the tree cover change based on the MOD44B data was significant ( $p \leq 0.05$ ) in each  
 357 study region, are shown in Figure @ref(fig:tctrend) for the time series to 2009 (left panels) and the time  
 358 series to 2015 (right panels). In both cases it can be seen that the area of negative cover change was  
 359 greater in the series to 2009 than in the series to 2015. But even the longer series indicates a large change  
 360 in the NSW/VIC region (bottom row). In the NSW/VIC region, much of the tree loss between 2002 and  
 361 2003 was concentrated in the Snowy Mountains close to the border of NSW and VIC. Tree cover loss  
 362 occurred in large parts of the QLD region between 2002 and 2005, but this tree loss was spatially less  
 363 concentrated. Most of the clearings appear in the centre of this region in the series to 2009. The tree cover  
 364 change map (@ref(fig:tctrend)) is consistent with the annual mean EVI trend map (based on DLCD data,  
 365 not shown here). However in the series to 2015 the tree cover loss seems to have dissappeared (top right  
 366 Figure @ref(fig:tctrend)) and suggests an increase in tree cover over the area. The implications of this  
 367 will be discussed in more detail in the discussion.



**Figure 8.** The maps show significant changes in tree cover identified from the MOD44B data between 2003 and 2009 (top left) and 2003 and 2015 (top right) in the NSW/VIC region and the Qld region (bottom left to 2009 and bottom right to 2015). The amount of change was calculated as the difference in tree cover before and after the specified land cover intervention and it is shown as the percentage of the ground area. Green colour indicates an increase in tree cover, while red colour indicates a decrease in tree cover.



## 2.2 Regression Model and significance of Vegetation Cover Changes

Generally the regression model only explains a limited amount of the rainfall variability. The model in Equation @ref(eq:model2) accounts for around 15%<sup>1</sup> of the rainfall variation in both regions. The residual analysis shows that the assumptions of the regression model are generally met (Figure @ref(fig:residuals)). The standardised residual plots, however, show some funnelling for the NSW/VIC regions, suggesting non constant variance. The residual patterns are consistent for all pixels within each region.

The model, however, confirms the importance of the climate drivers and the seasonality in Australian rainfall. Even at the grid level, the seasons and several of the climatic indices were significant ( $p \leq 0.05$ ) everywhere in both regions. The explaining power of the model is mostly due to these variables. The climate drivers (at lag zero) accounted, on average, for 6.7% of the rainfall variability in both the QLD region and the NSW/VIC region (see Figure @ref(fig:rsq) for the distribution of  $R^2$  in these two regions). These figures are within the upper bound of seasonal rainfall predictability by a SST anomaly field reported by (Westra and Sharma 2010).

Statistically significant long term trends were only observed in part of the NSW region (results not shown). However, this result might prove or disprove the existence a long term trend in rainfall. The overall time period is fairly short (Koutsoyiannis 2006) and more pixels in NSW/VIC could indicate a significant step change if the long term trend effect is not removed by the model. As trend free data is an important requirement for the step trend test, the trend term was kept in the regression model to ensure the detection of step change was not due to a possible long term trend.

The land cover variable in the model (Equation @ref(eq:model2)) aims to identify a step change in the rainfall before and after the observed change in land cover. The variable was only significant ( $p \leq 0.05$ ) for the rainfall estimates in some areas in NSW/VIC, as shown in Figure @ref(fig:LCp). However, the number of pixels where the landcover variable was significant was much greater for the series up to 2009, compared to the series to 2015. There was no direct relationship between the areas of bushfires in Figure @ref(fig:bushfire). No significant step change due to the land cover changes was found in rainfall in the QLD region in the series to 2015, but a small area of change was identified in the period to 2009 (left bottom panel).

Figure @ref(fig:LCp) for NSW/VIC suggests that a significant step change in rainfall related to land cover change was found in an area larger than where the bushfires has occurred. This might be showing a large scale effect that reaches beyond the vegetation cover change effect.


More generally, the model suggests that the tree cover has a positive impact on rainfall in both regions. The fitted coefficients for the Land cover change variable were consistently positive for the “tree” part of the series. It implies that rainfall was higher when the surface was covered by trees.

## 2.3 Step Trend Test


The spatial step trend test  $Z'$  scores for the Qld region (top row) and NSWVIC region (bottom row) are shown in Figure @ref(fig:steptest30). The figure also contains the period to 2009 (left panels) and the period to 2015 (right panels). This figure provides two types of information: the sign and the significance level. The sign indicates the direction of the step change, as listed in Table @ref(Zscore). In each region up to 2009, there is a broad area of positive  $Z'$  values which implies a decrease in rainfall. There appears to be little relationship between the locations where changes in tree cover are observed (Figure @ref(fig:tc.trend)) and the patterns in the  $Z'$  score. However, there is not necessarily a direct relationship as movement of air masses could mean that actual changes of rainfall are observed close by, but not necessarily exactly at areas with changes of landcover. There is once again a difference between the left panels and the right panels, indicating that considering a longer time series (to 2015) reduces the pattern and significance of  $Z'$  scores. In the QLD region to 2009, 21% of the pixels obtained a positive  $Z'$  score with  $p < 0.1$ , and for the period to 2015 no pixels with positive  $Z'$  score at  $p < 0.1$  occurred. In the NSW/VIC region to 2009, only 2.6% of pixels in the alpine area having a positive  $Z'$  score with  $p < 0.1$ . In general it is only a small proportion of both study regions.

The rainfall residuals of the two periods (before-change since 1979 and after-change) were also compared (Figure @ref(fig:meandiff)) using a simple t-test. From the boxplots it is difficult to see that there were significantly different ( $p < 0.05$ ) mean values between the “before” and “after” periods in both regions. For the Queensland locations, there was no significant difference if measured to 2009 and slightly


<sup>1</sup> Here the adjusted  $R^2$  was reported. Adjusted  $R^2$  is the coefficient of determination, a measurement of the amount of variability predicted by the model adjusting for the number of explanatory terms




**Figure 9.** The residual analysis of a sample pixel in the QLD region (top) and NSW/VIC region (bottom).




**Figure 10.** The performance of the regression model if rainfall is only modelled by the climate drivers. It shows the percentage of rainfall variability that can be explained by the climate drivers for the Qld and NSW/VIC region



**Figure 11.** The spatial distribution of significance of land cover step change variable in the GAM model predicting changes in rainfall in the both regions. The top row reflects NSW/VIC, while the bottom relates to the Qld study site. The left hand plots are the data up to 2009, while the right hand plots are for the full series. The p value reported is for the land cover variable in the model.



**Figure 12.** Spatial distribution of the step trend test  $Z'$  statistics in the two study sites. Panels on the top are for the NSW/Vic site, while panels of the bottom are for Qld. The left panels are for the series to 2009, while the right panels are for the full data series. Warm colours (yellow, orange and red) are for positive  $Z'$  values which indicate decreasing rainfall trend due to the land surface intervention. Cold colours (light blue to blue) are for negative  $Z'$  values which indicate increasing rainfall trend. The deeper the colour, the more significant the statistic.



**Figure 13.** Boxplots of annual rainfall residuals (estimated based on Equation 2 before and after the land cover intervention during 1979 - 2015 in the study regions. On average, the after period has a significantly lower annual rainfall residual in NSW/VIC, but a significantly higher annual rainfall residual in the Qld study area

less rain ( $p < 0.05$ ) before if measured to 2015 (  $2\text{mm/month}$ ). In contrast, for the NSW/VIC locations there was also slightly more rain ( $p < 0.05$ ) after the change if measured to 2015 (  $1\text{mm/month}$ ), but significantly less rain ( $p < 0.05$ ) after the change (  $9\text{mm/month}$ ) if measured to 2009.

A closer look at the rainfall data indicates that rainfall was consistently very low in 2006 for both regions and consistently high for 2010 @ref(fig:ts-mean). The low rainfall in 2006 was due to a weak El Niño and unrecovered conditions from the previous drought, in contrast, 2010 brought record drought breaking rains across Eastern Australia related to a La Niña (Dijk and Viney 2013). While the regression model has removed most of the effect of ENSO, the effect of these anomolous years could still be visible in the residuals if the response of the rainfall to the ENSO effect is non-linear, and for example the memory of the past drought is persistent. This can also explain the significant results in the previous method. To remove the possible outlier effects from 2006 and 2010, the two periods were compared excluding the 2006 and 2010 rainfall. In addition, the group of pixels showing a negative step change in rainfall ( $p < 0.10$ ) and the group of pixels indicating no change were analysed separately for the period to 2009. An unpaired unequal variance two sample t-test was applied to test whether the after period had lower mean rainfall than the before period, excluding the 2006 and 2010 years for the full period, and excluding 2006 for the period to 2009.

For the Queensland locations using the regression model residuals and excluding 2006 and 2010, there was no difference between the period before and after the landuse change for the full data series. However for the period to 2009, there is slightly more rain ( $p < 0.05$ ,  $2\text{mm/month}$ ) before than after the change. For the NSW/VIC locations for the same analyses, there is slightly more rain ( $p < 0.05$ ) after the change if measured to 2015 (  $2\text{mm/month}$ ), and for the period to 2009 there is still significantly less rain ( $p < 0.05$ ,  $5\text{mm/month}$ ). Focussing on the pixels with a negative step change (positive significant  $Z'$  score) and the monthly rainfall totals, only the period to 2009 is of interest, as the full period had too few significant pixels (@ref(fig:steptest30)). For both regions and excluding 2006, the Qld area had significantly more rain before than after at  $p < 0.05$ . For the NSW/VIC area, there was similarly significantly more rain before than after the change ( $p < 0.05$  and  $100\text{mm/month}$ ). For pixels where the  $Z'$  score was not significantly positive ( $p > 0.1$  and  $Z'$  score negative), the Qld area showed no significant difference in monthly rainfall totals before and after. For the NSW/VIC area there was still significantly less rain after the change ( $p < 0.05$ ) but difference in monthly rainfall totals was smaller than for positive and significant  $Z'$  scores: 24 mm/month less.

The choice of  $ns$  has some impact on the test results (Hirsch and Gilroy 1985). The cases of  $ns = 1$  and  $ns = 9$  were also tested. The results for  $ns = 1$  showed a lower number of significant pixels(at  $p < 0.10$ ) compared to the  $ns = 4$  test, with only 66 pixels in the Queensland area in the period to 2010 and a slightly higher 6 pixels in the period to 2015. No pixels were significant at  $p < 0.1$  in the NSW/VIC area. The results from the  $ns = 9$  test was not different from the  $ns = 4$  test. The power of the test does not change much after  $ns = 4$ , as shown by Hirsch and Gilroy (1985).

As part of the analysis, the “field significance” of the  $Z'$  score test was considered to better interpret the step change at regional scales from multiple local tests (Wilks 2006, Westra, Alexander, and Zwiers (2013)). Here, the bootstrapping resampling method from Westra, Alexander, and Zwiers (2013) was used to evaluate the field significance for the period up to 2009 (as the period to 2015 essentially showed no significant positive  $Z'$  scores). This means the spatial structure of the pixels was maintained, but the order of the years and months was changed by random resampling. For each resampling the test statistics identifies the percentage of the pixels with significant positive step change for the step trend test. The test statistics on 1000 resampled replicates were used to develop the distribution of these percentage values under the local null hypothesis that there was no step change. The results showed that none of the resampled series indicated any significant step trend for both regions, suggesting that the actual values found are for a unique series of years.

### 3 DISCUSSION

```
## Warning: package 'kableExtra' was built under R version 3.5.2
```

Overall the summary table of the results indicates that the effects of land clearing or bushfire on rainfall are not easily detected, even with a range of different statistical tests. In addition several of the tests appear to indicate contradictory results, in particular in the comparison between the two regions and between the two time periods.



**Table 3.** Summary table of all tests on the two regions and for the two time periods

Test	Qld 2009	Qld 2015	NSW/VIC 2009	NSW/VIC 2015
LC variable	small area on west side	no pixels significant	Large area	Smaller area than 2009
t-test regression \& model residuals	not significant	less rain before	less rain after	slightly less rain after
t-test residuals \& excluding 2006 and 2010	slightly more rain after	no significant difference	significantly less rain after	slightly more rain after
Z' score	21% of pixels at p ; 0.1	1 pixel at p ; 0.1	2.6% of pixels at p ; 0.1	no pixels at p ; 0.1
t-test pixels positive Z' score	significantly less rain after	NA	100 mm/month less rain after (p ; 0.05)	NA
t-test other pixels	no difference	NA	24 mm/month less rain after (p ; 0.05)	NA

Generally, empirical studies on LCC-precipitation interaction are conducted within an area with known land surface intervention (e.g. Otterman et al. 1990, Durieux, Machado, and Laurent (2003), Negri et al. (2004), Sato, Kimura, and Hasegawa (2007)). However, these locations are rare and difficult to isolate from real landscape change. Modelling studies are abundant, but are generally not linked to observed data. In this study we therefore tested the effect of land cover change across a broad region, rather than only for locations where changes were known to occur or have occurred. The advantage of the suggested approach is that it does not require a long time series of land cover data as this is usually unavailable. Furthermore, it does not assume a specific relationship between vegetation cover change and rainfall but allows the data to show this relationship, by applying the analysis to a broader area outside the boundary of the vegetation cover change. This approach is expected to provide a way to reduce the risk of a false positive paradox, by comparing results between areas with and without vegetation cover change.

While there is some indication that the observed landuse changes (Figure @ref(fig:tctrend)) cause a decrease in the rainfall, our analysis has not been able to give an unequivocal answer. There are several possible complicating factors in the data that could explain the differences between the two regions and the two time periods.


### 3.1 Rainfall variability

The first and most significant effect on the results remains the variability in the rainfall. The regression model used here is a simple model. We only consider the important effects of a historical trend, seasonality and climate drivers. The purpose of using the regression model is to remove the year to year and month to month variability in rainfall and therefore strengthen the land cover signal in the residuals. However, the model shows that trend, seasonality, ENSO and IOD together explain no more than 30% of the rainfall variability, and only around 7% on average is due to the climate drivers (Figure @ref(fig:rsq)). And while this is consistent with the literature e.g. Westra and Sharma (2010), a large amount of variation is left in the model residuals. Rainfall is generally considered a stochastic process (e.g. Fowler et al. 2005, Cowpertwait, Salinger, and Mullen (2009), Burton et al. (2010)) and clearly some of the variability is either a different response to a combination of climate factors (as interactions were not tested in the model), or a strongly non-stationary response to the climate drivers. The remaining variability in the residuals increases the difficulty to detect a change in rainfall. This is clearly demonstrated in the difference in land cover variable significance and the Z' score between the series to 2009 and the series to 2015. While 2009 was at the end of the drought, the series to 2015 includes the record breaking year 2010 (Figure @ref(fig:ts-mean)).

The severe bushfires in 2003 were also triggered by the extreme drought conditions during the millenium drought (Dijk and Viney 2013). Although the drought on rainfall has been accounted for in part by the model, a further delayed or cumulative impact of drought could be feeding into the local land-atmosphere interaction. As a result, the rainfall feedback to the vegetation cover change could be weak under the dry conditions, and this could have affected the result.

### 3.2 Vegetation dynamics

The second possible effect is the dynamic nature of the vegetation clearing and recovery. Although land clearing has occurred at a high rate and broad scale in Queensland (Department of Science, Information Technology and Innovation 2017), the clearing does not have a clear start and end point. QLD has a long history of land clearing. According to the series of SLATS reports on land cover changes in QLD released by the Queensland government, land clearing continued in and around the study region between 1988 - 2008. Major broad scale and high rate clearings occurred in 1999 - 2000 and 2002 - 2004 (Figure @ref(fig:slat)). And even though there was a decrease in land clearing post 2005, it is difficult to define a clear cut change in this region. The more continuous ongoing land clearing could have reduced the



**Figure 14.** Woody vegetation clearing rate in QLD for the two major bioregions in the study area. The data were obtained from SLATS (2017), and clearly indicate a sharp decrease in the clearing rate after 2005.

significance of a step change.

The two approaches used in this study indicate quite different results on the level of change in the NSW/VIC region. The regression model showed that a large area in the NSW/VIC region has experienced a significant land cover effect ( $p < 0.05$ ) on the rainfall after 2003. The effect in the step trend test (Z' scores) is much more subdued, even though it was able to detect significant changes ( $p < 0.1$ ) within the area in the period to 2009, but almost no effect in the period to 2015. In contrast, the results are almost reversed in the Qld area. While some of this could be due to the difficulty in removing year on year rainfall variability, some of this might also be due to vegetation regrowth in both locations.

The specific vegetation class in the Queensland area is well-known for rapid regrowth and "thickening" in favourable conditions (Gowen and Bray 2016), and this could explain the change in vegetation cover between 2009 and 2015 in the Qld area (Figure @ref(fig:tctrend)). In particular the favourable rainfall years of 2010 and 2011 would have boosted regrowth, increasing evapotranspiration and therefore decreasing the effect of the rainfall change.

The different causes of vegetation cover change in these two regions could lead to different post-change characteristics. The magnitude of EVI decreasing trends in the QLD region are less than in NSW region, as reported in the DLCD data. This is due to the lower tree density in the QLD region than in the NSW/VIC region before land surface interventions. The significant bushfires (Figure @ref(fig:fig2bushfire)) would have drastically reduced the vegetation cover and recovery was very slow in some areas within the NSW/VIC area. The persistent drought in the 2000s (Howden 2012, Dijk and Viney (2013)) delayed the regrowth of trees. On the other hand, replacing tree cover with pasture and crops in the Qld area might have a relatively subtle impact on the EVI.

### 3.3 Misalignment rainfall effects and vegetation clearing

Particularly in the NSW/VIC area, there is no direct overlap between the pixels with significant vegetation change (Figure @ref(fig:tctrend)) and the pixels with significant land cover variables or positive Z'scores (Figure @ref(fig:LCp) and Figure @ref(fig:steptest30)). While this seems possibly surprising at first, a plausible explanation could be the general movement of moisture and climate systems over the landscape,

544 resulting in a possible shift of where the moisture is evaporated and where rainfall occurs. Specifically for  
545 this reason, this study selected rather large spatial boxes to capture the overall response rather than the  
546 pixel by pixel comparison/

### 547 **3.4 Gridded monthly rainfall data**

548 The rainfall data used in this study is a gridded data set. This data set is robust and consistent over a  
549 long time series (from 1900 to current) and has a broad national wide coverage which can provide more  
550 information spatially. However, high cross correlation between pixels, due to the interpolation method  
551 generating this data set, can also introduce spatial noise. In the step trend test the cross correlation was  
552 accounted for. Some other methods are also available which can be used to perform a comparative trial.  
553 For example, (Narisma et al. 2007) applied a spatial Gaussian filter on a similar data set and used wavelet  
554 analysis to detect step change in rainfall. High quality station data is another option to test whether the  
555 observed spatial pattern in the step trend test results was not due to the gridded data itself. Resampling  
556 methods, such as bootstrapping and permutation (Wilks 1997, Kundzewicz and Robson (2004), Westra,  
557 Alexander, and Zwiers (2013)), can also be used to further assess the strength of significance of results  
558 and incorporate spatial and temporal patterns in the analysis. While the gridded data set is most useful  
559 in regions with sparse rain gauge networks, it can actually reduces information where the rain gauge  
560 density is high (Jones, Wang, and Fawcett 2009). In the NSW/VIC area, the coverage of rainfall stations  
561 is more intensive but they are mainly located in the valleys. The interpolated data might not be the best  
562 representation of the local rainfall.

### 563 **3.5 General approach**

564 Parametric tests are generally more powerful than nonparametric test in detecting a trend, when the data is  
565 normally distributed (Onoz and Bayazit 2003, Kundzewicz and Robson (2004)). As a non-parametric test,  
566 the step trend test has the advantages of distribution free and having no restriction on missing data (Hirsch  
567 and Gilroy 1985). This is particularly useful in rainfall analysis since rainfall data is usually skewed. On  
568 the other hand, the disadvantages of non-parametric tests, such as being limited to hypothesis testing and  
569 weaker in power, also hold for the step trend test (Whitley and Ball 2002).

570 Overall, the current study provides a clear approach build on several lines of evidence to provide some  
571 evidence to reject the null hypothesis (no step change in rainfall occurs as a result of tree cover loss).  
572 Limited by the available data, the time frame under study was chosen to include a long lasting drought  
573 period (Holper 2011, Dijk and Viney (2013)). The strong impact of this prolonged drought might have  
574 suppressed the land-atmosphere interaction and confused the cause and effect relation between rainfall  
575 and vegetation cover change. This could be one of the reasons that the LCC effects on the local climate  
576 found in other studies (e.g. G6rgen et al. 2006, McAlpine et al. (2007)) are not found to be significant  
577 here. Possible future work could focus on a non-drought period, or when a longer series of land cover data  
578 is available. While the power of the test can be improved with the longer length of the after-intervention  
579 period (Hirsch and Gilroy 1985), the dynamic nature of vegetation regrowth in this case study prevents  
580 this effect. A better approach might be to build a global study that investigates multiple locations where  
581 drastic landcover changes have taken place, which would also remove some of the climate variability  
582 effects due to the larger sample size.

## 583 **4 CONCLUSIONS**

584 In this study, we present an approach to identify if found some observational data based evidence, although  
585 not strong, that vegetation cover change has changed local rainfall. The semi-parametric method and  
586 non-parametric method did totally agree on detecting a significant step change in rainfall in the hot dry  
587 QLD region where land clearing has occurred. On the other hand, the bushfires in the humid, temperate  
588 mountain range in the NSW/VIC region has experienced reduced rainfall. But the dry spell also plays an  
589 important role in the results.

590 Drought has had a pronounced impact on the land surface condition during the study period, leading  
591 to significant reduction in vegetation and extreme events such as bushfires. The associated lack of rainfall  
592 and high temperatures may mask the step change in the vegetation. Hence, the signal of LCC feedback on  
593 rainfall is probably weaker under such regional dry conditions, as the impact of LCC on rainfall is mainly  
594 through changes in moisture convergence (G6rgen et al. 2006, Pitman and Hesse (2007)).

## 595 **5 ACKNOWLEDGMENTS**

596 CL was supported by an Australian Postgraduate Award for this work.

597 **6 APPENDIX**

598 **6.1 Summary of Data**

## REFERENCES

- ABARES. 2010. "Land Use of Australia, Version 4, 2005/2006."
- ABC News. 2003. "Kosciuszko Slow to Recover from Bushfires."
- Australian Bureau of Statistics. 2012. "Geography and Climate - Australia's Climate."
- Ben-Gai, T., A. Bitan, A. Manes, P. Alpert, and S. Rubin. 1998. "Spatial and Temporal Changes in Rainfall Frequency Distribution Patterns in Israel." *Theoretical and Applied Climatology* 61 (3-4): 177-90. <Go to ISI>://000078025000005.
- BoM. 2012a. "Australia - Climate of Our Continent."
- . 2012b. "Australian Climate Influence."
- Bosilovich, Michael G., and Jiun-Dar Chern. 2006. "Simulation of Water Sources and Precipitation Recycling for the MacKenzie, Mississippi, and Amazon River Basins." *Journal of Hydrometeorology* 7 (3): 312-29. <http://dx.doi.org/10.1175%2FJHM501.1>.
- Burton, A, HJ Fowler, S Blenkinsop, and CG Kilsby. 2010. "Downscaling Transient Climate Change Using a Neyman-Scott Rectangular Pulses Stochastic Rainfall Model." *Journal of Hydrology* 381 (1): 18-32.
- Chagnon, F. J. F., and R. L. Bras. 2005. "Contemporary Climate Change in the Amazon." *Geophysical Research Letters* 32 (13). <Go to ISI>://WOS:000230640100004 <http://www.agu.org/journals/gl/gl0513/2005GL022722>
- Chowdhury, R. K., and S. Beecham. 2010. "Australian Rainfall Trends and Their Relation to the Southern Oscillation Index." *Hydrological Processes* 24 (4). John Wiley & Sons, Ltd.: 504-14. <http://dx.doi.org/10.1002/hyp.7504>.
- Cowpertwait, PSP, Jim Salinger, and B Mullen. 2009. "A Spatial-Temporal Stochastic Rainfall Model for Auckland City: Scenarios for Current and Future Climates." *Journal of Hydrology (NZ)* 48: 95-109.
- Deo, RC, JI Syktus, CA McAlpine, and KK Wong. 2009. "The Simulated Impact of Land Cover Change on Climate Extremes in Eastern Australia." In *Proceedings of the 18th World Imacs Congress and Modsim09 International Congress on Modelling and Simulation*, 2035-41. Modelling; Simulation Society of Australia; New Zealand Inc.; International Association for Mathematics; Computers in Simulation.
- Department of Natural Resources and Mines. 2005. "Land Cover Change in Queensland 2001-2003, Incorporating 2001-2002 Adn 2002-2003 Change Periods: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Natural Resources; Mines.
- Department of Natural Resources and Water. 2007. "Land Cover Change in Queensland 2004-2005: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Natural Resources; Water.
- Department of Science, Information Technology and Innovation. 2017. "Land Cover Change in Queensland 2015-2016: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Science, Information Technology; Innovation.
- Dijk, Beck van, Albert I. J. M., and Neil R. Viney. 2013. "The Millennium Drought in Southeast Australia (2001-2009): Natural and Human Causes and Implications for Water Resources, Ecosystems, Economy, and Society." *Water Resources Research* 49 (2): 1040-57. doi:10.1002/wrcr.20123.
- Dirmeyer, P. A., K. L. Brubaker, and T. DelSole. 2009. "Import and Export of Atmospheric Water Vapor Between Nations." *Journal of Hydrology* 365 (1-2): 11-22. <Go to ISI>://000263575600003.
- Durieux, Laurent, Luiz Augusto Toledo Machado, and Henri Laurent. 2003. "The Impact of Deforestation on Cloud Cover over the Amazon Arc of Deforestation." *Remote Sensing of Environment* 86 (1): 132-40. <http://www.sciencedirect.com/science/article/pii/S0034425703000956>.
- Eltahir, E. A. B., and R. L. Bras. 1996. "Precipitation Recycling." *Reviews of Geophysics* 34 (3): 367-78. <Go to ISI>://A1996VD58700003.
- Fowler, HJ, CG Kilsby, PE O'connell, and A Burton. 2005. "A Weather-Type Conditioned Multi-Site Stochastic Rainfall Model for the Generation of Scenarios of Climatic Variability and Change." *Journal of Hydrology* 308 (1): 50-66.
- Gaertner, M. A., O. B. Christensen, J. A. Prego, J. Polcher, C. Gallardo, and M. Castro. 2001. "The Impact of Deforestation on the Hydrological Cycle in the Western Mediterranean: An Ensemble Study with Two Regional Climate Models." *Climate Dynamics* 17 (11): 857-73.
- Gallant, Ailie J. E., Kevin J. Hennessy, and James Risbey. 2007. "Trends in Rainfall Indices for Six Australian Regions: 1910 - 2005." *Australian Meteorological Magazine* 56: 223-39.
- Gimeno, Luis, Anita Drumond, Raquel Nieto, Ricardo M. Trigo, and Andreas Stohl. 2010. "On the Origin of Continental Precipitation." *Geophysical Research Letters* 37 (13). AGU. <http://dx.doi.org>

org/10.1029/2010GL043712.

Gowen, Rebecca, and Steven G. Bray. 2016. "Bioeconomic Modelling of Woody Regrowth Carbon Offset Options in Productive Grazing Systems." *The Rangeland Journal* 38 (3): 307–17. doi:https://doi.org/10.1071/RJ15084.

Görgen, K., A. H. Lynch, A. G. Marshall, and J. Beringer. 2006. "Impact of Abrupt Land Cover Changes by Savanna Fire on Northern Australian Climate." *Journal of Geophysical Research-Atmospheres* 111 (D19): D19106. <http://dx.doi.org/10.1029/2005JD006860>.

Hastie, Trevor, and Robert Tibshirani. 1986. "Generalized Additive Models." *Statistical Science*, 297–310.

Hipel, K.W., and A.I. McLeod. 1994. *Time Series Modelling of Water Resources and Environmental Systems*. Vol. 45. Elsevier Science Ltd.

Hirsch, Robert M., and Edward J. Gilroy. 1985. "Detectability of Step Trends in the Rate of Atmospheric Deposition of Sulfate." *JAWRA Journal of the American Water Resources Association* 21 (5): 773–84. <http://dx.doi.org/10.1111/j.1752-1688.1985.tb00171.x>.

Holper, Paul N. 2011. *Climate Change, Science Information Paper: Australian Rainfall-Past, Present and Future*. Canberra: CSIRO.

Howden, Saffron. 2012. "It's Official: Australia No Longer in Drought." *Brisbane Times*.

Hughes, Lesley. 2003. "Climate Change and Australia: Trends, Projections and Impacts." *Austral Ecology* 28 (4). Blackwell Science Pty: 423–43. <http://dx.doi.org/10.1046/j.1442-9993.2003.01300.x>.

Jeffrey, Stephen J., John O. Carter, Keith B. Moodie, and Alan R. Beswick. 2001. "Using Spatial Interpolation to Construct a Comprehensive Archive of Australian Climate Data." *Journal Article. Environmental Modelling & Software* 16 (4 SU-): 309–30. <http://www.sciencedirect.com/science/article/B6VHC-42SXF8F-2/2/b5c0a24b85149f40ba5d2419ddc0999c>.

Jones, D.A., W. Wang, and R. Fawcett. 2009. "High-Quality Spatial Climate Data-Sets for Australia." *Australian Meteorological and Oceanographic Journal* 58 (4): 233–48.

Junkermann, W., J. Hacker, T. Lyons, and U. Nair. 2009. "Land Use Change Suppresses Precipitation." *Atmospheric Chemistry and Physics* 9 (17): 6531–9. <Go to ISI>://WOS:000269778500018.

Kamruzzaman, M., S. Beecham, and A. V. Metcalfe. 2011. "Non-Stationarity in Rainfall and Temperature in the Murray Darling Basin." *Hydrological Processes* 25 (10). John Wiley & Sons, Ltd.: 1659–75. <http://dx.doi.org/10.1002/hyp.7928>.

Koutsoyiannis, Demetris. 2006. "Nonstationarity Versus Scaling in Hydrology." *Journal Article. Journal of Hydrology* 324 (1-4): 239–54. <http://www.sciencedirect.com/science/article/B6V6C-4HP6GJ9-1/2/67dcc2e8f8e75b7abb918617f5a79a37>.

Kucharski, Fred, Ning Zeng, and Eugenia Kalnay. 2013. "A Further Assessment of Vegetation Feedback on Decadal Sahel Rainfall Variability." *Climate Dynamics* 40 (5-6): 1453–66. <http://link.springer.com/article/10.1007/s00382-012-1397-x>.

Kuczera, George. 1987. "Prediction of Water Yield Reductions Following a Bushfire in Ash-Mixed Species Eucalypt Forest." *Journal of Hydrology* 94 (3-4): 215–36. doi:https://10.1016/0022-1694(87)90054-0.

Kundzewicz, Z.W., and A.J. Robson. 2004. "Change Detection in Hydrological Records-a Review of the Methodology/Revue Méthodologique de La Détection de Changements Dans Les Chroniques Hydrologiques." *Hydrological Sciences Journal* 49 (1).

Lymburner, Leo, Peter Tan, Norman Mueller, Richard Thackway, Adam Lewis, Medhavy Thankappan, Lucy Randall, Anisul Islam, and Udaya Senarath. 2010. "250 Metre Dynamic Land Cover Dataset." Geoscience Australia, Canberra.

Ma, H. Y., C. R. Mechoso, Y. Xue, H. Xiao, C. M. Wu, J. L. Li, and F. De Sales. 2011. "Impact of Land Surface Processes on the South American Warm Season Climate." *Climate Dynamics* 37 (1-2): 187–203. <Go to ISI>://WOS:000293403000012.

MacDonald, Glen M., and Roslyn A. Case. 2005. "Variations in the Pacific Decadal Oscillation over the Past Millennium." *Geophysical Research Letters* 32 (8): L08703. <http://dx.doi.org/10.1029/2005GL022478>.

McAlpine, C. A., J. Syktus, R. C. Deo, P. J. Lawrence, H. A. McGowan, I. G. Watterson, and S. R. Phinn. 2007. "Modeling the Impact of Historical Land Cover Change on Australia's Regional Climate."



708 *Geophysical Research Letters* 34 (22). <Go to ISI>://000251345100005.

709 Mei, R., and G. L. Wang. 2010. "Rain Follows Logging in the Amazon? Results from CAM3-CLM3." *Climate Dynamics* 34 (7-8): 983–96. <Go to ISI>://WOS:000278088400005.

710 Meneghini, Belinda, Ian Simmonds, and Ian N. Smith. 2007. "Association Between Australian Rainfall and the Southern Annular Mode." *International Journal of Climatology* 27 (1): 109–21. <http://dx.doi.org/10.1002/joc.1370>.

711 Murphy, Bradley F., and Bertrand Timbal. 2008. "A Review of Recent Climate Variability and Climate Change in Southeastern Australia." *International Journal of Climatology* 28 (7): 859–79. <http://dx.doi.org/10.1002/joc.1627>.

712 Nair, Udaysankar S., Y. Wu, J. Kala, T. J. Lyons, Sr. Pielke R. A., and J. M. Hacker. 2011. "The Role of Land Use Change on the Development and Evolution of the West Coast Trough, Convective Clouds, and Precipitation in Southwest Australia." *Journal of Geophysical Research-Atmospheres* 116 (D7). AGU: D07103. <http://dx.doi.org/10.1029/2010JD014950>.

713 Narisma, G.T., J.A. Foley, R. Licker, and N. Ramankutty. 2007. "Abrupt Changes in Rainfall During the Twentieth Century." *Geophysical Research Letters* 34 (6): L06710.

714 Negri, A. J., R. F. Adler, L. M. Xu, and J. Surratt. 2004. "The Impact of Amazonian Deforestation on Dry Season Rainfall." *Journal of Climate* 17 (6): 1306–19. <Go to ISI>://WOS:000220450100013 <http://journals.ametsoc.org/doi/pdf/10.1175/1520-0442%282004%29017%3C1306%3ATIOADO%3E2.0.CO%3B2>.

715 Nicholls, N. 2006. "Detecting and Attributing Australian Climate Change: A Review." *Australian Meteorological Magazine* 55 (3). Bureau of Meteorology.: 199–211.

716 Oleson, K. W., G. B. Bonan, S. Levis, and M. Vertenstein. 2004. "Effects of Land Use Change on North American Climate: Impact of Surface Datasets and Model Biogeophysics." *Climate Dynamics* 23 (2): 117–32.

717 Onoz, B., and M. Bayazit. 2003. "The Power of Statistical Tests for Trend Detection." *Turkish Journal of Engineering and Environmental Sciences* 27 (4): 247–51.

718 Otterman, J., A. Manes, S. Rubin, P. Alpert, and D. Starr. 1990. "An Increase of Early Rains in Southern Israel Following Land-Use Change?" *Boundary-Layer Meteorology* 53 (4): 333–51.

719 Pinto, E., Y. Shin, S. A. Cowling, and C. D. Jones. 2009. "Past, Present and Future Vegetation-Cloud Feedbacks in the Amazon Basin." *Climate Dynamics* 32 (6): 741–51. <Go to ISI>://WOS:000264118400001.

720 Pitman, A. J., and P. P. Hesse. 2007. "The Significance of Large-Scale Land Cover Change on the Australian Palaeomonsoon." *Quaternary Science Reviews* 26 (1-2): 189–200. <http://www.sciencedirect.com/science/article/B6VBC-4MC71SN-1/2/0e1d4dff6e15a1de79cefbfa612f9b37>.

721 Pitman, A. J., and R. Lorenz. 2016. "Scale Dependence of the Simulated Impact of Amazonian Deforestation on Regional Climate." *Environmental Research Letters* 11 (9): 094025. doi:10.1088/1748-9326/11/9/094025.

722 Pitman, A. J., G. T. Narisma, R. A. Pielke, and N. J. Holbrook. 2004. "Impact of Land Cover Change on the Climate of Southwest Western Australia." *Journal of Geophysical Research-Atmospheres* 109 (D18). <Go to ISI>://WOS:000224126500001.

723 R Core Team, 2018. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>.

724 Risbey, James S., Michael J. Pook, Peter C. McIntosh, Matthew C. Wheeler, and Harry H. Hendon. 2009. "On the Remote Drivers of Rainfall Variability in Australia." *Monthly Weather Review* 137 (10): 3233–53. <http://dx.doi.org/10.1175/2009MWR2861.1>.

725 Roy, David P, Luigi Boschetti, Christopher O Justice, and J Ju. 2008. "The Collection 5 Modis Burned Area Product—Global Evaluation by Comparison with the Modis Active Fire Product." *Remote Sensing of Environment* 112 (9). Elsevier: 3690–3707.

726 Roy, DP, Y Jin, PE Lewis, and CO Justice. 2005. "Prototyping a Global Algorithm for Systematic Fire-Affected Area Mapping Using Modis Time Series Data." *Remote Sensing of Environment* 97 (2). Elsevier: 137–62.

727 Roy, DP, PE Lewis, and CO Justice. 2002. "Burned Area Mapping Using Multi-Temporal Moderate Spatial Resolution Data—A Bi-Directional Reflectance Model-Based Expectation Approach." *Remote Sensing of Environment* 83 (1). Elsevier: 263–86.

728 Saha, Subodh K., Paul A. Dirmeyer, and Thomas N. Chase. 2016. "Investigating the Impact of

- Land-Use Land-Cover Change on Indian Summer Monsoon Daily Rainfall and Temperature During 1951-2005 Using a Regional Climate Model." *Hydrology and Earth System Sciences* 20 (5): 1765. <http://search.proquest.com/openview/c4242c9ecce075e3545390b1d00f2751/1?pq-origsite=gscholar&cbl=105724>.
- Saji, NH, B.N. Goswami, PN Vinayachandran, and T. Yamagata. 1999. "A Dipole Mode in the Tropical Indian Ocean." *Nature* 401 (6751): 360–63.
- Sato, T., F. Kimura, and A. S. Hasegawa. 2007. "Vegetation and Topographic Control of Cloud Activity over Arid/Semiarid Asia." *Journal of Geophysical Research-Atmospheres* 112 (D24). <Go to ISI>://WOS:000252013700002 <http://www.agu.org/journals/jd/jd0724/2006JD008129/2006JD008129.pdf>.
- Schepen, Andrew, Q. J. Wang, and David Robertson. 2012. "Evidence for Using Lagged Climate Indices to Forecast Australian Seasonal Rainfall." *Journal of Climate* 25 (4). American Meteorological Society: 1230–46.
- Semazzi, F. H. M., and Y. Song. 2001. "A GCM Study of Climate Change Induced by Deforestation in Africa." *Climate Research* 17 (2): 169–82.
- Smith, Ian N., and Bertrand Timbal. 2012. "Links Between Tropical Indices and Southern Australian Rainfall." *International Journal of Climatology* 32 (1): 33–40.
- Speer, Milton, Lance Leslie, and Alexandre Fierro. 2011. "Australian East Coast Rainfall Decline Related to Large Scale Climate Drivers." *Climate Dynamics* 36 (7): 1419–29. <http://dx.doi.org/10.1007/s00382-009-0726-1>.
- The State Government of Victoria. 2011. "Bushfire History."
- Timbal, B., and J.M. Arblaster. 2006. "Land Cover Change as an Additional Forcing to Explain the Rainfall Decline in the South West of Australia." *Geophysical Research Letters* 33 (7): L07717.
- Townshend, J.R.G., M. Carroll, C. Dimiceli, R. Sohlberg, M. Hansen, and R. DeFries. 2011. "Vegetation Continuous Fields MOD44B, 2001 Percent Tree Cover, Collection 5." University of Maryland, College Park, Maryland, 2001.
- Tozer, C. R., A. S. Kiem, and D. C. Verdon-Kidd. 2009. "On the Uncertainties Associated with Using Gridded Rainfall Data as a Proxy for Observed." Conference Paper. In *18th World Imacs Congress and Modsim09 International Congress on Modelling and Simulation.*, edited by R.S. Anderssen, R.D. Braddock, and L.T.H. Newham, 3886–92. <http://www.mssanz.org.au/modsim09/I13/beesley.pdf>.
- . 2012. "On the Uncertainties Associated with Using Gridded Rainfall Data as a Proxy for Observed." Journal Article. *Hydrol. Earth Syst. Sci.* 16 (5): 1481–99. doi:10.5194/hess-16-1481-2012.
- Trenberth, K. E. 1999. "Atmospheric Moisture Recycling: Role of Advection and Local Evaporation." *Journal of Climate* 12 (5): 1368–81. <Go to ISI>://000080145700002.
- Verdon, Danielle C., Adam M. Wyatt, Anthony S. Kiem, and Stewart W. Franks. 2004. "Multidecadal Variability of Rainfall and Streamflow: Eastern Australia." *Water Resources Research* 40 (10): W10201. <http://dx.doi.org/10.1029/2004WR003234>.
- Wang, Jingfeng, Frederic Chagnon, Earle Williams, Alan Betts, Nilton Renno, Luiz Machado, Gautam Bisht, Ryan Knox, and Rafael Bras. 2009. "Impact of Deforestation in the Amazon Basin on Cloud Climatology." *Proceedings of the National Academy of Sciences of the United States of America* 106 (10): 3670. <http://ezproxy.library.usyd.edu.au/login?url=http://search.proquest.com/docview/201417978?accountid=14757>.
- Westra, Seth, Lisa V. Alexander, and Francis W. Zwiers. 2013. "Global Increasing Trends in Annual Maximum Daily Precipitation." *Journal of Climate* 26 (11). American Meteorological Society: 3904–18. <http://dx.doi.org/10.1175/JCLI-D-12-00502.1>.
- Westra, Seth, and Ashish Sharma. 2010. "An Upper Limit to Seasonal Rainfall Predictability?" *Journal of Climate* 23 (12). American Meteorological Society: 3332–51. <http://ezproxy.library.usyd.edu.au/login?url=http://search.proquest.com/docview/224582143?accountid=14757>.
- Whitley, Elise, and Jonathan Ball. 2002. "Statistics Review 6: Nonparametric Methods." *Critical Care, London* 6 (6): 509–13.
- Wilks, D. S. 2006. "On 'Field Significance' and the False Discovery Rate." *Journal of Applied Meteorology and Climatology* 45 (9). American Meteorological Society: 1181–9. <http://ezproxy.library.usyd.edu.au/login?url=http://search.proquest.com/docview/224582143?accountid=14757>.

817 8&rft\_id=info:sid/ProQ%3Amilitary&rft\_val\_fmt=info:ofi/fmt:kev:mtx:journal&rft.genre=article&rft.jtitle=Journal+of+  
818 09-01&rft.volume=45&rft.issue=9&rft.spape=1181&rft.isbn=&rft.btitle=&rft.title=Journal+of+Applied+Meteorology+  
819 Wilks, DS. 1997. "Resampling Hypothesis Tests for Autocorrelated Fields." *Journal of Climate* 10  
820 (1): 65–82.  
821 Wood, S. N. 2011. "Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation  
822 of Semiparametric Generalized Linear Models." *Journal of the Royal Statistical Society (B)* 73 (1): 3–36.  
823 Zanchettin, Davide, Stewart W. Franks, Pietro Traverso, and Mario Tomasino. 2008. "On Enso  
824 Impacts on European Wintertime Rainfalls and Their Modulation by the Nao and the Pacific Multi-  
825 Decadal Variability Described Through the Pdo Index." *International Journal of Climatology* 28 (8). John  
826 Wiley & Sons, Ltd.: 995–1006. <http://dx.doi.org/10.1002/joc.1601>.  
827 Zeng, X. M., Z. H. Wu, S. Song, S. Y. Xiong, Y. Q. Zheng, Z. G. Zhou, and H. Q. Liu. 2012. "Effects  
828 of Land Surface Schemes on the Simulation of a Heavy Rainfall Event by WRF." *Chinese Journal of*  
829 *Geophysics-Chinese Edition* 55 (1): 16–28. <Go to ISI>://WOS:000300129300002.