

1 **Highlights**

- 2 • There are limited studies that analyse observational data of vegetation
3 feedback on rainfall
- 4 • Two sites with major step trend landuse change were analysed for
5 changes in rainfall
- 6 • Statistical change in rainfall detected after bushfire, but not after land
7 clearing

Analysis of land surface effects on local rainfall using observational data

C. Liang, R.W. Vervoort*, F.F. Van Ogtrop*

*Sydney Institute of Agriculture, School of Life and Environmental Sciences, The
University of Sydney*

Abstract

Analysis of observational data to identify relationships between rainfall and land cover change are scarce due to multiple environmental factors that cannot be controlled. In this study we investigated the relationship using statistical methods on data from best available sources at two sites in Australia. Gridded data of rainfall and tree cover were used as spatially corresponding local conditions. Large scale effects were represented by climatic indicators, such as SOI and IOD. Regression analysis and step trend tests were used to assess the effect of abrupt land surface intervention. At a Queensland site, dramatic tree cover change between 2002 - 2005 did not result in strong statistically significant precipitation changes. On the other hand, results from step trend test on 1998 - 2008 rainfall data in a bushfire affected NSW/VIC region has a better match with the tree cover change map. This indicates the method works better when a abrupt change in the data can be clearly identified. The results from the step trend test implied a positive relationship between the tree cover and the rainfall at 0.1 significance level.

Keywords: Observed data analysis, Rainfall change, Landcover feedback

*
Email address: willem.vervoort@sydney.edu.au (R.W. Vervoort)
Preprint submitted to *Journal of Hydrology* November 20, 2017

1. Introduction

Land use and land cover changes can lead to changes in the local climate. Some observational and many modelling studies have found cloud types and rainfall are correlated to large scale vegetation cover changes, such as deforestation in the Amazon and in the Sahel (Chagnon and Bras, 2005; Pinto et al., 2009; Wang et al., 2009; Mei and Wang, 2010; Kucharski et al., 2013; Pitman and Lorenz, 2016) and afforestation in south Israel (Otterman et al., 1990; Ben-Gai et al., 1998). In terms of observational studies, using airborne measurement in Western Australia, Junkermann et al. (2009) showed a significantly higher level of aerosols over an agricultural area compared to an adjacent natural vegetation. They suggested that a modification of aerosol concentrations due to deforestation could have contributed to a reduction of local rainfall, as more but smaller rain droplets were observed. Nair et al. (2011) reported from the Bunny Fence Experiment in Western Australia that local land use change altered the synoptic west coast trough dynamics and surface roughness, and this resulted in the observed rainfall decrease. Climate sensitivity to land cover change is also found in eastern Australia (McAlpine et al., 2007).

Rainfall over land is generally influenced by multiple factors as has been shown from multiple simulation studies. Locally, there are two main sources: moisture from advective atmospheric transport; and local evapotranspiration (Eltahir and Bras, 1996; Bosilovich and Chern, 2006; Dirmeyer et al., 2009; Gimeno et al., 2010). According to Trenberth (1999), the contribution of advective moisture partially depends on the availability of external moisture and atmospheric transport. On the longer time scale, such as monthly and

annually, large scale atmospheric dynamics are affected by large scale climate drivers. Many studies have reported significant relationships between rainfall in large parts of Australia and the El Niño-Southern Oscillation (ENSO) (Verdon et al., 2004; Risbey et al., 2009; Speer et al., 2011). ENSO can be used to represent longer term cycles in rainfall data such as drought. On the other hand, local ET is determined by local land surface characteristics. Local land surface characteristics further influence local scale atmospheric dynamics and hence the amount of rainfall, including contribution from both sources. Therefore land surface plays an important role in local rainfall.

Conventionally, large scale climate drivers are used as predictors for seasonal rainfall forecasts. According to the Australian Bureau of Meteorology (BoM), probabilistic seasonal outlooks are developed based on Australian rainfall and temperature as well as sea surface temperature records from the tropical Pacific and Indian Oceans (BoM, 2012c).

Different parts of the Australian continent can be more or less influenced by different climatic drivers (BoM, 2008). Using both SOI and PDO in a prediction model, Kamruzzaman et al. (2011) reported that PDO was rarely significant for rainfall stations in the MDB and on the southeast coast of Australia, while SOI was at least significant at the 1% level. In addition to SOI, Speer et al. (2011) found that the observed rainfall decrease in the southeast of NSW was linked to an increasingly positive SAM in 1976 - 1998. In southwest WA, the influence of any oceanic indices is small (Smith and Timbal, 2012).

Although climate drivers demonstrate some capability to predict Australian rainfall, there is still a large amount of unexplained variance. Westra

65 and Sharma (2010) pointed out that models based on global sea surface tem-
66 perature anomalies can only predict up to 14.7% of precipitation variance.
67 Recent studies suggest that land surface processes are important for pre-
68 dicting local rainfall (e.g. Ma et al., 2011; Zeng et al., 2012; Pitman and
69 Lorenz, 2016; Saha et al., 2016). However, they are mostly based on mod-
70 elling experiments and little evidence was reported from observations. Pit-
71 man et al. (2004) found a good match between observations and simulated
72 rainfall changes in southwest Western Australia ,forced by land cover change.
73 Timbal and Arblaster (2006) were able to reproduce the rainfall decline in
74 south west Australia by including land cover influence. Local land use change
75 might not be a primary, but is likely to be a secondary cause of rainfall change
76 (Nicholls, 2006). Therefore, land surface modification has, at least partially,
77 contributed to local rainfall variability.

78 Overall the number of observational data studies related to changes to
79 rainfall due to land cover change is limited. This is because there are some
80 fundamental experimental difficulties in both space (where does evaporated
81 water reappear as rainfall?) and in time (how much time does it take for
82 land cover change effects to appear or disappear?). Therefore, the aim of this
83 study is to use observational data at regional scales to investigate the cause
84 and effect relationship between land cover change and local rainfall using
85 empirical evidence. More specifically, we hypothesize that a step change on
86 the land cover on the surface will cause a step change in the rainfall and that
87 this can be identified statistically. To demonstrate the approach and this
88 effect we studied step changes in rainfall at two locations in Queensland and
89 NSW/Victoria where there are possible step changes in land cover change due

90 to land clearing and bushfires based on tree cover data. More specifically,
91 step changes in rainfall were identified statistically, which were subsequently
92 associated with land cover change through spatial comparison.

93 In this paper, after this section (the introduction), section 2 covers the
94 case study areas and the observed land use changed. Section 3 describes
95 the data used in the study in more detail. Section 4 details the statistical
96 methods and the underlying assumptions related to the modelling approach,
97 Section 5 gives the results, which are further discussed in section 6 and finally
98 section 7 offers the conclusions.

99 **2. Study regions and tree cover change**

100 In Australia, significant tree cover change has mainly occurred in the
101 north east of the continent and on the southeast coast, as well as in the
102 southwest of Western Australia. According to the National Dynamic Land
103 Cover Dataset (DLCD) (Lymburner et al., 2010), most of these areas have
104 experienced decreasing EVI between 2000 - 2008. Being a vegetation green-
105 ness index, the decreasing EVI values indicate lower biomass over time in
106 the tree cover regions. The EVI reduction is possibly due to land clearing,
107 bushfires or drought.

108 Two regions were selected where significant tree cover change was present.
109 One region is located in south central Queensland to the north of the Murray
110 Darling Basin (MDB) (site 1 in Figure 1). High rates of land clearing have
111 been reported in this region during the early 2000s (Department of Natural
112 Resources and Water, 2007). The second study region is located at the border
113 of New South Wales and Victoria, including the Snowy Mountains range (site




Figure 1: Selected study regions are highlighted by red rectangles in the main map (the red rectangle in the insert indicates the location of the main map). The types of tree cover in 2008 from the DLCD product is shown at the background. In site 1 (the QLD region), the tree cover is mostly sparse. In site 2 (the NSW/VIC region), many areas have open or close forest in which tree cover is denser.

2 in Figure 1). Severe bushfires occurred in this area and the surroundings in
early 2003 (see Figure 2). The 2003 bushfires were the largest and the worst
in this area for the last 60 years (The State Government of Victoria, 2011).
Two thirds of Kosciuszko national park was heavily burned and regrowth was
reported to be slow due to drought and cold conditions (ABC News, 2003)
and the type of species in this region. In both study regions, significant tree

120 cover loss has happened in the last decade, either permanently or temporarily.




Figure 2: Location of bushfires occurring in January 2003, in and around the NSW/VIC study region, as shown by the red pixels. The map shows large area in the Kosciuszko national park has been burned. Some locations in the southwest of ACT have also experienced intensive bushfires.

121 The two regions have different climate characteristics. The QLD region
122 is partially grassland and partially subtropical, while the NSW/VIC region
123 is mainly within the temperate zone, under the Köppen classification. Ac-
124 cording to Australian Bureau of Meteorology (BoM), the NSW/VIC region
125 receives 1000 - 2000 mm rainfall annually, which is more than double of
126 the rainfall in the QLD region. Evapotranspiration is similar in both regions.

127 Marine moisture and orographic effects are likely to be the main contributors
128 to rainfall in the southeast mountain areas of the NSW/VIC region.

129 The land use and land cover characteristics in the two regions are also
130 different. In the Queensland region, the tree cover is sparse over most of the
131 area. The MODIS satellite tree cover data (discussed in more detail in section
132 3) shows that tree cover in this region is generally below 20% of total ground
133 area. Grazing is the main activity in this region, with over 90% of land used
134 by the grazing industry (ABARES, 2010). Our starting assumption is that
135 the main cause of the EVI decline over large part of the region is due to land
136 clearing. Tree cover has been cleared at a massive scale over the last decade,
137 especially during 2002 - 2004.

138 The Kosciuszko national park is within the NSW/VIC region. Here tree
139 cover is denser with open or closed forest (the tree cover distribution is bi-
140 modal at 10 - 20% and 60 - 70%). The dominant species in the alpine area
141 are Snow Gum and large stand species such as Alpine Ash and Mountain
142 Gum in the sub-alpine area. These trees can reach a great height but they
143 take long time to grow. For example, Alpine Ash would need about 20 years
144 to mature. Although land clearing is not a major issue in this region, it is
145 vulnerable to fires and drought.

146 Therefore two types of land cover changes were studied. The reports
147 from the Queensland Statewide Land Cover and Trees Study (SLATS) (e.g.
148 Department of Natural Resources and Mines, 2005, 2006) were used to in-
149 vestigate the time and location of land clearing in the QLD region. The
150 MODIS burned area product, MCD45A1 (Roy et al., 2002, 2005, 2008), was
151 used to locate bushfires areas in the NSW/VIC region, with a grid resolution

152 of 500 m. MCD45A1 provides monthly burning information on all pixels,
 153 which helps to pinpoint an abrupt event. Due to the nature of the different
 154 land cover change, the post-change vegetation status in the two regions is
 155 expected to be different (see Figure 3).




Figure 3: The expected evolution of the land surface after trees have been removed in (a) the QLD region and (b) the NSW/VIC region.

156 Specifically, this study focused on the effect of 2003 - 2004 land clearings
 157 in the QLD region and the effect of 2003 bushfires in the NSW/VIC region.
 158 These events are expected to cause a step change in the local rainfall. The
 159 actual tree cover change at the pixel level during this time was derived from
 160 the 11-year MODIS data (discussed below). The difference of tree cover
 161 before and after the land disturbance was tested using a Student's t-test. As
 162 the length of the tree cover data is shorter than the length of the rainfall
 163 data, earlier land clearings in the QLD region cannot be identified spatially,
 164 hence they are excluded from the analysis.

165 3. Data

166 Two land surface data sets were used in this study. The main one was the
167 MOD44B product Global Vegetation Continuous Field dataset (version 5).
168 This dataset provides estimates of percent tree cover (percentage of ground
169 surface covered by trees) at a grid resolution of 250 m (Townshend et al.,
170 2011), which is finer than the earlier mentioned burned product MCD45A1.
171 The dataset is available on an annual time interval for the period of 2000
172 - 2010. The National Dynamic Land Cover Dataset (DLCD) (Lymburner
173 et al., 2010) from the Australian Collaborative Land Use Mapping Program
174 (ACLUMP) was used to verify the trend of vegetation cover change calculated
175 from the previous dataset. This dataset, developed by Geoscience Australia
176 and Australian Bureau of Agricultural and Resource Economics and Sciences
177 (ABARES), is the first nationally consistent and thematically comprehensive
178 land cover reference for Australia. The DLCD is based on the 16-day En-
179 hanced Vegetation Index (EVI), again from the MODIS satellite, between
180 April 2000 and April 2008. It also has a grid resolution of 250 m. The
181 dataset provides information on the final land cover types (as in 2008) and
182 estimated trend of EVI statistics (annual mean, maximum and minimum).

183 Rainfall data for Australia (Jones et al., 2009) was obtained from BoM.
184 The data has been projected onto a national $0.05^\circ \times 0.05^\circ$ grid (approximately
185 $5 \text{ km} \times 5 \text{ km}$). This gridded dataset was generated from station observa-
186 tions using an optimised Barnes successive correction technique. The Barnes
187 technique combines a weighted averaging process and defined topographical
188 information to estimate rainfall values between spatial points (BoM, 2009).
189 The resulting dataset provides additional information for data-sparse areas

like central Australia but reduces information in the data-rich areas, such as southeast Australia where station density is up to 20 per 100 km². The data is available on a monthly basis from 1900 to current. Here a subset of 30 years (1979 - 2008) was used. The study was conducted on monthly data, as a land cover change effect on annual rainfall might be negligible but can often found to be significant in particular months or seasons (e.g. Otterman et al., 1990; Gaertner et al., 2001; Semazzi and Song, 2001; Oleson et al., 2004; Deo et al., 2009).

Large scale climate drivers are represented by various climatic indices. The Southern Oscillation Index (SOI) is generally regarded as a good predictor for Australian rainfall (Risbey et al., 2009; Chowdhury and Beecham, 2010; Westra and Sharma, 2010), but its skill is weaker in some parts of Australia. For example the Southern Annular Mode (SAM) is found to be more important than ENSO in south Western Australia (Meneghini et al., 2007). The suitability of each index for the study regions is analysed in detail in section 4.1. The following climate indices were tested as predictors for local rainfall.

- Southern Oscillation Index (SOI). The Troup version of the monthly SOI series used in this study was obtained from BoM (available online at <http://www.bom.gov.au/climate/current/soihtm1.shtml>).
- Eastern, East Central and Central Tropical Pacific Sea Surface Temperatures (NINO 3, NINO 3.4 and NINO 4). Monthly SST anomalies are available from IRI/LDEO data library and the extended NINO dataset is used (available online at <http://iridl.ldeo.columbia.edu/SOURCES/.Indices/.nino/.EXTENDED/>).

- 215 • Pacific Decadal Oscillation (PDO). The Pacific Decadal Oscillation
216 is the leading principal component of monthly SST anomaly in the
217 North Pacific Ocean.. The monthly PDO series was provided by JISAO
218 (Joint Institute for the Study of the Atmosphere and Ocean, University
219 of Washington) (available online at [http://jisao.washington.edu/
220 pdo/PDO.latest](http://jisao.washington.edu/pdo/PDO.latest)).
- 221 • ~~The interaction of PDO and SOI (PDO×SOI) (Kamruzzaman et al.,~~
222 ~~2011).~~
- 223 • Indian Ocean Dipole (IOD). The Indian Ocean dipole is commonly mea-
224 sured by the difference between SST anomaly in the western (50 - 70°E
225 and 10°S-10°N) and eastern (90 - 110°E and 0 - 10°S) equatorial India
226 Ocean (Saji et al., 1999). Monthly IOD was obtained from JAMSTEC
227 (the Japan Agency for Marine-Earth Science and Technology) (avail-
228 able online at [229 DATA/dmi.monthly.txt](http://www.jamstec.go.jp/frcgc/research/d1/iod/)).

230 4. Statistical method

231 As shown in Figure 4, a step change is not obvious in the ~~time series~~
232 data, even ~~though the~~ data is deseasonalised and detrended. Hence, the step
233 changes in the rainfall were analysed using two different statistical methods
234 to provide a comparison. Both methods make use of a regression model to
235 remove variability in rainfall hypothesised to be caused by factors other than
236 vegetation cover change. In the first method, the tree cover change was im-
237 plemented as a factor variable in the regression model. In the second method,

238 a rank sum test (step trend test), was applied to the model residuals after
 239 effects of other major factors have been removed, assuming that vegetation
 240 cover change is the only factor explaining the non-random pattern in the
 241 rainfall residuals.






Figure 4: The deseasonalised and detrended rainfall over the 30 years period in (a) the QLD region and (b) the NSW/VIC region. The vertical red lines indicate the year of 2003, in which the studied land cover changes occurred. A change in the time series data is not obvious before and after the land cover changes.

242 4.1. Regression model

243 Rainfall is generally affected by several factors. Many studies suggest the
 244 large-scale climate drivers, which are related to the global or regional atmo-
 245 spheric air mass movement, are the dominant control on the overall rainfall
 246 (Maynard and Polcher, 2003; DeAngelis et al., 2010; Holper, 2011; Smith
 247 and Timbal, 2012). Long term trends are shown in Australian rainfall, and
 248 this has been linked to global climate change (Cai et al., 2014; Delworth and
 249 Zeng, 2014; Holper, 2011; Karoly, 2014). The difference in available radia-
 250 tion and hence ET in different seasons increase the variability in rainfall. The

251 land surface effect on rainfall can be confused by changes in these factors.
252 Hence a regression model was used to estimate the amount of variability in
253 rainfall that is due to these important factors, and isolate changes resulting
254 from local vegetation cover change. As highlighted in the introduction, the
255 Australian climate is influenced by sea surface temperature in the tropical
256 Pacific and Indian Oceans, as well as pressure systems in the Southern Ocean
257 (BoM, 2012b). Risbey et al. (2009) compared five large-scale drivers, includ-
258 ing ENSO (measured by SOI and the Tropical Pacific SSTs), IOD, SAM,
259 MJO¹ (Madden-Julian oscillation) and blocking, in relation to Australian
260 rainfall variability. They identified SOI as the most important index among
261 all indices tested for broad parts of Australia (including QLD and NSW/VIC)
262 in almost any season. In this study, up to two important indices identified
263 from the seven climatic indicators (see section 3) were used, for each study
264 region, as the explanatory variables in the model .

265 As a preliminary step, the correlations between rainfall and each climatic
266 index were analysed. Rainfall in each study region was first deseasonalised
267 and detrended using the seasonal decomposition function "stl" in R (R Devel-
268 opment Core Team., 2011).  Detrending is needed to remove the correlations
269 between trends in the data as discussed in Smith and Timbal (2012). The
270 cross-correlations between the deseasonalised and detrended rainfall and the 
271 climatic indices were tested using the Pearson's product moment correla-
272 tion method, assuming the relationships are linear. Although the optimal
273 technique for exploring the correlation with each index could be different as

¹MJO is a large scale eastward-propagating wave-like disturbance in equatorial latitudes (Risbey et al., 2009).

described in Risbey et al. (2009), ~~the~~ Pearson's method was ~~applied to~~ all indices for consistency. Because the PDO is a low frequency descriptor of the multi-decadal SST (MacDonald and Case, 2005; Zanchettin et al., 2008; Kamruzzaman et al., 2011) a longer period (108 years, from 1900 to 2008), instead of 30-year rainfall data, was used to estimate the correlation with PDO, up to lag 24. For the other indices, the 30-year rainfall data set was used.

Based on the correlation between the climatic indices and rainfall (as shown in Figure 5 and Figure 6), the following ~~results were found~~:

- In QLD, the correlation between rainfall and SOI at zero time lags is the highest across all indices, outweighing the other ENSO indicators.
- In NSW/VIC, ~~again~~ the SOI has the highest correlation with rainfall, closely followed by the IOD. Both occur at ~~the~~ zero time lags.

The above findings are consistent with previous studies. Although some indices are serially correlated with rainfall up to several months, the lag zero events have the most significant correlation coefficients. Concurrent climatic index series were generally found most useful in rainfall prediction (e.g. Risbey et al., 2009; Kamruzzaman et al., 2011). The correlations between the climatic indices and rainfall for each individual season have also been tested. SOI was used in the Qld prediction, SOI and IOD were used in the NSW/VIC prediction.

Rainfall in Australia shows strong seasonal patterns (Holper, 2011; Australian Bureau of Statistics, 2012). For example, the north part of the country is summer rainfall dominant with dry winter, while most of the southern part has a winter rainfall regime. This ~~character is given~~ by the movement




Figure 5: Cross-correlation of the six indices and rainfall in QLD study region. Where the correlations with PDO are sought, 108-year rainfall data (1900 - 2008) are used. Otherwise, 30-year rainfall data are used. The correlation with NINO 3 is not shown as it is very similar to but weaker than the case of NINO 3.4. The blue dashed lines indicate the 95% confidence interval.

299 of subtropical high pressure system, which dominates the Australian climate
 300 (BoM, 2012a). The seasonal component of rainfall has a periodic pattern so
 301 it is better modelled as a smooth term. A spline function was applied on the
 302 ~~months to define a smooth seasonal pattern~~ (Wood, 2011).

303 Long term trends in the regional rainfall in some parts of Australia are




Figure 6: Cross-correlation of the six indices and rainfall in NSW/VIC study region.

304 significant (Hughes, 2003; Gallant et al., 2007; Chowdhury and Beecham,
 305 2010). In the northern and eastern parts of the continent, increasing rainfall
 306 is reported over the last century (Hughes, 2003). The presence of such long
 307 term trends may be confused with the outcome of a step change in rainfall.
 308 A linear trend term was implemented in the model to remove any long term
 309 trend effect.

310 We assumed all the factors are additive components in determining rain-
 311 fall as in Kamruzzaman et al. (2011). Generally, monthly rainfall has a

312 skewed distribution so the normality assumption of the residuals in a general
 313 linear model could be violated. In this case, the rainfall model is expressed
 314 as a generalised additive model (GAM) (Hastie and Tibshirani, 1986) with
 315 a log link function $g()$, assuming the residuals are gamma distributed (see
 316 Figure 7).

$$g(E(\mathbf{R}_r)) = \beta_0 + s_1(D_{1,r}\mathbf{SOI}) + s_2(D_{2,r}\mathbf{IOD}) + s_3(\mathbf{Season}) + \beta_1\mathbf{Trend} + \epsilon_r \quad (1)$$

317
 318 The bold letters represent the time series vectors. The subscript r denotes
 319 the region, $r = \text{qld}$ or nswvic . β_u ($u=0, 1$) are the fitted coefficients in the
 320 model. s_v ($v=1, 2, 3$) are the smooth functions on the climatic indices and
 321 the season. $D_{1,r}$ and $D_{2,r}$ switch on/off the corresponding climatic index in
 322 the model as discussed previously.

$$D_{1,r} = 1 \quad (2)$$

324 as SOI was used in both regions as rainfall predictor.

$$D_{2,r} = \begin{cases} 1 & \text{for NSW/VIC} \\ 0 & \text{for QLD} \end{cases} \quad (3)$$

326 The linear long term trend in the rainfall data is modelled by $\mathbf{Trend}=1,2,3\dots n$,
 327 where n is the total number of months in the time series. \mathbf{Season} is the sea-
 328 sonal component which is represented by applying a smooth spline function
 329 on the month values (1-12). The monthly rainfalls in both regions appear to
 330 have a lag 1 autocorrelation, being 0.26 for QLD and 0.33 for NSW/VIC.
 331 The SOI and IOD terms are also modelled with spline functions. As the
 332 effect of large scale drivers on Australian rainfall is more likely to be seasonal

333 (Murphy and Timbal, 2008; Schepen et al., 2012), the spline function can
 334 closely reproduce the high and low impacts of the climatic indices.




Figure 7: Distribution of monthly rainfall in (a) QLD and (b) NSW/VIC. By using a Kolmogorov-Smirnov test with shape = 1 and 2.4 respectively, rainfalls in both regions are shown to have a gamma distribution.

335 4.2. Tree cover change as factor variable

336 One of the main difficulties in empirical observation studies on the ef-
 337 fect of land cover change on rainfall is the lack of continuous monitoring of
 338 land surface variables, or even, no specific variable can be defined to clearly
 339 represent the land surface process. Given the lack of a full picture of the
 340 land surface process, a factor variable was used in this study to represent the
 341 abrupt land surface change (see Equation 5). The change could be a result
 342 of either land clearing or bushfires as long as it is permanent or takes a long
 343 time to recover. Here we approached the problem with two different models.

344 In the first method, the tree cover change was used as a predictor in the
 345 regression model, represented by a factor variable **LC**. The significance of
 346 the coefficient of **LC**, denoted as β'_2 in Equation 5, can be determined by a
 347 ratio test.

$$348 \quad LC = \begin{cases} \text{Trees} \\ \text{Removed} \end{cases} \quad (4)$$

349 Therefore in both regions, land cover is “trees” for the period before land
 350 cover change and “removed” for the period after the change. Here we sim-
 351 ply assumed that vegetation cover change has occurred on every pixel. The
 352 remaining term ϵ_r is the amount of rainfall that is attributed to other un-
 353 specified factors and random errors. Hence the regression model becomes
 354

$$g(E(\mathbf{R}_r)) = \beta'_0 + s'_1(D_{1,r}\mathbf{SOI}) + s'_2(D_{2,r}\mathbf{IOD}) + s'_3\mathbf{Season} + \beta'_1\mathbf{Trend} + \beta'_2\mathbf{LC} + \epsilon'_r \quad (5)$$

355
 356 Vegetation cover changes occurred at different times in the two regions.
 357 In the QLD region, there no exact time can be assigned to the land clear-
 358 ing. Clearing has occurred between 2003 - 2004 according to SLATS reports.
 359 The information on the change in type of land cover during this time period
 360 is missing. Therefore, four scenarios were tested in the analysis. In these
 361 scenarios the “after change” period started from: (1) June 2003, (2) January
 362 2004, (3) June 2004 and (4) January 2005. In the NSW/VIC region, se-
 363 vere bushfires were reported in early January 2003. Hence the “tree” cover
 364 state was up to December 2002 then it was changed to “removed” state from
 365 January 2003. The regression model was run from 1979 for both regions.

366 4.3. Step trend test

367 As a second method, a step trend test was used to detect changes in rain-
368 fall after the vegetation cover change. This nonparametric statistical test
369 was modified from the Mann-Whitney Rank-Sum test by Hirsch and Gilroy
370 (1985). The test was developed to look for a step change in data which is
371 cross correlated. As the gridded rainfall dataset is based on an interpolation
372 method, this results in a high spatial correlation between neighbouring pix-
373 els. The step trend test is suitable for the analysis of such a dataset. The
374 advantages of this test are: (1) it does not depend on assumptions of the
375 data distribution; (2) it is not restricted to datasets with no missing data;
376 (3) it is robust and not as easily influenced by outliers and negative numbers
377 (Hirsch and Gilroy, 1985).

378 4.3.1. Rainfall residuals

379 The rainfall residuals from the regression model in Equation 1 were used
380 for this test. According to Hirsch and Gilroy (1985), using deseasonalised and
381 detrended data is important in a test to detect step change. Furthermore,
382 since rainfall is only partially attributed to local sources and conditions, noise
383 can be introduced by large scale dynamics and changes in other climatic
384 factors. The assumption is that the regression model should remove this
385 noise. Hence the model residuals ϵ'_r exclude the effects due to climate drivers,
386 seasonality and the long term trend. As a result local effects are assumed
387 to explain at least part of the variation in the residuals. The test, described
388 in the following section, associates changes in rainfall with the tree cover
389 changes.

390 *4.3.2. Mann-Whitney rank-sum statistic*

391 The step trend test is a modified version of the Mann-Whitney rank-sum
 392 statistic (Hirsch and Gilroy, 1985). As a nonparametric rank-based test, the
 393 Mann-Whitney test does not use the exact values of rainfall but depends on
 394 the ranks of the data. For each month, rainfall residuals of each year were
 395 ranked in an ascending order. The ranking of January rainfall in a sample
 396 pixel k in QLD is illustrated below:

	1998	1999	2000	2001	2002	2005	2006
Rainfall residuals:	-0.3	-60.9	-16.1	-71.7	111.1	-7.2	-60.5
Rank (R'_{1k}):	6	2	4	1	7	5	3

397 Therefore, the smallest value (including negative values) has rank 1 and
 398 the largest value has the maximum rank.

399 The before period and the after period formed two groups of samples.
 400 The split point of the two periods was based on the timing of the vegetation
 401 cover changes. In the QLD region, changes occurred anytime during 2003
 402 and 2004. In contrast to the previous method, the time period of where the
 403 land cover change occurred was excluded, as the nonparametric test allows
 404 missing data. Hirsch and Gilroy (1985) also pointed out that the power of
 405 the test is higher if the data of the change period is ignored. Hence 2003 and
 406 2004 were excluded from the analysis. As a result, the after-change period
 407 was 2005 - 2007.

408 In the case of NSW/VIC, the bushfires broke out in early January 2003.
 409 The change was within a relatively short period of the year. Therefore the
 410 after-change period in this region still started in January 2003. The before

411 period was set to five years (1998 - 2002) in both regions.

412 The rank of rainfall in month j year i in pixel k is denoted as R'_{ijk} . The
 413 sum of ranks of rainfall in month j in pixel k before the known intervention
 414 is:

$$415 \quad W_{jk} = \sum_{i=1}^{n_1} R'_{ijk}. \quad (6)$$

416 n_1 is the number of years before the land cover change. The expected value
 417 of W_{jk} is

$$418 \quad \mu_w = n_1(n_1 + n_2 + 1)/2 \quad (7)$$

419 n_2 is the number of years after the change. Hence the expected value of the
 420 rank sum before the intervention is the same for all months and all pixels.
 421 The sum of ranks for the whole time period is fixed, as $(n_1 + n_2)(n_1 + n_2 + 1)/2$.
 422 Knowing the rank-sum of one group the rank-sum of the other group can be
 423 easily derived. If the rainfall data is temporally and spatially independent,
 424 the variance of W_{jk} is

$$425 \quad \sigma_w^2 = n_1 \cdot n_2(n_1 + n_2 + 1)/m \quad (8)$$

426 where m is the number of months which is 12 in the case of a full year.

427 *4.3.3. Step trend test*

428 Instead of completing the Mann-Whitney U-test, Hirsch and Gilroy (1985)
 429 applied the rank-sum statistics in a standard normal Z test. The modified
 430 test can be used to detect step change and it accounts for serial and cross
 431 correlation in the data. In the case here, the deseasonalised and detrended
 432 data shows little autocorrelation in the time series but possesses strong cross
 433 correlation between neighbouring pixels, i.e. $R > 0.99$. Hence the covariance
 434 between pixels would need to be considered in the test.

435 The sum of W_{jk} for a block of ns pixels over the whole year, $\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}$,
 436 has a mean value of

$$437 \quad E\left(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}\right) = 12 \cdot ns \cdot \mu_W \quad (9)$$

438 and variance

$$439 \quad Var\left(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}\right) = \sum_{j=1}^{12} \sum_{k=1}^{ns} \sum_{h=1}^{ns} C(W_{jk}, W_{jh}). \quad (10)$$

440 $C(W_{jk}, W_{jh})$ is the covariance of the W statistics between pixel k and pixel
 441 h in month j . When $k = h$, $C(W_{jk}, W_{jh}) = \sigma_w^2$. When $k \neq h$,

$$442 \quad C(W_{jk}, W_{jh}) = \sigma_w^2 r(R_k, R_h) \quad (11)$$

443 where $r(R_k, R_h)$ is the product moment correlation coefficient of the concur-
 444 rent ranks in pixel k and h . Here r is calculated on the full time series in
 445 each pixel. In this analysis, the test was applied to a square block of four
 446 pixels each time. As argued by Hirsch and Gilroy (1985), $ns = 4$ is the most
 447 optimal solution to balance the cost and the gain in the test power.

448 The statistic of the step trend test is then defined as

$$449 \quad Z' = \frac{\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk} - 12 \cdot ns \cdot \mu_w}{\sqrt{Var(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk})}}. \quad (12)$$

450 The above statistic is written for a 12 month period. The statistic can also
 451 be used to test seasonal rainfall change or for other customized periods, by
 452 changing the value 12.

453 The null hypothesis (H_0) in this study is that there was no change in
 454 rainfall due to land surface intervention. The results of the step trend test
 455 can be interpreted according to the sign of Z' score (see Table 1). Z' is

Table 1: The interpretation of Z' score in the step trend test. Following Hipel and McLeod (1994, Chapter 23, P887).

$Z' > 0$	rainfall decreases after change
$Z' < 0$	rainfall increases after change
$Z' = 0$	rainfall does not change

456 normally distributed similar to the standard normal statistics Z . Hence it
457 can be compared to a standard normal distribution to determine the p value.

458 5. Results

459 5.1. Tree cover changes

460 The pixels with significant tree cover change in each study region are
461 shown in Figure 8, at the $p < 0.1$ significance level. Despite the limitations of
462 the length of this data series, it still indicates a large change in the NSW/VIC
463 region. In the NSW/VIC region, much of the tree loss between 2002 and 2003
464 was in the Snowy Mountains which are at the border of NSW and VIC. Tree
465 cover loss occurred in large parts of the QLD study region between 2002
466 and 2005. Most of the clearings appear in the centre of this region. The
467 tree cover change map is consistent with the annual mean EVI trend map
468 (based on DLCD data, map not shown here), which confirms these changes
469 are significant and persistent over the study period.




Figure 8: The maps show the areas with significant changes in tree cover (at $p < 0.1$) in (a) the QLD region and (b) the NSW/VIC region. The amount of change was calculated as the difference in tree covers before and after the specified land cover intervention and it is shown as the percentage of the ground area. Green colour indicates an increase in tree cover, while red colour indicates a decrease in tree cover.

470 5.2. Regression Model & Significance of Vegetation Cover Changes

471 The regression model does not explain much of the rainfall variability.
 472 The model in Equation 5 accounts for around 13%² of the rainfall variations
 473 in the QLD region and 19% in the NSW/VIC region on average. The resid-
 474 ual analysis shows that the assumptions of the regression model are generally
 475 met. The standardised residual plots, however, show some funnelling for the
 476 NSW/VIC regions, suggesting non constant variance. The residual analy-
 477 sis, based on one sample pixel from each region, in Figure 9 and Figure 10
 478 illustrates this. The residual patterns are consistent within each region.

²Here the adjusted R^2 was reported. Adjusted R^2 is the coefficient of determination, a measurement of the amount of variability predicted by the model adjusting for the number of explanatory terms

479 The model confirms the importance of the climate drivers and the sea-
 480 sonality in Australian rainfall. Even at the grid level, the seasons and the
 481 climatic indices were significant ($p < 0.05$) in both regions. The explaining
 482 power of the model is mostly due to these variables. The climate drivers (at
 483 lag zero) accounted, on average, for 6% of the rainfall variability in both the
 484 QLD region and the NSW/VIC region (see Figure 11 for the distribution
 485 of R^2 in these two regions). These figures are within the upper bound of
 486 seasonal rainfall predictability by SST anomaly field reported by Westra and
 487 Sharma (2010).




Figure 9: The residual analysis of a sample pixel in the QLD region.




Figure 10: The residual analysis of a sample pixel in the NSW/VIC region.

488 A statistically significant long term trend was not observed in either of
 489 the two regions. However, this result does not disprove the importance of a
 490 long term trend in rainfall. The overall time period is fairly short and more
 491 pixels in NSW/VIC would have a significant step change if the long term
 492 trend effect is not removed by the model. As trend free data is an important
 493 requirement for the step trend test, the trend term was kept in the regression
 494 model to ensure the detection of step change was not due to a possible long
 495 term trend.

496 The land cover variable implies a step change with different values before




Figure 11: The distribution of R^2 when rainfall is only modelled by the climate drivers. It shows the percentage of rainfall variability that can be explained by the climate drivers. In the case of (a) the QLD region, SOI was the only climatic index considered. In (b) the NSW/VIC region, SOI and IOD were used.




Figure 12: The spatial distribution of significance of vegetation cover changes in the two study regions. The significance of vegetation cover changes is assessed in the regression model. The figures report the p values of the coefficient of the land cover variable in the model at different significance level. The plot of the QLD region is for the scenario that the after change period started from January 2005, as the other scenarios don't have any p value less than 0.2.

497 and after the land cover intervention. This variable was only significant
 498 ($p = 0.05$) for the rainfall estimates in some areas in NSW/VIC, as shown

499 in Figure 12 (b). The effect of tree removal in the NSW/VIC region was
500 highly significant in the area from the Snowy Mountains range in the south
501 to the west side of Australian Capital Territory (ACT), highlighted by the
502 red colour in Figure 12 (b). However no significant step change (at $p < 5\%$)
503 due to the land cover changes was found in rainfall of the QLD region in any
504 of the four scenarios.

505 In the NSW/VIC region, the locations of rainfall change show some agree-
506 ment with the locations of vegetation cover change. Comparing the results
507 in Figure 12 (b) to Figure 8 (b), the alpine area with massive tree cover
508 loss is located inside the highlighted region. The results show that the step
509 change in rainfall also occurred in the cotter river catchment which was heav-
510 ily burned in 2003 bushfires (refer to Figure 2). However, Figure 12 (b) also
511 shows that a significant step change in rainfall was found in an area larger
512 than where the bushfires has occurred. From this point of view, the results
513 might be showing a large scale effect that extends beyond the vegetation
514 cover change effect.

515 More generally, the model also shows that the tree cover has a positive
516 impact on rainfall. The fitted coefficients for the reference state "trees" were
517 consistently positive for the pixels whose step changes of rainfall were signif-
518 icant. It implies that rainfall was higher when the surface was covered by
519 trees. This was confirmed by applying a two sample Student's t-test on the
520 annual rainfall of the before-change and after-change periods.

521 5.3. Step Trend Test

522 The spatial step trend test Z' scores are shown in Figure 13. This figure
523 provides two types of information: the sign and the significance level. The




Figure 13: Spatial distribution of the step trend test Z' statistics in the two study sites. The test was conducted on the 30-year rainfall data from 1979 to 2009. Warm colours (yellow, orange and red) are for positive Z' values which indicate decreasing rainfall trend due to the land surface intervention. Cold colours (light blue to blue) are for negative Z' values which indicate increasing rainfall trend. The deeper the colour, the more significant the statistic.

sign indicates the direction of the step change, as listed in Table 1. In each region, there is a broad area of positive Z' values which implies a decrease in rainfall. In QLD, the areas of positive Z' , especially when p is smaller than 0.1, almost align with the locations reported to have heavy land clearing (see Figure 14). However the vegetation cover change map in Figure 8 (a) does not indicate a clear similar pattern. In the Snowy Mountains area and the west of ACT, where severe bushfires occurred in 2003, positive Z' values were also found. It again indicates that loss of tree cover was related to a rainfall decrease. Nevertheless the observed step change in rainfall was not strong. At the 5% significance level, changes in a number of pixels in the alpine area are possibly where the tree recovery is slow. In the QLD region, 35 pixels obtained a Z' score with $p < 5\%$. In the NSW/VIC region, there are only

18 pixels in the alpine area having a Z' score with $p < 5\%$. They are a very small proportion of both study regions. As the variability of the monthly rainfall, and the variability of rainfall in Australia is considerable (Delworth and Zeng, 2014), detecting significant trends at $p < 5\%$ could be difficult. Therefore, an analysis based on $p < 10\%$ is presented in the figures. At $p = 10\%$, 278 pixels in the QLD region had a significant positive Z' score, while 142 pixels in the NSW/VIC region were significant. In both cases this was a bit over 10% of the total area.




Figure 14: Woody vegetation clearing rate in QLD. The maps and figures are obtained from Department of Natural Resources and Water (2007). The red rectangle is the boundary of the QLD study region.

The rainfall residuals of the two periods (before-change since 1998 and after-change) were also compared (Figure 15). The annual values of the




Figure 15: Boxplots of annual rainfall (estimated based on rainfall residuals from Equation 1) before and after the land cover intervention during 1998 - 2008 in the study regions. On average, the after period has a lower annual rainfall amount and with outliers of small values.

11-year period (1998 - 2008) were calculated for each pixel. The boxplots indicate that there were different mean values between the “before” and “after” periods in both regions. There are small outliers in the after period in NSW/VIC. A close look at the rainfall data revealed that rainfall was consistently very low in 2006 for the whole region. The low rainfall in 2006 was due to a weak El Niño and memory effects of the previous drought. While

the regression model has removed most of the effect of ENSO, the cumulative drought effect could still be visible in the residuals if the response of the rainfall to the ENSO effect is non-linear, and the memory of the past drought is persistent. This can also explain the significant results in the previous method. To remove the outlier effects, the two periods were compared excluding the 2006 rainfall. An F test confirmed that the variances were different between the before and after periods. Hence an unpaired unequal variance two sample t-test was applied to test whether the after period has lower mean rainfall than the before period, for the group of pixels showing a negative step change in rainfall ($p = 0.05$) and the group of pixels with no change. The rainfall in the after period was lower than the before period ($p < 0.05$) for pixels showing a negative step change. For those pixels without a step change, there was no statistical evidence that the two periods have different means. Hence the t-test results are consistent with the step trend test results.

The choice of ns has some impact on the test results, as shown by Hirsch and Gilroy (1985). The cases of $ns = 1$ and $ns = 9$ were also tested. When $ns = 1$, 18 pixels in the QLD region obtained a Z' score at the 5% significance level. In the NSW/VIC region, the detection of negative step change ($p = 0.05$) reduced to 16 pixels. On the other hand, when $ns = 9$, the results were similar to the case of $ns = 4$. In this case, the changes ($p = 0.05$) were detected in 35 pixels in the QLD region and 18 pixels in the NSW/VIC region. The power of the test does not change much after $ns = 4$, as shown by Hirsch and Gilroy (1985). The cause of the somewhat inconsistent results in the QLD region between $ns = 1$ and $ns = 4$ is unclear. This could possibly

577 be because the higher ns values smooth the individual pixel results.

578 The "field significance" of the test is considered to make inferences about
579 the step change at regional scales from multiple local tests (Wilks, 2006;
580 Westra et al., 2013). Here, the bootstrapping resampling method from Wes-
581 tra et al. (2013) was modified to evaluate the field significance. The spa-
582 tial structure of the pixels was maintained, while the order of the years and
583 months, were changed by random resampling. The test statistic identifies the
584 percentage of the pixels with significant step change, positive and negative
585 respectively, for the step trend test. The test statistics on 1000 resampled
586 replicates were used to develop the distribution of these percentage values
587 under the local null hypothesis that there was no step change.

588 The bootstrapping resampling technique was applied on both $ns = 1$ and
589 $ns = 4$ cases. In the $ns = 4$ case, a spatial moving block bootstrapping
590 was used, in which the change of time sequence within the 2×2 block was
591 consistent. The distribution of the test statistics is highly concentrated at
592 zero with a skew to the right. In the outputs, negative step changes in the
593 cases of $ns = 1$ and $ns = 4$, the test statistics on the observed time series
594 generally fall within the null hypothesis distribution (the cases of $ns = 4$ are
595 shown in Figure 16 and Figure 17). However, the bootstrap distribution is
596 very wide and highly skewed, with step change values close to 0. The result
597 for both the positive Queensland and positive NSW/VIC $ns = 4$ step change
598 tests are located much further to the right on the tail of the distribution,
599 away from 0 and give a stronger indication of a change. The results showed
600 that overall the chance of detecting a rainfall change was small, which is
601 related to the strong natural variability of the rainfall in Australia. However,

602 the location of the positive tests on the tail of the distribution suggests some
 603 support for the hypothesis that vegetation cover change affects local rainfall.




Figure 16: Percentage of pixels showing statistically significant positive Z' values (left) and negative Z' values (right) in QLD at $p < 0.1$. The histogram shows the distribution of results from 1000 bootstrap resampling of the rainfall time series. The red dot represents the results from the observed data.

604 6. Summary and Discussion

605 Generally, empirical studies on LCC-precipitation interaction are con-
 606 ducted within an area with known land surface intervention (e.g. Otterman
 607 et al., 1990; Durieux et al., 2003; Negri et al., 2004; Sato et al., 2007). How-
 608 ever, these locations are rare and difficult to isolate from real landscape
 609 change. In this study we therefore tested the effect of land cover change
 610 across a broad region, rather than only for locations where changes were
 611 known to occur or have occurred. The advantage of this approach is that it
 612 does not require a long time series of land cover data which is usually un-
 613 available. Furthermore, it does not assume a specific relationship between




Figure 17: Percentage of pixels showing statistically significant positive Z' values (left) and negative Z' values (right) in NSW/VIC at $p < 0.1$. The histogram shows the distribution of results from 1000 bootstrap resampling of the rainfall time series. The red dot represents the results from the observed data.

614 vegetation cover change and rainfall but allows the data to show this rela-
615 tionship, by applying the analysis to a broader area outside the boundary
616 of the vegetation cover change. This approach is expected to provide a way
617 to reduce the risk of a false positive paradox, by comparing results between
618 areas with and without vegetation cover change.

619 Parametric tests are generally more powerful than nonparametric test in
620 detecting a trend, when the data is normally distributed (Onoz and Bayazit,
621 2003; Kundzewicz and Robson, 2004). As a non-parametric test, the step
622 trend test has the advantages of distribution free and having no restriction
623 on missing data (Hirsch and Gilroy, 1985). This is particularly useful in
624 rainfall analysis since rainfall data is usually skewed. On the other hand, the
625 disadvantages of non-parametric tests, such as being limited to hypothesis
626 testing and weaker in power, also hold for the step trend test (Whitley and

627 Ball, 2002).

628 The regression model used here is a simple model. We only consider the
629 important effects of historical trend, seasonality and climate drivers. Fur-
630 thermore, no more than two large scale climatic indices were used in order to
631 avoid over-parameterisation and multiple cross-correlations between climatic
632 indices. A purpose of the regression model is to remove the variability in
633 rainfall that is due to these known important factors. The model shows that
634 seasonality, ENSO and IOD together explain no more than 20% of the rain-
635 fall variability, and around 6 - 8% on average is due to the climate drivers.
636 This is consistent with the literature (e.g. Westra and Sharma, 2010). A
637 large amount of variation is left in the model residuals, which is likely due
638 to random factors. Rainfall is generally considered a stochastic process (e.g.
639 Fowler et al., 2005; Cowpertwait et al., 2009; Burton et al., 2010). The high
640 variability increases the difficulty to detect a change in rainfall.

641 There was some evidence that a step change has occurred in the QLD
642 region. The semi-parametric model could not identify any step change in the
643 rainfall data. On the other hand, the step trend test suggested a couple of
644 locations (300 - 500 km²) might have experienced rainfall change ($p = 0.05$).
645 The results from the step trend test also indicated a possible widespread
646 change, but the spatial pattern of change is not consistent with the tree
647 cover change map. Although land clearings in QLD have occurred at a high
648 rate and broad scale (Department of Natural Resources and Water, 2008),
649 this study cannot confirm its impact on the local rainfall. At this location,
650 land cover change was not found to change the mean rainfall significantly in
651 some other studies (e.g. Narisma and Pitman, 2003; McAlpine et al., 2007).

652 The characteristics of vegetation cover changes in QLD might increase the
 653 difficulty in detecting a step change. QLD has a long history of land clearing.
 654 According to the series of SLATS reports on land cover changes in QLD re-
 655 leased by the Queensland government, land clearing continued in and around
 656 the study region between 1988 - 2008. Major broad scale and high rate clear-
 657 ings occurred in 1999 - 2000 and 2002 - 2004. It is therefore difficult to define
 658 a clear cut change in this region. The more continuous ongoing land clear-
 659 ing could have reduced the significance of a step change. The two methods
 660 indicate quite different results on the level of change in the NSW/VIC re-
 661 gion. The semi-parametric model showed that a large area in the NSW/VIC
 662 region has experienced significant step change ($p = 0.05$) in rainfall after
 663 2003. But this result is likely to be affected by the low rainfall in 2006 as
 664 the area is larger than the bushfires region. The step trend test was able to
 665 detect significant changes ($p = 0.05$) in about 400 km² areas in the Snowy
 666 Mt. and some possible changes along the bushfires region ($p = 0.1$). This
 667 might be due to the possible small size of the step change and/or a short
 668 after-fire period (Hirsch and Gilroy, 1985). The vegetation cover change due
 669 to bushfires might have possibly changed the rainfall but the evidence was
 670 not strong either.

671 The bushfires locations highlighted in the analysis results is an interesting
 672 outcome. The results might be due to a drought effect. The 2002 - 2003
 673 drought has affected rainfall in a large part of Australia, particularly the
 674 MDB (Nicholls, 2004). The severe bushfires in 2003 were also triggered by
 675 the extreme drought conditions. Although the dry episode effect on rainfall
 676 has been accounted for in part by the SOI, further impact of drought could be

677 passed through the local land-atmosphere interaction. However, this impact
678 might not be statistically significant, even under an extreme event such as
679 bushfires. Overall, the rainfall feedback to the vegetation cover change could
680 be weak under dry conditions.

681 The different causes of vegetation cover change in these two regions could
682 lead to different post-change characteristics. The magnitude of EVI decreasing
683 trends in the QLD region are less than in NSW region, as reported in the
684 DLCD data. This is due to the lower tree density in the QLD region than
685 in the NSW/VIC region before land surface interventions. Wild fires might
686 have totally damaged the vegetation cover and recovery was very slow in some
687 areas within the Snowy Mt. The persistent drought in the 2000s (Howden,
688 2012) has delayed the regrowth of trees. On the other hand, replacing tree
689 cover with pasture and crops might have a relatively subtle impact on the
690 EVI. On the other hand, regrowth of trees can be expected after fires or as
691 drought condition is relieved; but clearings for agricultural purpose impose
692 permanent or semi-permanent changes to the land surface.

693 The rainfall data used in this study is a gridded data set. This data
694 set is robust and consistent over a long time series (from 1900 to current)
695 and has a broad national wide coverage which can provide more information
696 spatially. However, high cross correlation between pixels due to the interpolation
697 method generating this data set can also introduce spatial noise. Here
698 the cross correlation has been account for in the step trend test. Some other
699 methods are also available which can be used to perform a comparative trial.
700 For example, Narisma et al. (2007) applied a spatial Gaussian filter on a similar
701 data set and used wavelet analysis to detect step change in rainfall. High

quality station data is another option to test whether the observed spatial pattern in the step trend test results was not due to the gridded data itself. Resampling methods, such as bootstrapping and permutation (Wilks, 1997; Kundzewicz and Robson, 2004; Westra et al., 2013), can also be used to further assess the strength of significance of results and incorporate spatial and temporal patterns in the analysis. We are also aware that the gridded data set is most useful in regions with sparse rain gauge networks but it actually reduces information where the rain gauge density is high (Jones et al., 2009). In the Snowy Mountains area, the coverage of rainfall stations is intensive but they are mainly located in the valleys. The interpolated data might not best represent the local rainfall.

Overall, the inclusion of the anomalous rainfall year 2006, could have influenced the results, and this means that strong conclusions cannot be drawn.

However, the current study provides some evidence to reject the null hypothesis (no step change in rainfall is due to the tree cover loss). Limited by the available data, the time frame under study was chosen within a long lasting drought period (Holper, 2011). The strong impact of this prolonged drought might have suppressed the land-atmosphere interaction and confused the cause and effect relation between rainfall and vegetation cover change. This could be one of the reasons that the LCC effects found in other studies (Görgen et al., 2006; McAlpine et al., 2007, e.g.) are not found to be significant here. So possible future work can be conducted for a non-drought period, when a longer series of land cover data is available. This approach should be further assessed in different scenarios of wet period and/or af-

727 forestation. Continuous monitoring of land surface conditions is important
728 for future research. The power of the test can also be improved with the
729 longer length of the after-intervention period (Hirsch and Gilroy, 1985). But
730 then the question is what is an appropriate length for land cover change
731 study, when ongoing change is possible, such as tree regrowth or cropping.
732 This will require further research.

733 **7. Conclusion**

734 In this study, we found some observationl data based evidence, although
735 not strong, that vegetation cover change has changed local rainfall. The
736 semi-parametric method and non-parametric method did totally agree on
737 detecting a significant step change in rainfall in the hot dry QLD region
738 where land clearing has occurred. On the other hand, the bushfires in the
739 humid, temperate mountain range in the NSW/VIC region has experienced
740 reduced rainfall. But the dry spell also plays an important role in the results.

741 Drought has had a pronounced impact on the land surface condition dur-
742 ing the study period, leading to significant reduction in vegetation and ex-
743 treme events such as bushfires. The associated lack of rainfall and high
744 temperatures may mask the step change in the vegetation. Hence, the sig-
745 nal of LCC feedback on rainfall is probably weaker under such regional dry
746 conditions, as the impact of LCC on rainfall is mainly through changes in
747 moisture convergence (Görge et al., 2006; Pitman and Hesse, 2007).

748 **8. acknowledgments**

749 CL was supported by an Australian Postgraduate Award for this work.

750 **Appendix A. Summary of Data**

Table A.2: Summary of data.

Data	Source	Resolution		Analysis period
		Temporal	Spatial	
Percent tree cover	MOD44B	Annual	250m	2000-2010
Trend of vegetation cover change	DLCD (2009)	Onetime	250m	Trend of Apr 2000 - Apr 2008
Rainfall	AWAP gridded rainfall data	Monthly	$0.05^{\circ} \times 0.05^{\circ}$	Jan 1979- Dec 2008
SOI	BoM	Monthly	N/A	Jan 1979- Dec 2008
NINO 3, 3.4, 4	IRI/LDEO data library	Monthly	N/A	Jan 1979- Dec 2008
PDO	NOAA	Monthly	N/A	Jan 1979- Dec 2008
IOD	POAMA-2 dataset	Monthly	N/A	Jan 1979- Dec 2008

751 ABARES, 2010. Land use of Australia, Version 4, 2005/2006.

752 ABC News, 2003. Kosciuszko slow to recover from bushfires. 30 Oct,
753 2003. Available online at url[http://www.abc.net.au/news/2003-](http://www.abc.net.au/news/2003-10-30/kosciuszko-slow-to-recover-from-bushfires/1501722)
754 10-30/kosciuszko-slow-to-recover-from-bushfires/1501722, viewed on
755 06/06/2012.

756 Australian Bureau of Statistics, 24/05 2012. Geography and climate - Aus-
757 tralia's climate.

758 Ben-Gai, T., Bitan, A., Manes, A., Alpert, P., Rubin, S., 1998. Spatial and
759 temporal changes in rainfall frequency distribution patterns in Israel. The-
760 oretical and Applied Climatology 61 (3-4), 177–190.
761 URL <GotoISI>://000078025000005

762 BoM, 2009. Gridded Monthly Rainfall Metadata. Australian Bureau of
763 Meteorology.
764 URL [http://www.bom.gov.au/climate/austmaps/](http://www.bom.gov.au/climate/austmaps/metadata-monthly-rainfall.shtml)
765 [metadata-monthly-rainfall.shtml](http://www.bom.gov.au/climate/austmaps/metadata-monthly-rainfall.shtml)

766 BoM, 2012a. Australia - climate of our continent. Available online
767 at [http://www.bom.gov.au/lam/climate/levelthree/ausclim/zones.](http://www.bom.gov.au/lam/climate/levelthree/ausclim/zones.htm)
768 [htm](http://www.bom.gov.au/lam/climate/levelthree/ausclim/zones.htm), viewed on 12/03/2012.

769 BoM, 2012b. Australian climate influence. Available online at
770 [http://www.bom.gov.au/watl/about-weather-and-climate/](http://www.bom.gov.au/watl/about-weather-and-climate/australian-climate-influences.shtml)
771 [australian-climate-influences.shtml](http://www.bom.gov.au/watl/about-weather-and-climate/australian-climate-influences.shtml), viewed on 17/01/2012.

772 BoM, 2012c. Background information, National rainfall outlook. Available

773 online at http://www.bom.gov.au/climate/ahead/rain_ahead.shtml,
774 viewed on 03/01/2012.

775 BoM, m., 2008. Weather drivers in Western Australia.
776 Available online at [http://www.mla.com.au/files/](http://www.mla.com.au/files/1c39e89d-f3fa-4915-b72b-9d66008ab4bf/)
777 [1c39e89d-f3fa-4915-b72b-9d66008ab4bf/](http://www.mla.com.au/files/1c39e89d-f3fa-4915-b72b-9d66008ab4bf/), viewed on 03/02/2012.

778 Bosilovich, M. G., Chern, J.-D., June 01, 2006 2006. Simulation of water
779 sources and precipitation recycling for the MacKenzie, Mississippi, and
780 Amazon River Basins. *Journal of Hydrometeorology* 7 (3), 312–329.
781 URL <http://dx.doi.org/10.1175%2FJHM501.1>

782 Burton, A., Fowler, H., Blenkinsop, S., Kilsby, C., 2010. Downscaling tran-
783 sient climate change using a neyman-scott rectangular pulses stochastic
784 rainfall model. *Journal of Hydrology* 381 (1), 18–32.

785 Cai, W., Purich, A., Cowan, T., van Rensch, P., Weller, E., Jan. 2014. Did
786 Climate Change-Induced Rainfall Trends Contribute to the Australian
787 Millennium Drought? *Journal of Climate* 27 (9), 3145–3168.
788 URL [http://journals.ametsoc.org/doi/abs/10.1175/](http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-13-00322.1)
789 [JCLI-D-13-00322.1](http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-13-00322.1)

790 Chagnon, F. J. F., Bras, R. L., Jul 2005. Contemporary climate change in
791 the Amazon. *Geophysical Research Letters* 32 (13).
792 URL [http://www.agu.org/](http://www.agu.org/journals/gl/g10513/2005GL022722/2005GL022722.pdf)
793 [journals/gl/g10513/2005GL022722/2005GL022722.pdf](http://www.agu.org/journals/gl/g10513/2005GL022722/2005GL022722.pdf)

794 Chowdhury, R. K., Beecham, S., 2010. Australian rainfall trends and their
795 relation to the Southern Oscillation Index. *Hydrological Processes* 24 (4),

796 504–514.

797 URL <http://dx.doi.org/10.1002/hyp.7504>

798 Cowpertwait, P., Salinger, J., Mullen, B., 2009. A spatial-temporal stochastic
 799 rainfall model for Auckland City: Scenarios for current and future climates.
 800 Journal of Hydrology (NZ) 48, 95–109.

801 DeAngelis, A., Dominguez, F., Fan, Y., Robock, A., Kustu, M. D., Robin-
 802 son, D., Aug 2010. Evidence of enhanced precipitation due to irrigation
 803 over the Great Plains of the United States. Journal of Geophysical
 804 Research-Atmospheres 115 (D15).

805 URL [http://www.agu.org/
 806 journals/jd/jd1015/2010JD013892/2010JD013892.pdf](http://www.agu.org/journals/jd/jd1015/2010JD013892/2010JD013892.pdf)

807 Delworth, T. L., Zeng, F., Aug. 2014. Regional rainfall decline in Australia
 808 attributed to anthropogenic greenhouse gases and ozone levels. Nature
 809 Geoscience 7 (8), 583–587.

810 URL [http://www.nature.com/ngeo/journal/v7/n8/full/ngeo2201.
 811 html?foxtrotcallback=true](http://www.nature.com/ngeo/journal/v7/n8/full/ngeo2201.html?foxtrotcallback=true)

812 Deo, R., Syktus, J., McAlpine, C., Wong, K., 2009. The simulated impact of
 813 land cover change on climate extremes in eastern australia. In: Proceed-
 814 ings of the 18th World IMACS Congress and MODSIM09 International
 815 Congress on Modelling and Simulation. Modelling and Simulation Soci-
 816 ety of Australia and New Zealand Inc. and International Association for
 817 Mathematics and Computers in Simulation, pp. 2035–2041.

818 Department of Natural Resources and Mines, 2005. Land cover change in

819 Queensland 2001-2003, incorporating 2001-2002 adn 2002-2003 change pe-
820 riods: A statewide landcover and trees study (SLATS) report. Tech. rep.,
821 Department of Natural Resources and Mines, Brisbane, Feb, 2005.

822 Department of Natural Resources and Mines, 2006. Land cover change in
823 Queensland 2003-2004: A statewide landcover and trees study (SLATS)
824 report. Tech. rep., Department of Natural Resources and Mines, Brisbane,
825 Feb, 2006.

826 Department of Natural Resources and Water, 2007. Land cover change in
827 Queensland 2004-2005: A statewide landcover and trees study (SLATS)
828 report. Tech. rep., Department of Natural Resources and Water, Brisbane,
829 Feb, 2007.

830 Department of Natural Resources and Water, 2008. Land cover change in
831 Queensland 2005-06: A statewide landcover and trees study (SLATS) re-
832 port. Tech. rep., Department of Natural Resources and Water, Brisbane,
833 Feb, 2008.

834 Dirmeyer, P. A., Brubaker, K. L., DelSole, T., Feb 2009. Import and export
835 of atmospheric water vapor between nations. *Journal of Hydrology* 365 (1-
836 2), 11–22.

837 URL <GotoISI>://000263575600003

838 Durieux, L., Machado, L. A. T., Laurent, H., 2003. The impact of deforesta-
839 tion on cloud cover over the Amazon arc of deforestation. *Remote Sensing*
840 of Environment 86 (1), 132–140.

841 URL [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S0034425703000956)
842 [S0034425703000956](http://www.sciencedirect.com/science/article/pii/S0034425703000956)

843 Eltahir, E. A. B., Bras, R. L., Aug 1996. Precipitation recycling. *Reviews of*
844 *Geophysics* 34 (3), 367–378.

845 URL <GotoISI>://A1996VD58700003

846 Fowler, H., Kilsby, C., O’connell, P., Burton, A., 2005. A weather-type con-
847 ditioned multi-site stochastic rainfall model for the generation of scenarios
848 of climatic variability and change. *Journal of Hydrology* 308 (1), 50–66.

849 Gaertner, M. A., Christensen, O. B., Prego, J. A., Polcher, J., Gallardo, C.,
850 Castro, M., 2001. The impact of deforestation on the hydrological cycle in
851 the western Mediterranean: An ensemble study with two regional climate
852 models. *Climate Dynamics* 17 (11), 857–873.

853 Gallant, A. J. E., Hennessy, K. J., Risbey, J., 2007. Trends in rainfall indices
854 for six Australian regions: 1910 - 2005. *Australian Meteorological Magazine*
855 56, 223 – 239.

856 Gimeno, L., Drumond, A., Nieto, R., Trigo, R. M., Stohl, A., 2010. On the
857 origin of continental precipitation. *Geophysical Research Letters* 37 (13).
858 URL <http://dx.doi.org/10.1029/2010GL043712>

859 Görden, K., Lynch, A. H., Marshall, A. G., Beringer, J., 2006. Impact of
860 abrupt land cover changes by savanna fire on northern Australian climate.
861 *Journal of Geophysical Research-Atmospheres* 111 (D19), D19106.
862 URL <http://dx.doi.org/10.1029/2005JD006860>

- 863 Hastie, T., Tibshirani, R., 1986. Generalized additive models. Statistical sci-
864 ence, 297–310.
- 865 Hipel, K., McLeod, A., 1994. Time series modelling of water resources and
866 environmental systems. Vol. 45. Elsevier Science Ltd.
- 867 Hirsch, R. M., Gilroy, E. J., 1985. Detectability of step trends in the rate of
868 atmospheric deposition of sulfate. JAWRA Journal of the American Water
869 Resources Association 21 (5), 773–784.
870 URL <http://dx.doi.org/10.1111/j.1752-1688.1985.tb00171.x>
- 871 Holper, P. N., 2011. Climate change, science information paper: Australian
872 rainfall-past, present and future. Canberra: CSIRO.
- 873 Howden, S., 2012. It’s official: Australia no longer in
874 drought. Brisbane Times 27 April, 2012. Available online at
875 [http://www.brisbanetimes.com.au/environment/weather/](http://www.brisbanetimes.com.au/environment/weather/its-official-australia-no-longer-in-drought-20120427-1xpsp.html)
876 [its-official-australia-no-longer-in-drought-20120427-1xpsp.](http://www.brisbanetimes.com.au/environment/weather/its-official-australia-no-longer-in-drought-20120427-1xpsp.html)
877 [html](http://www.brisbanetimes.com.au/environment/weather/its-official-australia-no-longer-in-drought-20120427-1xpsp.html), viewed on 23/08/2013.
- 878 Hughes, L., 2003. Climate change and Australia: Trends, projections and
879 impacts. Austral Ecology 28 (4), 423–443.
880 URL <http://dx.doi.org/10.1046/j.1442-9993.2003.01300.x>
- 881 Jones, D., Wang, W., Fawcett, R., 2009. High-quality spatial climate data-
882 sets for australia. Australian Meteorological and Oceanographic Journal
883 58 (4), 233–248.
- 884 Junkermann, W., Hacker, J., Lyons, T., Nair, U., 2009. Land use change

885 suppresses precipitation. *Atmospheric Chemistry and Physics* 9 (17), 6531–
886 6539.
887 URL <GotoISI>://WOS:000269778500018

888 Kamruzzaman, M., Beecham, S., Metcalfe, A. V., 2011. Non-stationarity
889 in rainfall and temperature in the Murray Darling Basin. *Hydrological*
890 *Processes* 25 (10), 1659–1675.
891 URL <http://dx.doi.org/10.1002/hyp.7928>

892 Karoly, D. J., Aug. 2014. Climate change: Human-induced rainfall changes.
893 *Nature Geoscience*; London 7 (8), 551–552.
894 URL [https://search.proquest.com/docview/1656589632/citation/](https://search.proquest.com/docview/1656589632/citation/8DB40BDF8E794FC4PQ/1)
895 [8DB40BDF8E794FC4PQ/1](https://search.proquest.com/docview/1656589632/citation/8DB40BDF8E794FC4PQ/1)

896 Kucharski, F., Zeng, N., Kalnay, E., 2013. A further assessment of vegetation
897 feedback on decadal Sahel rainfall variability. *Climate dynamics* 40 (5-6),
898 1453–1466.
899 URL [http://link.springer.com/article/10.1007/](http://link.springer.com/article/10.1007/s00382-012-1397-x)
900 [s00382-012-1397-x](http://link.springer.com/article/10.1007/s00382-012-1397-x)

901 Kundzewicz, Z., Robson, A., 2004. Change detection in hydrological records-
902 a review of the methodology/revue méthodologique de la détection de
903 changements dans les chroniques hydrologiques. *Hydrological sciences jour-*
904 *nal* 49 (1).

905 Lymburner, L., Tan, P., Mueller, N., Thackway, R., Lewis, A., Thankap-
906 pan, M., Randall, L., Islam, A., Senarath, U., 2010. 250 metre Dynamic

907 Land Cover Dataset. Geoscience Australia, Canberra, digital data access
 908 on 28/12/2011 from www.ga.gov.au/earth-observation/landcover.

909 Ma, H. Y., Mechoso, C. R., Xue, Y., Xiao, H., Wu, C. M., Li, J. L., De Sales,
 910 F., Jul 2011. Impact of land surface processes on the South American warm
 911 season climate. *Climate Dynamics* 37 (1-2), 187–203.
 912 URL <GotoISI>://WOS:000293403000012

913 MacDonald, G. M., Case, R. A., 2005. Variations in the pacific decadal os-
 914 cillation over the past millennium. *Geophysical Research Letters* 32 (8),
 915 L08703.
 916 URL <http://dx.doi.org/10.1029/2005GL022478>

917 Maynard, K., Polcher, J., Apr 2003. Impact of land-surface processes on
 918 the interannual variability of tropical climate in the LMD GCM. *Climate*
 919 *Dynamics* 20 (6), 613–633.
 920 URL <GotoISI>://WOS:000183022400004

921 McAlpine, C. A., Syktus, J., Deo, R. C., Lawrence, P. J., McGowan, H. A.,
 922 Watterson, I. G., Phinn, S. R., Nov 2007. Modeling the impact of historical
 923 land cover change on Australia’s regional climate. *Geophysical Research*
 924 *Letters* 34 (22).
 925 URL <GotoISI>://000251345100005

926 Mei, R., Wang, G. L., Jun 2010. Rain follows logging in the Amazon? Results
 927 from CAM3-CLM3. *Climate Dynamics* 34 (7-8), 983–996.
 928 URL <GotoISI>://WOS:000278088400005

- 929 Meneghini, B., Simmonds, I., Smith, I. N., 2007. Association between Aus-
 930 tralian rainfall and the Southern Annular Mode. *International Journal of*
 931 *Climatology* 27 (1), 109–121.
 932 URL <http://dx.doi.org/10.1002/joc.1370>
- 933 Murphy, B. F., Timbal, B., 2008. A review of recent climate variability and
 934 climate change in southeastern Australia. *International Journal of Clima-*
 935 *tology* 28 (7), 859–879.
 936 URL <http://dx.doi.org/10.1002/joc.1627>
- 937 Nair, U. S., Wu, Y., Kala, J., Lyons, T. J., Pielke, R. A., S., Hacker, J. M.,
 938 2011. The role of land use change on the development and evolution of the
 939 west coast trough, convective clouds, and precipitation in southwest Aus-
 940 tralia. *Journal of Geophysical Research-Atmospheres* 116 (D7), D07103.
 941 URL <http://dx.doi.org/10.1029/2010JD014950>
- 942 Narisma, G., Foley, J., Licker, R., Ramankutty, N., 2007. Abrupt changes in
 943 rainfall during the twentieth century. *Geophysical Research Letters* 34 (6),
 944 L06710.
- 945 Narisma, G. T., Pitman, A. J., 2003. The impact of 200 years of land cover
 946 change on the Australian near-surface climate. *Journal of Hydrometeorol-*
 947 *ogy* 4 (2), 424–436.
- 948 Negri, A. J., Adler, R. F., Xu, L. M., Surratt, J., Mar 2004. The impact of
 949 Amazonian deforestation on dry season rainfall. *Journal of Climate* 17 (6),
 950 1306–1319.
 951 URL [http://journals.ametsoc.](http://journals.ametsoc.org/doi/10.1175/JCLI122003)

952 org/doi/pdf/10.1175/1520-0442%282004%29017%3C1306%3ATI0AD0%
 953 3E2.0.CO%3B2

954 Nicholls, N., 2004/04/01 2004. The changing nature of australian droughts.
 955 Climatic Change 63 (3), 323–336.
 956 URL <http://dx.doi.org/10.1023/B%3ACLIM.0000018515.46344.6d>

957 Nicholls, N., 2006. Detecting and attributing Australian climate change: A
 958 review. Australian Meteorological Magazine 55 (3), 199–211.

959 Oleson, K. W., Bonan, G. B., Levis, S., Vertenstein, M., 2004. Effects of
 960 land use change on North American climate: impact of surface datasets
 961 and model biogeophysics. Climate Dynamics 23 (2), 117–132.

962 Onoz, B., Bayazit, M., 2003. The power of statistical tests for trend detection.
 963 Turkish Journal of Engineering and Environmental Sciences 27 (4), 247–
 964 251.

965 Otterman, J., Manes, A., Rubin, S., Alpert, P., Starr, D., 1990. An increase of
 966 early rains in Southern Israel following land-use change? Boundary-Layer
 967 Meteorology 53 (4), 333–351.

968 Pinto, E., Shin, Y., Cowling, S. A., Jones, C. D., May 2009. Past, present and
 969 future vegetation-cloud feedbacks in the Amazon Basin. Climate Dynamics
 970 32 (6), 741–751.
 971 URL <GotoISI>://WOS:000264118400001

972 Pitman, A. J., Hesse, P. P., 2007. The significance of large-scale land cover
 973 change on the Australian palaeomonsoon. Quaternary Science Reviews

974 26 (1-2), 189–200.

975 URL [http://www.sciencedirect.com/science/article/](http://www.sciencedirect.com/science/article/B6VBC-4MC71SN-1/2/0e1d4dff6e15a1de79cefbfa612f9b37)

976 [B6VBC-4MC71SN-1/2/0e1d4dff6e15a1de79cefbfa612f9b37](http://www.sciencedirect.com/science/article/B6VBC-4MC71SN-1/2/0e1d4dff6e15a1de79cefbfa612f9b37)

977 Pitman, A. J., Lorenz, R., 2016. Scale dependence of the simulated impact

978 of Amazonian deforestation on regional climate. *Environmental Research*

979 *Letters* 11 (9), 094025.

980 URL <http://stacks.iop.org/1748-9326/11/i=9/a=094025>

981 Pitman, A. J., Narisma, G. T., Pielke, R. A., Holbrook, N. J., 2004. Impact of

982 land cover change on the climate of southwest Western Australia. *Journal*

983 *of Geophysical Research-Atmospheres* 109 (D18).

984 URL <GotoISI>://WOS:000224126500001

985 R Development Core Team., 2011. R: A Language and Environment for Sta-

986 tistical Computing. R Foundation for Statistical Computing, Vienna, Aus-

987 tria, ISBN 3-900051-07-0.

988 URL <http://www.R-project.org/>

989 Risbey, J. S., Pook, M. J., McIntosh, P. C., Wheeler, M. C., Hendon, H. H.,

990 2009/10/01 2009. On the remote drivers of rainfall variability in Australia.

991 *Monthly Weather Review* 137 (10), 3233–3253.

992 URL <http://dx.doi.org/10.1175/2009MWR2861.1>

993 Roy, D., Jin, Y., Lewis, P., Justice, C., 2005. Prototyping a global algorithm

994 for systematic fire-affected area mapping using modis time series data.

995 *Remote sensing of environment* 97 (2), 137–162.

- 996 Roy, D., Lewis, P., Justice, C., 2002. Burned area mapping using multi-
 997 temporal moderate spatial resolution data—a bi-directional reflectance
 998 model-based expectation approach. *Remote Sensing of Environment* 83 (1),
 999 263–286.
- 1000 Roy, D. P., Boschetti, L., Justice, C. O., Ju, J., 2008. The collection 5 modis
 1001 burned area product—global evaluation by comparison with the modis
 1002 active fire product. *Remote Sensing of Environment* 112 (9), 3690–3707.
- 1003 Saha, S. K., Dirmeyer, P. A., Chase, T. N., 2016. Investigating the impact of
 1004 land-use land-cover change on Indian summer monsoon daily rainfall and
 1005 temperature during 1951-2005 using a regional climate model. *Hydrology
 1006 and Earth System Sciences* 20 (5), 1765.
- 1007 URL [http://search.proquest.com/openview/
 1008 c4242c9ecce075e3545390b1d00f2751/1?pq-origsite=gscholar&
 1009 cb1=105724](http://search.proquest.com/openview/c4242c9ecce075e3545390b1d00f2751/1?pq-origsite=gscholar&cb1=105724)
- 1010 Saji, N., Goswami, B., Vinayachandran, P., Yamagata, T., 1999. A dipole
 1011 mode in the tropical Indian Ocean. *Nature* 401 (6751), 360–363.
- 1012 Sato, T., Kimura, F., Hasegawa, A. S., Dec 2007. Vegetation and topographic
 1013 control of cloud activity over arid/semiarid Asia. *Journal of Geophysical
 1014 Research-Atmospheres* 112 (D24).
- 1015 URL [http://www.agu.org/
 1016 journals/jd/jd0724/2006JD008129/2006JD008129.pdf](http://www.agu.org/journals/jd/jd0724/2006JD008129/2006JD008129.pdf)
- 1017 Schepen, A., Wang, Q. J., Robertson, D., 2012 Feb 15 2012. Evidence for

1018 using lagged climate indices to forecast australian seasonal rainfall. Journal
 1019 of Climate 25 (4), 1230–1246.

1020 Semazzi, F. H. M., Song, Y., 2001. A GCM study of climate change induced
 1021 by deforestation in Africa. Climate Research 17 (2), 169–182.

1022 Smith, I. N., Timbal, B., 2012. Links between tropical indices and southern
 1023 Australian rainfall. International Journal of Climatology 32 (1), 33–40.

1024 Speer, M., Leslie, L., Fierro, A., 2011. Australian east coast rainfall decline
 1025 related to large scale climate drivers. Climate Dynamics 36 (7), 1419–1429.
 1026 URL <http://dx.doi.org/10.1007/s00382-009-0726-1>

1027 The State Government of Victoria, 2011. Bushfire his-
 1028 tory. Available online at [http://www.dse.vic.gov.au/](http://www.dse.vic.gov.au/fire-and-other-emergencies/major-bushfires-in-victoria)
 1029 [fire-and-other-emergencies/major-bushfires-in-victoria](http://www.dse.vic.gov.au/fire-and-other-emergencies/major-bushfires-in-victoria), viewed
 1030 on 12/12/2011.

1031 Timbal, B., Arblaster, J., 2006. Land cover change as an additional forcing
 1032 to explain the rainfall decline in the south west of Australia. Geophysical
 1033 Research Letters 33 (7), L07717.

1034 Townshend, J., Carroll, M., Dimiceli, C., Sohlberg, R., Hansen, M.,
 1035 DeFries, R., 2011. Vegetation continuous fields MOD44B, 2001
 1036 Percent tree cover, Collection 5. University of Maryland, Col-
 1037 lege Park, Maryland, 2001, digital data accessed on 10/02/2012
 1038 from [https://lpdaac.usgs.gov/products/modis_products_table/](https://lpdaac.usgs.gov/products/modis_products_table/vegetation_continuous_fields/yearly_13_global_500m/mod44b)
 1039 [vegetation_continuous_fields/yearly_13_global_500m/mod44b](https://lpdaac.usgs.gov/products/modis_products_table/vegetation_continuous_fields/yearly_13_global_500m/mod44b).

1040 Trenberth, K. E., May 1999. Atmospheric moisture recycling: Role of advec-
 1041 tion and local evaporation. *Journal of Climate* 12 (5), 1368–1381.
 1042 URL <GotoISI>://000080145700002

1043 Verdon, D. C., Wyatt, A. M., Kiem, A. S., Franks, S. W., 2004. Multidecadal
 1044 variability of rainfall and streamflow: Eastern Australia. *Water Resources*
 1045 *Research* 40 (10), W10201.
 1046 URL <http://dx.doi.org/10.1029/2004WR003234>

1047 Wang, J., Chagnon, F., Williams, E., Betts, A., Renno, N., Machado,
 1048 L., Bisht, G., Knox, R., Bras, R., 2009. Impact of deforestation in the
 1049 Amazon basin on cloud climatology. *Proceedings of the National Academy*
 1050 *of Sciences of the United States of America* 106 (10), 3670.
 1051 URL [http://ezproxy.library.usyd.edu.au/login?url=http:](http://ezproxy.library.usyd.edu.au/login?url=http://search.proquest.com/docview/201417978?accountid=14757)
 1052 [//search.proquest.com/docview/201417978?accountid=14757](http://search.proquest.com/docview/201417978?accountid=14757)

1053 Westra, S., Alexander, L. V., Zwiers, F. W., 2013/06/01 2013. Global in-
 1054 creasing trends in annual maximum daily precipitation. *Journal of Climate*
 1055 26 (11), 3904–3918.
 1056 URL <http://dx.doi.org/10.1175/JCLI-D-12-00502.1>

1057 Westra, S., Sharma, A., 2010. An upper limit to seasonal rainfall predictabil-
 1058 ity? *Journal of Climate* 23 (12), 3332–3351.
 1059 URL [http://ezproxy.library.usyd.edu.au/login?url=http:](http://ezproxy.library.usyd.edu.au/login?url=http://search.proquest.com/docview/814127501?accountid=14757http://DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+Predictability%3F&title=Journal+of+Climate&issn=0894-8755&)
 1060 [//search.proquest.com/docview/814127501?accountid=14757http:](http://search.proquest.com/docview/814127501?accountid=14757http://DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+Predictability%3F&title=Journal+of+Climate&issn=0894-8755&)
 1061 [//DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=](http://DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+Predictability%3F&title=Journal+of+Climate&issn=0894-8755&)
 1062 [article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+](http://DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+Predictability%3F&title=Journal+of+Climate&issn=0894-8755&)
 1063 [Predictability%3F&title=Journal+of+Climate&issn=0894-8755&](http://DD8GH5YX7K.search.serialssolutions.com/?SS_Source=3&genre=article&sid=ProQ:&atitle=An+Upper+Limit+to+Seasonal+Rainfall+Predictability%3F&title=Journal+of+Climate&issn=0894-8755&)

1064 date=2010-06-01&volume=23&issue=12&spage=3332&SS_docid=
 1065 814127501&author=Westra%2C+Seth%3BSharma%2C+Ashish
 1066 Whitley, E., Ball, J., 2002. Statistics review 6: Nonparametric methods.
 1067 Critical Care, London 6 (6), 509–513.
 1068 Wilks, D., 1997. Resampling hypothesis tests for autocorrelated fields. Jour-
 1069 nal of Climate 10 (1), 65–82.
 1070 Wilks, D. S., Sep 2006 2006. On "field significance" and the false discovery
 1071 rate. Journal of Applied Meteorology and Climatology 45 (9), 1181–1189.
 1072 URL <http://journals.ametsoc.org/doi/abs/10.1175/JAM2404.1>
 1073 Wood, S. N., 2011. Fast stable restricted maximum likelihood and marginal
 1074 likelihood estimation of semiparametric generalized linear models. Journal
 1075 of the Royal Statistical Society (B) 73 (1), 3–36.
 1076 Zanchettin, D., Franks, S. W., Traverso, P., Tomasino, M., 2008. On enso
 1077 impacts on european wintertime rainfalls and their modulation by the nao
 1078 and the pacific multi-decadal variability described through the pdo index.
 1079 International Journal of Climatology 28 (8), 995–1006.
 1080 URL <http://dx.doi.org/10.1002/joc.1601>
 1081 Zeng, X. M., Wu, Z. H., Song, S., Xiong, S. Y., Zheng, Y. Q., Zhou, Z. G.,
 1082 Liu, H. Q., Jan 2012. Effects of land surface schemes on the simulation
 1083 of a heavy rainfall event by WRF. Chinese Journal of Geophysics-Chinese
 1084 Edition 55 (1), 16–28.
 1085 URL <GotoISI>://WOS:000300129300002