

Detecting the impact of land cover change on observed rainfall.

Chun X. Liang¹, Floris F. van Ogtrop¹, and R. Willem Vervoort¹

¹Sydney Institute of Agriculture, The University of Sydney, NSW 2006

Corresponding author:

R. Willem Vervoort¹

Email address: willem.vervoort@sydney.edu.au

ABSTRACT

Analysis of observational data to identify relationships between rainfall and land cover change are difficult due to multiple environmental factors that cannot be strictly controlled. In this study we present a methodology to investigate the relationship using statistical methods on data from best available sources at two sites in Australia. Gridded data of rainfall and tree cover were used as spatially corresponding local conditions. Large scale effects were represented by climatic indicators, such as SOI and IOD. Regression analysis and step trend tests were used to assess the effect of abrupt land surface intervention. At a Queensland site, significant tree cover change between 2002 - 2005 did not result in strong statistically significant precipitation changes. On the other hand, results from a bushfire affected NSW/VIC region suggests significant changes in the rainfall. This indicates the method works better when a abrupt change in the data can be clearly identified. The results from the step trend test implied a positive relationship between the tree cover and the rainfall at 0.1 significance level in both locations in data up to 2009. However, high rainfall variability and possible regrowth meant that no significant changes were observed in longer time series to 2015.

Warning: package 'kableExtra' was built under R version 3.5.2

INTRODUCTION

Land use and land cover changes can lead to changes in the local climate. Empirical and modelling studies have found that cloud types and rainfall are correlated to large scale vegetation cover changes, such as deforestation in the Amazon and in the Sahel (Chagnon and Bras 2005; Pinto et al. 2009; Wang et al. 2009; Mei and Wang 2010; Kucharski, Zeng, and Kalnay 2013; Pitman and Lorenz 2016) and afforestation in south Israel (Otterman et al. 1990; Ben-Gai et al. 1998). Using airborne measurement in Western Australia, Junkermann et al. (2009) showed a significantly higher level of aerosols over an agricultural area compared to an adjacent natural vegetation. They suggested that a modification of aerosol concentrations due to deforestation could have contributed to a reduction of local rainfall, as more, but smaller rain droplets were observed. Nair et al. (2011) reported from the Bunny Fence Experiment in Western Australia that local land use change altered the synoptic west coast trough dynamics and surface roughness, and this resulted in an observed rainfall decrease. Maximum temperatures were also found to be sensitive to land cover change in eastern Australia (McAlpine et al. 2007).

Overall the number of empirical studies analyzing changes to rainfall due to land cover change from observational data is limited. Most of the studies mentioned previously were either model simulations, or comparisons of modelled data with observations. This is because there are some fundamental experimental difficulties in both space (where does evaporated water reappear as rainfall?) and in time (how much time does it take for land cover change effects to appear or disappear?). In addition, in many areas across the globe, rainfall variability is related to a complex set of interactions, of which land use change might only be a minor component.

Locally, there are two main sources that generate rainfall: moisture from advective atmospheric transport; and local evapotranspiration (Eltahir and Bras 1996; Bosilovich and Chern 2006; Dirmeyer, Brubaker, and DelSole 2009; Gimeno et al. 2010). The local evapotranspiration component is the

component considered to be affected by land use change (Eltahir and Bras 1996). According to Trenberth (1999), the contribution of advective moisture partially depends on the availability of external moisture and atmospheric transport. On the longer time scale, such as monthly and annually, large scale atmospheric dynamics are affected by large scale climate drivers. For example, many studies have reported significant relationships between rainfall in large parts of Australia and the El Niño-Southern Oscillation (ENSO) (Verdon et al. 2004; Risbey et al. 2009; Speer, Leslie, and Fierro 2011). In contrast, local ET is determined by local land surface characteristics, which influence local scale atmospheric dynamics and hence the amount of rainfall, including contribution from both main sources.

Although climate drivers demonstrate some capability to predict Australian rainfall, there is still a large amount of unexplained variance. Westra and Sharma (2010) pointed out that models based on global sea surface temperature anomalies can only predict up to 14.7% of annual precipitation variance. Some of the remaining variance could be due to land surface processes as suggested in studies predicting local rainfall (e.g. Ma et al. 2011; Zeng et al. 2012; Pitman and Lorenz 2016; Saha, Dirmeyer, and Chase 2016). However, most are based on modelling experiments and few empirical observational studies have been reported. However, Pitman et al. (2004) found a good match between observations and simulated rainfall changes in southwest Western Australia, forced by land cover change. Timbal and Arblaster (2006) were able to reproduce the rainfall decline in south west Australia by including land cover influence. In addition, local land use change might not be a primary, but is likely to be a secondary cause of rainfall change (Nicholls 2006).

Therefore, the aim of this study is to use a statistical approach on rainfall data at regional scales to investigate the cause and effect relationship between land cover change and local rainfall, which is demonstrated in many modelling studies. More specifically, we hypothesize that a step change on the land cover on the surface will cause a step change in the rainfall. To demonstrate this we study changes in observed rainfall over time at a Queensland and NSW/Victoria location where there are possible step changes in land cover change due to land clearing and bush fires. The methodology uses statistical approaches to identify changes in rainfall, which are subsequently associated with land cover change through spatial comparison.

In this paper, after this section (the introduction), section 2 covers the case study areas and the observed land use change. Section 3 describes the data used in the study in more detail. Section 4 details the statistical methods and the underlying assumptions related to the modelling approach, Section 5 gives the results, which are further discussed in section 6 and finally section 7 offers the conclusions.

1 STUDY REGIONS AND TREE COVER CHANGE

In Australia, significant tree cover change has mainly occurred in the north east and south east of the continent, as well as in the southwest of Western Australia. According to the National Dynamic Land Cover Dataset (DLCD) (Lymburner et al. 2010), most of these areas have experienced decreases in the Enhanced Vegetation Index (EVI) post 2000 derived from satellite data. As an index for vegetation greenness, the decreasing values indicate lower biomass over time in the tree cover regions. The possible EVI reduction might be due to land clearing, bush fires or drought.

Two regions were selected where significant tree cover change since 2000 was reported. The first region is located in south central Queensland partly covering the north of the Murray Darling Basin (MDB) (site 1 in Figure 1). High rates of land clearing have been reported in this region during the early 2000s (Department of Natural Resources and Water 2007). The second study region is located at the border of New South Wales and Victoria, and includes the Snowy Mountain ranges (site 2 in Figure 1). Severe bush fires occurred in this area in early 2003 (see Figure 2). The 2003 bush fires were the largest and the worst in the last 60 years (The State Government of Victoria 2011). Two thirds of Kosciuszko national park was heavily burned and regrowth was reported to be slow due to drought and cold conditions (ABC News 2003), and the type of species in this region. However, in the longer term, after an early high transpiration period a recovery of pre-fire evapotranspiration would be expected (Kuczera 1987). For the purpose of this study, significant tree cover loss has happened in both study areas in the last decade, either permanently or temporarily.

The two regions have different characteristics. The QLD region is partially grassland and subtropical, while the NSW/VIC region is mainly within the temperate zone, under the Köppen classification. According to Australian Bureau of Meteorology (BoM), the NSW/VIC region receives 1000 - 2000 mm rainfall annually, which is more than double the annual rainfall in the QLD region. Evapotranspiration is similar




Figure 1. Selected study regions are highlighted by red rectangles in the main map (the red rectangle in the insert indicates the location of the main map). The types of tree cover in 2008 from the DLCD product is shown at the background. In site 1 (the QLD region), the tree cover is mostly sparse. In site 2 (the NSW/VIC region), many areas have open or close forest where the tree cover is denser.

in both regions. Marine moisture and orographic effects are likely to be the main contributors to rainfall in the southeast mountain areas of the NSW/VIC region.

The land use and land cover characteristics in the two regions are also different. In the Queensland region, the tree cover is sparse over most of the area. The MODIS satellite tree cover data (discussed in more detail in section 3) shows that tree cover in this region is generally below 20% of total ground area. Grazing is the main activity in this region, with over 90% of land used by the grazing industry (ABARES 2010). Our starting assumption is that the main cause of the EVI decline over large part of the region is due to land clearing. Tree cover has been cleared at a massive scale over the last decade, especially during 2002 - 2004. The reports from the Queensland Statewide Land Cover and Trees Study (SLATS) (e.g. Department of Natural Resources and Mines 2005; Department of Science, Information Technology and Innovation 2017) were used to investigate the time and location of the land clearing in the QLD region.

The Kosciuszko national park is within the NSW/VIC region. Here tree cover is denser with open or even closed forest (the tree cover distribution is bimodal at 10 - 20% and 60 - 70%). The dominant species in the alpine area are Snow Gum and large stand species such as Alpine Ash and Mountain Gum in the sub-alpine area. These trees can reach a great height but they take long time to grow. For example, Alpine Ash would need about 20 years to mature. Although land clearing is not a major issue in this region, it is vulnerable to fires and drought. The MODIS burned area product, MCD45A1 (Roy, Lewis, and Justice 2002; Roy et al. 2005; Roy et al. 2008), was used to locate bush fires areas in the NSW/VIC region, with a grid resolution of 500 m. MCD45A1 provides monthly burning information on all pixels, which helps to pinpoint an abrupt event.

Due to the difference in nature of the land cover change in the two regions, the post-change vegetation status is hypothesized to be different as well (see Figure 3).

The overall hypothesis is that the effect of 2003 - 2004 land clearings in the QLD region and the 2003 bush fires in the NSW/VIC region cause a step change in the local rainfall. The actual tree cover change during this time at the pixel level was derived from the 15-year MODIS data (discussed below). The difference of tree cover before and after the land disturbance was tested using a Student's t-test. As the length of the tree cover data is shorter than available rainfall data, earlier land clearing in the QLD region




Figure 2. Location of bushfires occurring in January 2003, in and around the NSW/VIC study region, as shown by the red pixels. The map shows the large area in the Kosciuszko national park that has been burned. Some locations in the southwest of the Australian Capital Territory (ACT) have also experienced intensive bushfires.




Figure 3. The expected evolution of the land surface after trees have been removed in (a) the QLD region and (b) the NSW/VIC region.

cannot be identified spatially, hence they are excluded from the analysis.

2 DATA

Several land surface data sets were used in this study. The main one was the MOD44B product Global Vegetation Continuous Field data set (version 5). This data set provides estimates of percent tree cover (percentage of ground surface covered by trees) at a grid resolution of 250 m (Townshend et al. 2011), which is finer than the earlier mentioned burned product MCD45A1. The data set is available on an annual time interval for the study period of 2000 - 2015. The tree cover data was produced from 16-day Terra MODIS Land Surface Reflectance data and Land Surface Temperature (Townshend et al. 2011). The National Dynamic Land Cover Dataset (DLCD) (Lymburner et al. 2010) from the Australian Collaborative Land Use Mapping Program (ACLUMP) was used to verify the trend of vegetation cover change calculated from the previous data set. This data set, developed by Geoscience Australia and Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES), is the first nationally consistent and thematically comprehensive land cover reference for Australia. The DLCD is based on the 16-day Enhanced Vegetation Index (EVI), again from the MODIS satellite, between April 2000 and April 2015. It also has a grid resolution of 250 m. The data set provides information on the final land cover types (as in 2015) and estimated trend of EVI statistics (annual mean, maximum and minimum).

The SILO Rainfall product data for Australia was used (Jeffrey et al. 2001) (available online at <https://silo.longpaddock.qld.gov.au/>). The data has been projected onto a national 0.05° by 0.05° grid (approximately 5 km by 5 km). This gridded data set was generated from station observations using spline interpolation and kriging (Jeffrey et al. 2001). The data has been compared to other gridded products and observed data and is generally of high quality (Tozer, Kiem, and Verdon-Kidd 2009; Tozer, Kiem, and Verdon-Kidd 2012). The data is available on a daily and monthly basis from 1889 to current. Here a subset of 36 years (1979 - 2015) was used. The study was conducted on monthly data, as a land cover change effect on annual rainfall might be negligible, but can often be significant in particular months or seasons (e.g. Otterman et al. 1990; Gaertner et al. 2001; Semazzi and Song 2001; Oleson et al. 2004; Deo et al. 2009).

Large scale climate drivers are represented by various climatic indices. The Southern Oscillation Index (SOI) is generally regarded as a good predictor of Australian rainfall (Risbey et al. 2009; Chowdhury and Beecham 2010; Westra and Sharma 2010), but its skill is weaker in some parts of Australia. For example

the Southern Annular Mode (SAM) is found to be more important than ENSO in south Western Australia (Meneghini, Simmonds, and Smith 2007). The testing of the suitability of each index for the regions of interest is described in a later section. The following climate indices were used as candidate predictors for local rainfall.

- Southern Oscillation Index (SOI). The Troup version of the monthly SOI series used in this study was obtained from BoM (available online at <http://www.bom.gov.au/climate/current/soihtml.shtml>).
- Eastern, East Central and Central Tropical Pacific Sea Surface Temperatures (NINO 3, NINO 3.4 and NINO 4). Monthly SST anomalies are available from IRI/LDEO data library and the extended NINO data set is used (available online at <http://iridl.ldeo.columbia.edu/SOURCES/.Indices/.nino/.EXTENDED/>).
- Pacific Decadal Oscillation (PDO). The Pacific Decadal Oscillation is the leading principal component of monthly SST anomaly in the North Pacific Ocean. The monthly PDO series was provided by JISAO (Joint Institute for the Study of the Atmosphere and Ocean, University of Washington) (available online at <http://jisao.washington.edu/pdo/PDO.latest>).
- Indian Ocean Dipole (IOD). The Indian Ocean dipole is commonly measured by the difference between SST anomaly in the western (50 - 70°E and 10°S-10°N) and eastern (90 - 110°E and 0 - 10°S) equatorial India Ocean (Saji et al. 1999). Monthly IOD was obtained from JAMSTEC (the Japan Agency for Marine-Earth Science and Technology) (available online at <http://www.jamstec.go.jp/frcgc/research/dl/iod/DATA/dmi.monthly.txt>).

3 STATISTICAL METHOD

As an initial analysis a simple boxplot and t-test is used to analyse whether there is a significant change in tree cover in time, and before and after the suspected change in the regions.

To assess the actual causal relationship between the tree cover and the rainfall a flexible regression model is applied (discussed in detail below). A step change is not directly obvious in the time series of the rainfall anomalies (Figure 4) for both regions, even though the data is deseasonalised and detrended. In this study we apply different statistical methods to analyse the effect of tree cover change on rainfall. Both methods make use of a regression model to remove year-on-year variability in rainfall to strengthen the tree cover change signal.

In the first method, the tree cover change is implemented as a factor variable in the regression model, and the significance of this variable is tested. In the second method, a rank sum test (step trend test), is applied to the regression model residuals after effects of other major factors were removed. This assumes that after removal of all climate, long term linear trend and seasonal variation, the vegetation cover change is the only factor explaining the non-random pattern in the rainfall residuals.

3.0.1 Regression model

As highlighted in the introduction, the Australian climate is influenced by sea surface temperatures in the tropical Pacific and Indian Oceans, as well as pressure systems in the Southern Ocean (BoM 2012b). Risbey et al. (2009) compared five large-scale drivers, including ENSO (measured by SOI and the Tropical Pacific Sea surface temperatures (SSTs)), IOD, SAM, MJO (Madden-Julian oscillation) and blocking, in relation to Australian rainfall variability. The MJO is a large scale eastward-propagating wave-like disturbance located around equatorial latitudes (Risbey et al. 2009). They identified SOI as the most important index among all climate indices tested for broad parts of Australia (including QLD and NSW/VIC) in almost any season. In this study, four climate indices were selected from the main climatic indicators (see section Data) and used as the explanatory variables in the model for each study region. A further complicating factor is the influence of the “millenium drought” over the study period and in particular the change to wet conditions in 2010 - 2011 (Dijk and Viney 2013). Therefore, the spatially averaged monthly rainfall in the Murray Darling Basin (MDB, downloaded from the Bureau of Meteorology) was used to explain the year-on-year variation in the rainfall in the regions. As climate alone are unlikely able to fully explain the temporal variation in rainfall in the regions, including the




Figure 4. The deseasonalised and detrended rainfall over the 30 years period in (a) the QLD region and (b) the NSW/VIC region. The vertical red lines indicate the year of 2003, in which the studied land cover changes occurred. A change in the time series data is not obvious before and after the land cover changes.




Figure 5. Cross-correlation of six climate indices and rainfall in QLD study region. For the PDO analysis, 108-year rainfall data (1900 - 2008) are used. Otherwise, 36-year rainfall data are used. The correlation with NINO 3 is not shown as it is very similar to but weaker than for NINO 3.4.

drought (Dijk and Viney 2013; Westra and Sharma 2010), this variable was included. Since both regions at least partly overlap with the MDB, the average rainfall across was assumed to be a useful explaining variable.

Correlations between rainfall and each climate index were analysed. Rainfall in each study region was first deseasonalised and detrended using the seasonal decomposition function `ds` in the package `deseasonalise` in R (R Core Team, 2018). Using detrended data gives a better indication of the underlying correlation compared to the correlation between trends in the data (Smith and Timbal 2012). The cross-correlations between the deseasonalised and detrended rainfall and the climatic indices were tested using the Pearson's product moment correlation method, assuming the relationships are linear. Although the optimal technique for exploring the correlation with each index could be different as described in Risbey et al. (2009), the Pearson's method was applied to all indices for consistency. Because the PDO describes the multi-decadal SST with lower frequency (MacDonald and Case 2005; Zanchettin et al. 2008; Kamruzzaman, Beecham, and Metcalfe 2011), instead of 37-year rainfall data, a longer period (115 years, from 1900 to 2015) was used to estimate the correlation with PDO, up to lag 24. For the other indices, the 37-year data was used.

Based on the correlation between the climatic indices and rainfall in the regions (as shown in Figure 5 and Figure 6), it can be concluded that:

- In QLD, the correlation between rainfall and SOI at zero time lags is the strongest across all indices,




Figure 6. Cross-correlation of six climate indices and rainfall in NSW/VIC study region.

outweighing the other ENSO indicators. IOD and PDO have a weak influence in QLD.

- In NSW/VIC, again the SOI has the strongest correlation with rainfall, followed by the IOD. For both, the strongest correlations occur at the zero time lags.
- in both cases PDO had the weakest correlations, and this factor was not further considered as a predictor.
- in general, for the better correlated indices, strongest correlations occurred at zero time lags.

The above findings are consistent with previous studies. Although some indices are serially correlated with rainfall up to several months, the lag zero events have the greatest correlation coefficients. Using multiple climatic index series was generally found most useful in rainfall prediction (e.g. Risbey et al. 2009; Kamruzzaman, Beecham, and Metcalfe 2011).

Rainfall in Australia shows strong seasonal patterns (Holper 2011; Australian Bureau of Statistics 2012). For example, the north part of the country is summer rainfall dominant with a dry winter, while most of the southern part has a winter rainfall regime. This character is driven by the movement of subtropical high pressure systems which dominate the Australian climate (BoM 2012a). These characteristic summer and winter rainfall patterns mean that the seasonal component of rainfall has a periodic pattern which should be included in the model.

Long term trends in the regional rainfall in some parts of Australia are significant (Hughes 2003; Gallant, Hennessy, and Risbey 2007; Chowdhury and Beecham 2010). In the northern and eastern parts of the continent, increasing rainfall is reported over the last century (Hughes 2003). The presence of such long term trends may be confused with the outcome of a step change in rainfall. As a result a linear trend term was implemented in the model to remove any long term effect.

We assumed all the factors are additive smooth components in determining rainfall following Kamruzzaman, Beecham, and Metcalfe (2011). In this case, the rainfall model is a generalised additive model (GAM) (Hastie and Tibshirani 1986) with a log link function $g(\cdot)$ and assuming the residuals are gamma distributed (see Figure 7). This means all predictors are modelled as smooth functions, in this case using the shrinkage version of the cubic regression splines (Wood 2011). All splines were limited to 3 knots in flexibility (Wood 2011), to reduce the risk of overfitting.

$$g(E(\mathbf{R}_r)) = \beta_0 + s_1(\mathbf{MDB}_{\text{monthlyRain}} + s_2(\mathbf{SOI}) + s_3(\mathbf{IOD}) + s_4(\mathbf{Nino3.4}) + s_5(\mathbf{Nino4}) + s_6(\mathbf{Season}) + \beta_1 \mathbf{Trend} + \varepsilon_r \quad (1)$$

The bold letters represent the time series vectors. The region is indicated by r , while β_u ($u=0, 1$) are the fitted coefficients in the model. s_v ($v=1, 2, 3, \dots$) are the smooth penalized cubic regression spline functions on the climatic indices and the season. $\mathbf{MDB}_{\{\text{monthlyRain}\}}$ is the spatially averaged monthly rainfall across the Murray Darling Basin. Apart from dropping PDO as a predictor, all other climate indices were included, allowing the model to select the appropriate predictors.

A possible linear long term trend in the rainfall data is modelled by $\mathbf{Trend} = 1, 2, 3, \dots, n$, where n is the total number of months in the time series. \mathbf{Season} is the seasonal component. The climatic terms are also modelled with smooth functions. The effect of large scale drivers on Australian rainfall is more likely to be seasonal (Murphy and Timbal 2008; Schepen, Wang, and Robertson 2012), and the smooth spline function is more flexible in reproducing the variability in impacts of the climatic indices.

3.1 Tree cover change as factor variable

One of the main difficulties in empirical observation studies related to the effect of land cover change on rainfall is the lack of continuous monitoring of land surface variables. Moreover, no specific variable can possibly be defined that can clearly represent the land surface process. Given the lack of a full picture of the land surface process, a factor variable was used in the regression model to represent the abrupt land surface change (see Equation (1)). The change could be a result of either land clearing or bush fires as long as it is permanent or takes a long time to recover. As indicated we approached the problem in two different models ways.




Figure 7. Distribution of monthly rainfall in (a) QLD and (b) NSW/VIC. Using a Kolmogorov-Smirnov test with shape = 1 and 2.4 for QLD and NSW/VIC respectively, rainfall in both regions can be modelled as a gamma distribution.

271 In the first method, the tree cover change was used as a predictor in the regression model, represented
 272 by a factor variable **LC**. The significance of the coefficient of **LC**, denoted as β'_5 in Equation (1), can be
 273 determined by a ratio test.

$$LC = \begin{cases} \text{Trees} \\ \text{Removed} \end{cases} \quad (2)$$

274 Therefore in both regions, land cover is “trees” for the period before land cover change and “removed”
 275 for the period after the change. Here we simply assumed that vegetation cover change has occurred on
 276 every pixel. The remaining term ε_r is the amount of rainfall that is attributed to other unspecified factors
 277 and random errors. Hence the regression model becomes

$$g(E(\mathbf{R}_r)) = \beta'_0 + s'_1(\mathbf{MDB}_{\text{monthlyRain}} + s'_2(\mathbf{SOI}) + s'_3(\mathbf{IOD}) + s'_4(\mathbf{Nino3.4}) + s'_5(\mathbf{Nino4}) + s'_6\mathbf{Season} + \beta'_1\mathbf{Trend} + \beta'_2\mathbf{LC} + \varepsilon'_r \quad (3)$$

278 One of the difficulties is to point an exact time to the changes in the vegetation cover in the two regions.
 279 In the QLD region, no exact time can be assigned to the land clearing. According to the SLATS reports,
 280 the most substantial clearing occurred between 2003 - 2004 . However, the information on the change in
 281 type of land cover during the time period is missing. Therefore, four scenarios were initially tested in the
 282 analysis (**see supplementary material**). In the NSW/VIC region, severe bush fires were reported in early
 283 January 2003. Hence the “tree” cover state was up to December 2002 then it was changed to “removed”
 284 state from January 2003. As a starting date, the regression model was run from 1979 for both regions.

285 3.2 Step trend test

286 To support the regression analysis, a Mann-Whitney Rank-Sum step trend test was used to detect changes
 287 in rainfall as a result of vegetation cover change. This specific nonparametric statistical test was modified
 288 from the Mann-Whitney U test by Hirsch and Gilroy (1985) and can identify a step change in data which
 289 is cross-correlated. The gridded rainfall dataset used in this study has a high spatial correlation between
 290 neighbouring pixels due to the underlying interpolation method. The advantages of using the Rank-Sum
 291 test are: (1) it does not depend on assumptions of the data distribution; (2) it is not restricted to datasets
 292 with no missing data; (3) it is robust and not as easily influenced by outliers and negative numbers (Hirsch
 293 and Gilroy 1985).

294 The rainfall residuals from the regression model in Equation (3) were used for the step trend test. To
 295 detect a step change, using deseasonalised and detrended data is important (Hirsch and Gilroy 1985).
 296 Furthermore, as rainfall can only be partially attributed to local sources and conditions, other effects are
 297 introduced by large scale dynamics and changes in climatic factors. The assumption is that the regression
 298 model should remove these effects, and additionally deseasonalise and detrend the rainfall data. As
 299 a result, the local landuse effects are amplified in the remaining residuals. The test, described in the
 300 following sub section, subsequently associates trends in the rainfall residuals with tree cover changes.

301 3.2.1 Mann-Whitney rank-sum statistic

302 As indicated, the step trend test is a modified version of the Mann-Whitney U statistic (Hirsch and Gilroy
 303 1985). As a nonparametric rank-based test, the Mann-Whitney test does not use the exact values of rainfall
 304 but depends on the ranks of the data. For each month, rainfall residuals of each year were ranked in an
 305 ascending order. The ranking of January rainfall in a sample pixel k in QLD is illustrated below:

306 Therefore, the smallest or most negative value has rank 1 and the largest value has the maximum rank.

307 The before and after period in the data formed two groups of samples. The split point of the two
 308 periods was based on the timing of the vegetation cover changes. In the QLD region, changes occurred
 309 anytime during 2003 and 2004. In contrast to the previous method, the time period covering the land
 310 cover change was excluded, as the nonparametric test allows missing data. Hirsch and Gilroy (1985) also
 311 pointed out that the power of the test is higher if the data of the change period is ignored. Hence 2003
 312 and 2004 were excluded from the analysis. As a result, the after-change period was 2005 - 2015 for the
 313 Queensland location.

Table 1. Example of ranking rainfall residuals

Year	Rainfall residuals	Rank \$R'_{ijk}\$
1998	-0.3	6
1999	-60.9	2
2000	-16.1	4
2001	-71.7	1
2002	111.1	7
2005	-7.2	5
2006	-60.5	3

In the case of NSW/VIC, the bushfires broke out in early January 2003. The change was within a relatively short period of the year. Therefore the after change period in this region still started in January 2003. Following Hirsch and Gilroy (1985), the before period was set to five years (1998 - 2002) in both regions. The length of the after period is also difficult for the NSW/VIC region, as the regrowth would at some point have impacted the local effects.

The rank of rainfall in month j year i in pixel k is denoted as R'_{ijk} . The sum of ranks of rainfall in month j in pixel k before the known intervention is:

$$W_{jk} = \sum_{i=1}^{n_1} R'_{ijk}. \quad (4)$$

n_1 is the number of years before the land cover change. The expected value of W_{jk} is

$$\mu_w = n_1(n_1 + n_2 + 1)/2 \quad (5)$$

n_2 is the number of years after the change. Hence the expected value of the rank sum before the intervention is the same for all months and all pixels. The sum of ranks for the whole time period is fixed, as $(n_1 + n_2)(n_1 + n_2 + 1)/2$. In this study, since there are only two groups (before and after), knowing the rank-sum of one group is the same as knowing the rank-sum of the other group. If the rainfall data is temporally and spatially independent, the variance of W_{jk} is

$$\sigma_w^2 = n_1 \cdot n_2(n_1 + n_2 + 1)/m \quad (6)$$

where m is the number of months which is 12 in the case of a full year.

3.2.2 Step trend test

Instead of completing the Mann-Whitney U-test, Hirsch and Gilroy (1985) applied a standard normal Z test to the rank-sum statistics. As highlighted, this modified test accounts for serial and cross correlation in the data. In the case here, the deseasonalised and detrended data shows little autocorrelation in the time series but possesses strong cross correlation between neighbouring pixels, i.e. $R > 0.99$.

The sum of W_{jk} for a block of ns pixels over the whole year, $\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}$, has mean

$$E\left(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}\right) = 12 \cdot ns \cdot \mu_w \quad (7)$$

and variance

$$Var\left(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk}\right) = \sum_{j=1}^{12} \sum_{k=1}^{ns} \sum_{h=1}^{ns} C(W_{jk}, W_{jh}). \quad (8)$$

Table 2. The interpretation of Z' score in the step trend test

$Z' \leq 0$ & rainfall decreases after change
$Z' \geq 0$ & rainfall increases after change
$Z' = 0$ & rainfall does not change

$C(W_{jk}, W_{jh})$ is the covariance of the W statistics between pixel k and pixel h in month j . When $k = h$,
 $C(W_{jk}, W_{jh}) = \sigma_w^2$. When $k \neq h$,

$$C(W_{jk}, W_{jh}) = \sigma_w^2 r(R_k, R_h) \quad (9)$$

where $r(R_k, R_h)$ is the product moment correlation coefficient of the concurrent ranks in pixel k and h . Here r is calculated on the full time series in each pixel. In the analysis, the test was applied to a square block of four pixels each time. As argued by Hirsch and Gilroy (1985), $ns = 4$ is the most optimal solution to balance the cost and the gain in the test power.

The statistic of the step trend test is then defined as

$$Z' = \frac{\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk} - 12 \cdot ns \cdot \mu_w}{\sqrt{\text{Var}(\sum_{j=1}^{12} \sum_{k=1}^{ns} W_{jk})}} \quad (10)$$

The above statistic is written for a 12 month period. By changing the value 12, it can also be used to test seasonal rainfall change or for other customized periods.

The null hypothesis (H_0) in this study is that there was no change in rainfall due to land surface intervention. The results of the step trend test can be interpreted according to the sign of the Z' score (see Table 2, Chapter 23, P887 (Hipel and McLeod 1994)). Z' is normally distributed similar to the standard normal statistics Z . Hence it can be compared to a standard normal distribution to determine the p value.

4 RESULTS

4.1 Tree cover change

The pixels, where the tree cover change based on the MOD44B data was significant ($p \leq 0.05$) in each study region, are shown in Figure 8 for the NSW/VIC region (left panel) and the QLD region (right panel). In the QLD region there actually has generally been an increase in tree cover after the clearing of native vegetation stopped. In the NSW/VIC region, much of the tree loss between 2002 and 2003 was concentrated in the Snowy Mountains close to the border of NSW and VIC, which is also evident in the figure. Tree cover loss occurred in large parts of the QLD region between 2002 and 2005, but this tree loss was spatially less concentrated.

4.2 Regression Model and significance of Vegetation Cover Changes

Generally the regression model only explains a limited amount of the rainfall variability (Figure @reg(fig:rsqmodel)). The model in Equation (3) accounts for an average around 33% of the variation in the NSW/VIC region and about 50% of the variation in the QLD region as indicated by the adjusted r^2 . The adjusted r^2 is the coefficient of determination, a measurement of the amount of variability predicted by the model adjusting for the number of explanatory terms. The residual analysis shows that the assumptions of the regression model are generally met (Figure 10). The standardised residual plots, however, show some funnelling for the NSW/VIC regions, suggesting non constant variance. The residual patterns are consistent for all pixels within each region.

In terms of predictors of the local rainfall in the model, logically, the average rainfall across the larger Murray Darling Basin is highly significant. This confirms that this variable is a good reflection of the year on year variability in the rainfall. The model also confirms the importance of the climate drivers and the seasonality in Australian rainfall, as many of these variables were significant. Even at the grid level, the seasons and several of the climatic indices were significant ($p \leq 0.05$) everywhere in both regions.




Figure 8. The maps show significant changes in tree cover identified from the MOD44B data between 2003 and 2015 in the NSW/VIC region (right) and the Qld region (left). The amount of change was calculated as the difference in tree cover before and after the specified land cover intervention and it is shown as the percentage of the ground area. Green colour indicates an increase in tree cover, while red colour indicates a decrease in tree cover.




Figure 9. The distribution of the variance explained (adjusted r^2) by the full regression model

371 The climate drivers (at lag zero) accounted, on average, for 6.5% of the rainfall variability in both the
372 QLD region and the NSW/VIC region (see Figure 11 for the distribution of adjusted r^2 in these two
373 regions). These figures are within the upper bound of seasonal rainfall predictability by a SST anomaly
374 field reported by Westra and Sharma (2010).

375 There were generally no statistically significant long term trends in both regions. However, this result
376 might not prove or disprove the existence a long term trend in rainfall. The overall time period is fairly
377 short (Koutsoyiannis 2006) and more pixels in NSW/VIC could indicate a significant step change if the
378 long term trend effect is not removed by the model. As trend free data is an important requirement for the
379 step trend test, the trend term was kept in the regression model to ensure the detection of step change was
380 not due to a possible long term trend.

381 The land cover variable in the model (Equation (3)) aims to identify a step change in the rainfall
382 before and after the observed change in land cover. The variable was mainly significant ($p \leq 0.05$) for the
383 rainfall estimates in some areas in NSW/VIC, as shown in Figure 12 (right panel). However, the number
384 of pixels where the landcover variable was significant was much greater in NSW/VIC compared to the
385 QLD region. There was also some relationship between the areas of bushfires in Figure 2. Only a very
386 small area with a significant step change due to the land cover changes was found in rainfall in the QLD
387 region (left panel)

388 More generally, the model for NSW/VIC suggests that the land cover variable has a positive impact
389 on rainfall in both regions. The fitted coefficients for the Land cover change variable were consistently
390 positive for the “tree” part of the series. It implies that rainfall was higher when the surface was covered
391 by trees.

392 4.3 Step Trend Test

393 The spatial step trend test based on the regression residuals without the landcover variable generates Z'
394 scores. The results for the Qld region (left) and NSWVIC region (right) are shown in Figure 13. This
395 figure provides two types of information: the sign and the significance level. The sign indicates the
396 direction of the step change in the residuals of the regression, as listed in Table 2. In both regions there
397 are areas of positive Z' values which imply a decrease in rainfall, but this is stronger in the NSW/VIC
398 area than in the QLD area. For the NSW/VIC area there appears to be a reasonable relationship between
399 the locations where changes in tree cover are observed (Figure @ref(fig:tc_trend)) and the patterns in the
400 Z' score. However, there is not necessarily a direct relationship as movement of air masses could mean
401 that actual changes of rainfall are observed close by, but not necessarily exactly at areas with changes of
402 landcover. There is once again a difference between the left panel and the right panel, indicating only
403 a few significant of Z' scores in the QLD region wich matches the earlier significance in the landcover
404 variable in the full regresion. In the QLD region, only 0.2% of the pixels obtained a positive Z' score with
405 $p < 0.1$. In the NSW/VIC region 3.7% of pixels have a positive Z' score with $p < 0.1$. In general it is
406 only a small proportion of both study regions.

407 The rainfall regression residuals without the landcover variable for the two regions (before-change
408 since 1979 and after-change) were also compared (Figure 14) using a simple t-test. From the boxplots it
409 is difficult to see that there were significantly different ($p < 0.05$) mean rainfall residual values between
410 the “before” and “after” periods in both regions. For the Queensland locations, the mean monthly
411 rainfall residuals were slightly lower ($p < 0.05$) after the change in landcover ($0.45mm/month$). For the
412 NSW/VIC locations there was larger decrease in the mean monthly rainfall residual ($p < 0.05$) after the
413 change($1.5mm/month$).

414 The choice of the number of pixels averaged in the step trend test (ns) has some impact on the test
415 results (Hirsch and Gilroy 1985). The cases of $ns = 1$ and $ns = 9$ were also tested. The results for $ns = 1$
416 showed a lower number of significant pixels(at $p < 0.10$) compared to the $ns = 4$ test. The results from
417 the $ns = 9$ test was not different from the $ns = 4$ test. We therefore decided to stay with $ns = 4$. The
418 power of the test does not change much after $ns = 4$, as shown by Hirsch and Gilroy (1985).

419 As part of the analysis, the “field significance” of the Z' score test was considered to improve
420 the interpretation of the step change at regional scales from multiple local tests (Wilks 2006, Westra,
421 Alexander, and Zwiers (2013)). Here, the bootstrapping resampling method from Westra, Alexander,
422 and Zwiers (2013) was used to evaluate the field significance. This means the spatial structure of the
423 pixels was maintained, but the order of the years and months was changed by random resampling. For
424 each resampling, the test statistic identifies the percentage of the pixels with a significant positive or




Figure 10. The residual analysis of a sample pixel in the QLD region (top) and NSW/VIC region (bottom).




Figure 11. The performance of the regression model if rainfall is only modelled by the climate drivers. It shows the percentage of rainfall variability that can be explained by the climate drivers for the Qld and NSW/VIC region




Figure 12. The spatial distribution of significance of land cover step change variable in the GAM model predicting changes in rainfall in the both regions. The left graph relates to the QLD region, while the right hand plot reflects the NSW/VIC region. The outlines of relevant Australian states and the Murray Darling Basin are indicated in grey. The p value reported is for the land cover variable in the model.




Figure 13. Spatial distribution of the step trend test Z' statistics in the two study sites. The panel on the left is for the QLD region, while the panel of the right is for NSW/VIC. Warm colours (yellow, orange and red) are for positive Z' values which indicate decreasing rainfall trend due to the land surface change. Cold colours (light blue to blue) are for negative Z' values which indicate increasing rainfall trend. The deeper the colour, the more significant the statistic.




Figure 14. Boxplots of annual rainfall residuals (estimated based on Equation 2 before and after the land cover intervention during 1979 - 2015 in the study regions. On average, the after period has a significantly lower annual rainfall residual in NSW/VIC, but a significantly higher annual rainfall residual in the Qld study area




Figure 15. Field significance results showing the distribution of percentage of significant Z' scores for the resampled series (blue bars), and the percentage significant Z' scores for the original series of rainfall years (red circle)

negative step change for the step trend test. The test statistics on 1000 resampled replicates were used to develop the distribution of these percentage values, and the observed fraction of significant Z' scores (at $p < 0.1$) is plotted on the distribution (Figure 15). The results support the earlier Z' score results. Only the percentage significant positive Z' scores for the NSW/VIC region falls on the tail of the field significance distribution, while all the others are well within the distribution. This suggests that the percentage of significant positive Z' scores in the NSW/VIC region is least likely due to a unique series of rainfall years, but is most likely due to changes in landcover. In other words, in the NSW/VIC region it is most likely that the change in landcover caused a decrease in the rainfall.

5 DISCUSSION

Overall the summary table of the results indicates that the effects of land clearing or bushfire on rainfall are consistent across the range of different statistical tests. In all cases, the data for the NSW/VIC region indicates the strongest evidence that the reduction in tree cover (the land cover change) resulted in a local decrease in rainfall. In contrast the data for the QLD region only indicates weak evidence that the landclearing resulted in reduction in the rainfall.

Generally, empirical studies on LCC-precipitation interaction are conducted within an area with known land surface intervention (e.g. Otterman et al. 1990, Durieux, Machado, and Laurent (2003), Negri et al. (2004), Sato, Kimura, and Hasegawa (2007)). However, these locations are rare and difficult to

Table 3. Summary table of all tests on the two regions

Test	Qld	NSW/VIC
LC variable	Small area of significant pixels \& in the centre	Large area possibly aligning \& with bushfire affected area.
t-test regression \& model residuals	Slightly lower mean residual ($p < 0.05$) \& after change (0.5 mm/month)	Lower mean residual ($p < 0.50$) \& after change (1.5 mm/month)
Positive Z' score	0.2\% of pixels at $p < 0.1$	3.7\% of pixel at $p < 0.1$
Field significance \& /& of Z'scores	Both positive and negative \& \& within random distribution	Positive \& outside random distribution.

isolate from real landscape change. Modelling studies are abundant (e.g. Chagnon and Bras 2005; Pinto et al. 2009; Wang et al. 2009), but these are generally not directly linked to observed data. In this study we tested the effect of land cover change across a broad region, which included locations where changes were known to occur or have occurred. The advantage of the suggested approach is that it does not require a long time series of land cover data as this is usually unavailable. Furthermore, it does not assume a specific relationship between vegetation cover change and rainfall but allows the data to show this relationship, by applying the analysis to a broader area outside the boundary of the vegetation cover change. This approach is expected to provide a way to reduce the risk of a false positive paradox, by comparing results between areas with and without vegetation cover change.

While there is some indication that the observed landuse changes (Figure 8) cause a decrease in the rainfall, there are several possible complicating factors in the data that require discussion.

5.1 Rainfall variability

Rainfall in Australia is highly variable from year to year, and the time period in this study included a severe drought (Dijk and Viney 2013), as well as the drought breaking years 2010 and 2011. The purpose of using the regression model is to remove the year to year and month to month variability in rainfall and therefore strengthen the land cover signal in the residuals. We used the spatially averaged monthly rainfall timeseries for the larger Murray Darling Basin (Figure 1) to remove the general variability. However, overall the model does not explain more than 50% (QLD) and 30% (NSW/VIC) of the rainfall variability, and only around 7% on average is due to the climate drivers (Figure 11). And while this is consistent with the literature e.g. Westra and Sharma (2010), this means some variation due to external factors could still be left in the residuals. Rainfall is generally considered a stochastic process (e.g. Fowler et al. 2005, Cowpertwait, Salinger, and Mullen (2009), Burton et al. (2010)) and some of the variability could either be a different response to a combination of climate factors (as interactions were not tested in the model), or a non-stationary response to the climate drivers. The severe bushfires in 2003 were also triggered by the extreme drought conditions during the millenium drought (Dijk and Viney 2013). Although the drought on rainfall has been accounted for in part by the model, a further delayed or cumulative impact of the drought could be feeding into the local land-atmosphere interaction. As a result, the rainfall feedback to the vegetation cover change could be weak under the dry conditions between 2001 and 2009, and this could have affected the result.

5.2 Vegetation dynamics

The second possible effect is the dynamic nature of the vegetation clearing and recovery, especially for the QLD region. Although land clearing has occurred at a high rate and broad scale in Queensland (Department of Science, Information Technology and Innovation 2017), the clearing does not have a clear start and end point. QLD has a long history of land clearing. According to the series of SLATS reports on land cover changes in QLD released by the Queensland government, land clearing continued in and around the study region between 1988 - 2008. Major broad scale and high rate clearings occurred in 1999 - 2000 and 2002 - 2004 (Figure 16). And even though there was a decrease in land clearing post 2005, it is difficult to define a clear cut change in this region. The specific vegetation class in the Queensland area is well-known for rapid regrowth and "thickening" in favourable conditions (Gowen and Bray 2016), and this could explain the change in vegetation cover between 2009 and 2015 in the Qld area (Figure 8). In particular the favourable rainfall years of 2010 and 2011 would have boosted regrowth, increasing evapotranspiration and therefore decreasing the effect of the rainfall change.

The different causes of vegetation cover change in these two regions could lead to different post-change characteristics. The trends in EVI in the QLD region are less than in NSW/VIC region, based on the DLCD data. This is most likely due to the lower tree density in the QLD region than in the NSW/VIC region before land surface interventions. In contrast, the significant bushfires (Figure 2) would have drastically reduced the vegetation cover and recovery was very slow in some areas within the NSW/VIC




Figure 16. Woody vegetation clearing rate in QLD for the two major bioregions in the study area. The data were obtained from SLATS (2017), and clearly indicate a sharp decrease in the clearing rate after 2005.

area. The persistent drought in the 2000s (Howden 2012,Dijk and Viney (2013)) delayed the regrowth of trees. Conversely, replacing tree cover with pasture and crops in the Qld area might only have a relatively subtle impact on the EVI.

5.3 Gridded monthly rainfall data

The rainfall data used in this study is gridded. This data set is robust and consistent over a long time series (from 1900 to current) and has a broad national wide coverage which can provide more information spatially (Jeffrey et al. 2001; Tozer, Kiem, and Verdon-Kidd 2009). However, high cross correlation between pixels, due to the interpolation method in this data set (Jeffrey et al. 2001), can also introduce spatial noise. In the step trend test the cross correlation was accounted for. However, some other methods are also available, which could be used to perform a comparative trial. For example, (Narisma et al. 2007) applied a spatial Gaussian filter on a similar data set and used wavelet analysis to detect step change in rainfall. High quality station data is another option to test whether the observed spatial pattern in the step trend test results was not due to the gridded data itself. Resampling methods, such as bootstrapping and permutation (used in this study) (Wilks 1997,Kundzewicz and Robson (2004),Westra, Alexander, and Zwiers (2013)), can also be used to further assess the strength of the significance of results . While the gridded data set is most useful in regions with sparse rain gauge networks, such as in this study, it can actually reduces information where the rain gauge density is high (Jones, Wang, and Fawcett 2009). In the NSW/VIC region, the coverage of rainfall stations is more intensive, but they are mainly located in the valleys. As a result, the interpolated data might be a limited representation of the true local rainfall.

5.4 General approach

Parametric tests are generally more powerful than nonparametric test in detecting a trend, when the data is normally distributed (Onoz and Bayazit 2003,Kundzewicz and Robson (2004)). As a non-parametric test, the step trend test has the advantages of being distribution free and having no restriction on missing data (Hirsch and Gilroy 1985). This is particularly useful in rainfall analysis since rainfall data is usually highly skewed. On the other hand, the disadvantages of non-parametric tests, such as being limited to

514 hypothesis testing and being weaker in power, also hold for the step trend test (Whitley and Ball 2002).
515 Overall, the current study provides a statistical data based approach building on several lines of
516 evidence to reject the null hypothesis (no step change in rainfall occurs as a result of tree cover loss) at
517 least for the NSW/VIC region. Limited by the available data, the time frame under study included a long
518 lasting drought period (Holper 2011, Dijk and Viney (2013)). The strong impact of this prolonged drought
519 might have suppressed the land-atmosphere interaction and modified the cause and effect relationship
520 between rainfall and vegetation cover change. This could be one of the reasons that the LCC effects on
521 the local climate found in other studies (e.g. Görgen et al. 2006, McAlpine et al. (2007)) are not as strong
522 and do not appear in the QLD region. Possible future work could focus on a non-drought period, or using
523 a different series of land cover data, possibly on another continent. The recent large bushfires on the US
524 continent come to mind as a possible candidate study. While the power of the test can be improved with the
525 longer length of the after-intervention period (Hirsch and Gilroy 1985), the dynamic nature of vegetation
526 regrowth in this case study also affects this effect. A better approach might be to build a global study
527 that investigates multiple locations where drastic landcover changes have taken place, which would also
528 remove some of the climate variability effects due to the larger sample size.

529 **6 CONCLUSIONS**

530 In this study, we present a statistical data based approach to identify the impact of a change in land cover
531 on local rainfall. The results, based on gridded rainfall data found, although not strong, that a reduction in
532 vegetation cover is likely to have reduced local rainfall for a large area in NSW/VIC affected by bushfires.
533 However, land clearing in QLD was unlikely to have reduced rainfall over the same time period.

534 Drought can have a pronounced impact on the land surface condition during the study period, also
535 leading to significant reduction in vegetation cover and extreme events such as bushfires. The lack
536 of rainfall and associated high temperatures may mask the impact on rainfall of a step change in the
537 vegetation cover. Hence, the signal of LCC feedback on rainfall is probably weaker under such regional
538 dry conditions, as the impact of LCC on rainfall is mainly through changes in moisture convergence
539 (Görgen et al. 2006, Pitman and Hesse (2007)).

540 **7 ACKNOWLEDGMENTS**

541 CL was supported by an Australian Postgraduate Award for this work.

542 **8 APPENDIX**

543 **8.1 Summary of Data**

Table 4. Summary of data.

Data	Source	Resolution		Analysis period
		Temporal	Spatial	
Percent tree cover	MOD44B	Annual	250m	2000-2010
Trend of vegetation cover change	DLCD (2009)	Onetime	250m	Trend of Apr 2000 - Apr 2008
Rainfall	AWAP gridded rainfall data	Monthly	$0.05^{\circ} \times 0.05^{\circ}$	Jan 1979- Dec 2008
SOI	BoM	Monthly	N/A	Jan 1979- Dec 2008
NINO 3, 3.4, 4	IRI/LDEO data library	Monthly	N/A	Jan 1979- Dec 2008
PDO	NOAA	Monthly	N/A	Jan 1979- Dec 2008
IOD	POAMA-2 dataset	Monthly	N/A	Jan 1979- Dec 2008

REFERENCES

- ABARES. 2010. "Land Use of Australia, Version 4, 2005/2006."
- ABC News. 2003. "Kosciuszko Slow to Recover from Bushfires."
- Australian Bureau of Statistics. 2012. "Geography and Climate - Australia's Climate."
- Ben-Gai, T., A. Bitan, A. Manes, P. Alpert, and S. Rubin. 1998. "Spatial and Temporal Changes in Rainfall Frequency Distribution Patterns in Israel." *Theoretical and Applied Climatology* 61 (3-4): 177-90. <Go to ISI>://000078025000005.
- BoM. 2012a. "Australia - Climate of Our Continent."
- . 2012b. "Australian Climate Influence."
- Bosilovich, Michael G., and Jiun-Dar Chern. 2006. "Simulation of Water Sources and Precipitation Recycling for the MacKenzie, Mississippi, and Amazon River Basins." *Journal of Hydrometeorology* 7 (3): 312-29. <http://dx.doi.org/10.1175%2FJHM501.1>.
- Burton, A, HJ Fowler, S Blenkinsop, and CG Kilsby. 2010. "Downscaling Transient Climate Change Using a Neyman-Scott Rectangular Pulses Stochastic Rainfall Model." *Journal of Hydrology* 381 (1): 18-32.
- Chagnon, F. J. F., and R. L. Bras. 2005. "Contemporary Climate Change in the Amazon." *Geophysical Research Letters* 32 (13).
- Chowdhury, R. K., and S. Beecham. 2010. "Australian Rainfall Trends and Their Relation to the Southern Oscillation Index." *Hydrological Processes* 24 (4). John Wiley & Sons, Ltd.: 504-14. <http://dx.doi.org/10.1002/hyp.7504>.
- Cowpervait, PSP, Jim Salinger, and B Mullen. 2009. "A Spatial-Temporal Stochastic Rainfall Model for Auckland City: Scenarios for Current and Future Climates." *Journal of Hydrology (NZ)* 48: 95-109.
- Deo, RC, JI Syktus, CA McAlpine, and KK Wong. 2009. "The Simulated Impact of Land Cover Change on Climate Extremes in Eastern Australia." In *Proceedings of the 18th World Imacs Congress and Modsim09 International Congress on Modelling and Simulation*, 2035-41. Modelling; Simulation Society of Australia; New Zealand Inc.; International Association for Mathematics; Computers in Simulation.
- Department of Natural Resources and Mines. 2005. "Land Cover Change in Queensland 2001-2003, Incorporating 2001-2002 Adn 2002-2003 Change Periods: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Natural Resources; Mines.
- Department of Natural Resources and Water. 2007. "Land Cover Change in Queensland 2004-2005: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Natural Resources; Water.
- Department of Science, Information Technology and Innovation. 2017. "Land Cover Change in Queensland 2015-2016: A Statewide Landcover and Trees Study (SLATS) Report." Brisbane: Department of Science, Information Technology; Innovation.
- Dijk, Beck van, Albert I. J. M., and Neil R. Viney. 2013. "The Millennium Drought in Southeast Australia (2001-2009): Natural and Human Causes and Implications for Water Resources, Ecosystems, Economy, and Society." *Water Resources Research* 49 (2): 1040-57. doi:10.1002/wrcr.20123.
- Dirmeyer, P. A., K. L. Brubaker, and T. DelSole. 2009. "Import and Export of Atmospheric Water Vapor Between Nations." *Journal of Hydrology* 365 (1-2): 11-22. <Go to ISI>://000263575600003.
- Durieux, Laurent, Luiz Augusto Toledo Machado, and Henri Laurent. 2003. "The Impact of Deforestation on Cloud Cover over the Amazon Arc of Deforestation." *Remote Sensing of Environment* 86 (1): 132-40.
- Eltahir, E. A. B., and R. L. Bras. 1996. "Precipitation Recycling." *Reviews of Geophysics* 34 (3): 367-78. <Go to ISI>://A1996VD58700003.
- Fowler, HJ, CG Kilsby, PE O'connell, and A Burton. 2005. "A Weather-Type Conditioned Multi-Site Stochastic Rainfall Model for the Generation of Scenarios of Climatic Variability and Change." *Journal of Hydrology* 308 (1): 50-66.
- Gaertner, M. A., O. B. Christensen, J. A. Prego, J. Polcher, C. Gallardo, and M. Castro. 2001. "The Impact of Deforestation on the Hydrological Cycle in the Western Mediterranean: An Ensemble Study with Two Regional Climate Models." *Climate Dynamics* 17 (11): 857-73.
- Gallant, Ailie J. E., Kevin J. Hennessy, and James Risbey. 2007. "Trends in Rainfall Indices for Six Australian Regions: 1910 - 2005." *Australian Meteorological Magazine* 56: 223-39.
- Gimeno, Luis, Anita Drumond, Raquel Nieto, Ricardo M. Trigo, and Andreas Stohl. 2010. "On the Origin of Continental Precipitation." *Geophysical Research Letters* 37 (13). AGU. <http://dx.doi.org>.

org/10.1029/2010GL043712.

Gowen, Rebecca, and Steven G. Bray. 2016. "Bioeconomic Modelling of Woody Regrowth Carbon Offset Options in Productive Grazing Systems." *The Rangeland Journal* 38 (3): 307–17. doi:https://doi.org/10.1071/RJ15084.

Görgen, K., A. H. Lynch, A. G. Marshall, and J. Beringer. 2006. "Impact of Abrupt Land Cover Changes by Savanna Fire on Northern Australian Climate." *Journal of Geophysical Research-Atmospheres* 111 (D19): D19106. <http://dx.doi.org/10.1029/2005JD006860>.

Hastie, Trevor, and Robert Tibshirani. 1986. "Generalized Additive Models." *Statistical Science*, 297–310.

Hipel, K.W., and A.I. McLeod. 1994. *Time Series Modelling of Water Resources and Environmental Systems*. Vol. 45. Elsevier Science Ltd.

Hirsch, Robert M., and Edward J. Gilroy. 1985. "Detectability of Step Trends in the Rate of Atmospheric Deposition of Sulfate." *JAWRA Journal of the American Water Resources Association* 21 (5): 773–84. <http://dx.doi.org/10.1111/j.1752-1688.1985.tb00171.x>.

Holper, Paul N. 2011. *Climate Change, Science Information Paper: Australian Rainfall-Past, Present and Future*. Canberra: CSIRO.

Howden, Saffron. 2012. "It's Official: Australia No Longer in Drought." *Brisbane Times*.

Hughes, Lesley. 2003. "Climate Change and Australia: Trends, Projections and Impacts." *Austral Ecology* 28 (4). Blackwell Science Pty: 423–43. <http://dx.doi.org/10.1046/j.1442-9993.2003.01300.x>.

Jeffrey, Stephen J., John O. Carter, Keith B. Moodie, and Alan R. Beswick. 2001. "Using Spatial Interpolation to Construct a Comprehensive Archive of Australian Climate Data." *Journal Article. Environmental Modelling & Software* 16 (4 SU-): 309–30. <http://www.sciencedirect.com/science/article/B6VHC-42SXF8F-2/2/b5c0a24b85149f40ba5d2419ddc0999c>.

Jones, D.A., W. Wang, and R. Fawcett. 2009. "High-Quality Spatial Climate Data-Sets for Australia." *Australian Meteorological and Oceanographic Journal* 58 (4): 233–48.

Junkermann, W., J. Hacker, T. Lyons, and U. Nair. 2009. "Land Use Change Suppresses Precipitation." *Atmospheric Chemistry and Physics* 9 (17): 6531–9. <Go to ISI>://WOS:000269778500018.

Kamruzzaman, M., S. Beecham, and A. V. Metcalfe. 2011. "Non-Stationarity in Rainfall and Temperature in the Murray Darling Basin." *Hydrological Processes* 25 (10). John Wiley & Sons, Ltd.: 1659–75. <http://dx.doi.org/10.1002/hyp.7928>.

Koutsoyiannis, Demetris. 2006. "Nonstationarity Versus Scaling in Hydrology." *Journal Article. Journal of Hydrology* 324 (1-4): 239–54. <http://www.sciencedirect.com/science/article/B6V6C-4HP6GJ9-1/2/67dcc2e8f8e75b7abb918617f5a79a37>.

Kucharski, Fred, Ning Zeng, and Eugenia Kalnay. 2013. "A Further Assessment of Vegetation Feedback on Decadal Sahel Rainfall Variability." *Climate Dynamics* 40 (5-6): 1453–66. <http://link.springer.com/article/10.1007/s00382-012-1397-x>.

Kuczera, George. 1987. "Prediction of Water Yield Reductions Following a Bushfire in Ash-Mixed Species Eucalypt Forest." *Journal of Hydrology* 94 (3-4): 215–36. doi:https://10.1016/0022-1694(87)90054-0.

Kundzewicz, Z.W., and A.J. Robson. 2004. "Change Detection in Hydrological Records-a Review of the Methodology/Revue Méthodologique de La Détection de Changements Dans Les Chroniques Hydrologiques." *Hydrological Sciences Journal* 49 (1).

Lymburner, Leo, Peter Tan, Norman Mueller, Richard Thackway, Adam Lewis, Medhavy Thankappan, Lucy Randall, Anisul Islam, and Udaya Senarath. 2010. "250 Metre Dynamic Land Cover Dataset." Geoscience Australia, Canberra.

Ma, H. Y., C. R. Mechoso, Y. Xue, H. Xiao, C. M. Wu, J. L. Li, and F. De Sales. 2011. "Impact of Land Surface Processes on the South American Warm Season Climate." *Climate Dynamics* 37 (1-2): 187–203. <Go to ISI>://WOS:000293403000012.

MacDonald, Glen M., and Roslyn A. Case. 2005. "Variations in the Pacific Decadal Oscillation over the Past Millennium." *Geophysical Research Letters* 32 (8): L08703. <http://dx.doi.org/10.1029/2005GL022478>.

McAlpine, C. A., J. Syktus, R. C. Deo, P. J. Lawrence, H. A. McGowan, I. G. Watterson, and S. R. Phinn. 2007. "Modeling the Impact of Historical Land Cover Change on Australia's Regional Climate."

653 *Geophysical Research Letters* 34 (22). <Go to ISI>://000251345100005.

654 Mei, R., and G. L. Wang. 2010. "Rain Follows Logging in the Amazon? Results from CAM3-CLM3." *Climate Dynamics* 34 (7-8): 983–96. <Go to ISI>://WOS:000278088400005.

655 Meneghini, Belinda, Ian Simmonds, and Ian N. Smith. 2007. "Association Between Australian Rainfall and the Southern Annular Mode." *International Journal of Climatology* 27 (1): 109–21. <http://dx.doi.org/10.1002/joc.1370>.

656 Murphy, Bradley F., and Bertrand Timbal. 2008. "A Review of Recent Climate Variability and Climate Change in Southeastern Australia." *International Journal of Climatology* 28 (7): 859–79. <http://dx.doi.org/10.1002/joc.1627>.

657 Nair, Udaysankar S., Y. Wu, J. Kala, T. J. Lyons, Sr. Pielke R. A., and J. M. Hacker. 2011. "The Role of Land Use Change on the Development and Evolution of the West Coast Trough, Convective Clouds, and Precipitation in Southwest Australia." *Journal of Geophysical Research-Atmospheres* 116 (D7). AGU: D07103. <http://dx.doi.org/10.1029/2010JD014950>.

658 Narisma, G.T., J.A. Foley, R. Licker, and N. Ramankutty. 2007. "Abrupt Changes in Rainfall During the Twentieth Century." *Geophysical Research Letters* 34 (6): L06710.

659 Negri, A. J., R. F. Adler, L. M. Xu, and J. Surratt. 2004. "The Impact of Amazonian Deforestation on Dry Season Rainfall." *Journal of Climate* 17 (6): 1306–19.

660 Nicholls, N. 2006. "Detecting and Attributing Australian Climate Change: A Review." *Australian Meteorological Magazine* 55 (3). Bureau of Meteorology.: 199–211.

661 Oleson, K. W., G. B. Bonan, S. Levis, and M. Vertenstein. 2004. "Effects of Land Use Change on North American Climate: Impact of Surface Datasets and Model Biogeophysics." *Climate Dynamics* 23 (2): 117–32.

662 Onoz, B., and M. Bayazit. 2003. "The Power of Statistical Tests for Trend Detection." *Turkish Journal of Engineering and Environmental Sciences* 27 (4): 247–51.

663 Otterman, J., A. Manes, S. Rubin, P. Alpert, and D. Starr. 1990. "An Increase of Early Rains in Southern Israel Following Land-Use Change?" *Boundary-Layer Meteorology* 53 (4): 333–51.

664 Pinto, E., Y. Shin, S. A. Cowling, and C. D. Jones. 2009. "Past, Present and Future Vegetation-Cloud Feedbacks in the Amazon Basin." *Climate Dynamics* 32 (6): 741–51. <Go to ISI>://WOS:000264118400001.

665 Pitman, A. J., and P. P. Hesse. 2007. "The Significance of Large-Scale Land Cover Change on the Australian Palaeomonsoon." *Quaternary Science Reviews* 26 (1-2): 189–200.

666 Pitman, A. J., and R. Lorenz. 2016. "Scale Dependence of the Simulated Impact of Amazonian Deforestation on Regional Climate." *Environmental Research Letters* 11 (9): 094025. doi:10.1088/1748-9326/11/9/094025.

667 Pitman, A. J., G. T. Narisma, R. A. Pielke, and N. J. Holbrook. 2004. "Impact of Land Cover Change on the Climate of Southwest Western Australia." *Journal of Geophysical Research-Atmospheres* 109 (D18). <Go to ISI>://WOS:000224126500001.

668 R Core Team, 2018. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>.

669 Risbey, James S., Michael J. Pook, Peter C. McIntosh, Matthew C. Wheeler, and Harry H. Hendon. 2009. "On the Remote Drivers of Rainfall Variability in Australia." *Monthly Weather Review* 137 (10): 3233–53. <http://dx.doi.org/10.1175/2009MWR2861.1>.

670 Roy, David P, Luigi Boschetti, Christopher O Justice, and J Ju. 2008. "The Collection 5 Modis Burned Area Product—Global Evaluation by Comparison with the Modis Active Fire Product." *Remote Sensing of Environment* 112 (9). Elsevier: 3690–3707.

671 Roy, DP, Y Jin, PE Lewis, and CO Justice. 2005. "Prototyping a Global Algorithm for Systematic Fire-Affected Area Mapping Using Modis Time Series Data." *Remote Sensing of Environment* 97 (2). Elsevier: 137–62.

672 Roy, DP, PE Lewis, and CO Justice. 2002. "Burned Area Mapping Using Multi-Temporal Moderate Spatial Resolution Data—A Bi-Directional Reflectance Model-Based Expectation Approach." *Remote Sensing of Environment* 83 (1). Elsevier: 263–86.

673 Saha, Subodh K., Paul A. Dirmeyer, and Thomas N. Chase. 2016. "Investigating the Impact of Land-Use Land-Cover Change on Indian Summer Monsoon Daily Rainfall and Temperature During 1951-2005 Using a Regional Climate Model." *Hydrology and Earth System Sciences* 20 (5): 1765.

707 [http://search.proquest.com/openview/c4242c9ecce075e3545390b1d00f2751/](http://search.proquest.com/openview/c4242c9ecce075e3545390b1d00f2751/1?pq-origsite=gscholar&cbl=105724)
708 [1?pq-origsite=gscholar&cbl=105724](http://search.proquest.com/openview/c4242c9ecce075e3545390b1d00f2751/1?pq-origsite=gscholar&cbl=105724).

709 Saji, NH, B.N. Goswami, PN Vinayachandran, and T. Yamagata. 1999. "A Dipole Mode in the
710 Tropical Indian Ocean." *Nature* 401 (6751): 360–63.

711 Sato, T., F. Kimura, and A. S. Hasegawa. 2007. "Vegetation and Topographic Control of Cloud
712 Activity over Arid/Semiarid Asia." *Journal of Geophysical Research-Atmospheres* 112 (D24).

713 Schepen, Andrew, Q. J. Wang, and David Robertson. 2012. "Evidence for Using Lagged Climate
714 Indices to Forecast Australian Seasonal Rainfall." *Journal of Climate* 25 (4). American Meteorological
715 Society: 1230–46.

716 Semazzi, F. H. M., and Y. Song. 2001. "A GCM Study of Climate Change Induced by Deforestation
717 in Africa." *Climate Research* 17 (2): 169–82.

718 Smith, Ian N., and Bertrand Timbal. 2012. "Links Between Tropical Indices and Southern Australian
719 Rainfall." *International Journal of Climatology* 32 (1): 33–40.

720 Speer, Milton, Lance Leslie, and Alexandre Fierro. 2011. "Australian East Coast Rainfall Decline
721 Related to Large Scale Climate Drivers." *Climate Dynamics* 36 (7): 1419–29. [http://dx.doi.org/](http://dx.doi.org/10.1007/s00382-009-0726-1)
722 [10.1007/s00382-009-0726-1](http://dx.doi.org/10.1007/s00382-009-0726-1).

723 The State Government of Victoria. 2011. "Bushfire History."

724 Timbal, B., and J.M. Arblaster. 2006. "Land Cover Change as an Additional Forcing to Explain the
725 Rainfall Decline in the South West of Australia." *Geophysical Research Letters* 33 (7): L07717.

726 Townshend, J.R.G., M. Carroll, C. Dimiceli, R. Sohlberg, M. Hansen, and R. DeFries. 2011. "Veg-
727 etation Continuous Fields MOD44B, 2001 Percent Tree Cover, Collection 5." University of Maryland,
728 College Park, Maryland, 2001.

729 Tozer, C. R., A. S. Kiem, and D. C. Verdon-Kidd. 2009. "On the Uncertainties Associated with
730 Using Gridded Rainfall Data as a Proxy for Observed." Conference Paper. In *18th World Imacs Congress*
731 *and Modsim09 International Congress on Modelling and Simulation.*, edited by R.S. Anderssen, R.D.
732 Braddock, and L.T.H. Newham, 3886–92. <http://www.mssanz.org.au/modsim09/I13/reesley.pdf>.

733 ———. 2012. "On the Uncertainties Associated with Using Gridded Rainfall Data as a Proxy for
734 Observed." Journal Article. *Hydrol. Earth Syst. Sci.* 16 (5): 1481–99. doi:10.5194/hess-16-1481-2012.

735 Trenberth, K. E. 1999. "Atmospheric Moisture Recycling: Role of Advection and Local Evaporation."
736 *Journal of Climate* 12 (5): 1368–81. <Go to ISI>://000080145700002.

737 Verdon, Danielle C., Adam M. Wyatt, Anthony S. Kiem, and Stewart W. Franks. 2004. "Multidecadal
738 Variability of Rainfall and Streamflow: Eastern Australia." *Water Resources Research* 40 (10): W10201.
739 <http://dx.doi.org/10.1029/2004WR003234>.

740 Wang, Jingfeng, Frederic Chagnon, Earle Williams, Alan Betts, Nilton Renno, Luiz Machado, Gautam
741 Bisht, Ryan Knox, and Rafael Bras. 2009. "Impact of Deforestation in the Amazon Basin on Cloud
742 Climatology." *Proceedings of the National Academy of Sciences of the United States of America* 106 (10):
743 3670.

744 Westra, Seth, Lisa V. Alexander, and Francis W. Zwiers. 2013. "Global Increasing Trends in Annual
745 Maximum Daily Precipitation." *Journal of Climate* 26 (11). American Meteorological Society: 3904–18.
746 <http://dx.doi.org/10.1175/JCLI-D-12-00502.1>.

747 Westra, Seth, and Ashish Sharma. 2010. "An Upper Limit to Seasonal Rainfall Predictability?"
748 *Journal of Climate* 23 (12). American Meteorological Society: 3332–51.

749 Whitley, Elise, and Jonathan Ball. 2002. "Statistics Review 6: Nonparametric Methods." *Critical*
750 *Care, London* 6 (6): 509–13.

751 Wilks, D. S. 2006. "On 'Field Significance' and the False Discovery Rate." *Journal of Applied*
752 *Meteorology and Climatology* 45 (9). American Meteorological Society: 1181–9.

753 Wilks, DS. 1997. "Resampling Hypothesis Tests for Autocorrelated Fields." *Journal of Climate* 10
754 (1): 65–82.

755 Wood, S. N. 2011. "Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation
756 of Semiparametric Generalized Linear Models." *Journal of the Royal Statistical Society (B)* 73 (1): 3–36.

757 Zanchettin, Davide, Stewart W. Franks, Pietro Traverso, and Mario Tomasino. 2008. "On Enso
758 Impacts on European Wintertime Rainfalls and Their Modulation by the Nao and the Pacific Multi-
759 Decadal Variability Described Through the Pdo Index." *International Journal of Climatology* 28 (8). John
760 Wiley & Sons, Ltd.: 995–1006. <http://dx.doi.org/10.1002/joc.1601>.

761 Zeng, X. M., Z. H. Wu, S. Song, S. Y. Xiong, Y. Q. Zheng, Z. G. Zhou, and H. Q. Liu. 2012. "Effects

762 of Land Surface Schemes on the Simulation of a Heavy Rainfall Event by WRF.” *Chinese Journal of*
763 *Geophysics-Chinese Edition* 55 (1): 16–28. <Go to ISI>://WOS:000300129300002.