


Vignette EcoHydro2D v0.0.1

Willem Vervoort

3 September 2016

Introduction

This is the vignette demonstrating the alpha version of the ecohydr2d package. The package contains functions based on the ecohydrological work from Vervoort and van der Zee (2012) but extended to include a 1-D models included in a 2-D grid overlying a single cell groundwater system modelled with a separate fortran groundwater model developed by Prof. Toon Leijnse (gwt.exe).



Vervoort R.W. and S.E.A.T.M. van der Zee (2012) On stochastic modelling of groundwater uptake in semi-arid water-limited systems: root density and seasonality effects. *Ecohydrology* doi:10.1002/eco.1288.

First example: uniform rainfall

This first examples shows how the program is set up and how a simple simulation of a basic model with 10 grid cells with a uniform vegetation can be set up. This includes the set up of the input files for gwt.exe as well as the overall model setup and analysis of the output

Step 1: setup of packages and working directory

This currently assumes that the package EcoHydro2D is installed and can be loaded. Once I have worked out how Git and devtools work together I will change this.

```
require(ggplot2)
```

```
## Loading required package: ggplot2
```

```
require(EcoHydro2D)
```

```
## Loading required package: EcoHydro2D
```

```
require(knitr)
```

```
## Loading required package: knitr
```

```
## Warning: package 'knitr' was built under R version 3.2.5
```

```
today <- format(Sys.Date(), "%Y%m%d")
```

Step 2 define inputs: uniform rainfall for two years

We will start with a scenario which simulates uniform rainfall and uniform potential ET on a 10 cell linear grid in the x-direction (so the y-dimension is 1). Units in the model are cm and days.

The first step will be to define the three input time series that are needed for the model:

- Rainfall, uniform 0.25 cm/day
 - Streamflow (which is set to 0 here)
 - potential ET, uniform 0.5 cm/day
- The date values are arbitrary.
The model is still quite slow, so starting off with only 3 months of data.

```
days <- seq(as.Date("2005-01-01"), as.Date("2005-3-31"), by=1)
Rain_uni <- data.frame(Dates = days, Rain = rep(0.1, length(days)))
# start with no water in Stream
Stream0 <- data.frame(Dates = days, Height = rep(0, length(days)))
# and uniform ETp
ETp_uni <- data.frame(Dates = days, ET = rep(0.5, length(days)))
```

2.1 define groundwater model parameters

One of the things to do is to define the parameters for the groundwater model. This is slightly cryptic, but is described in more detail in the manual of gwt.exe which is provided in the documents folder. Currently the important values are hidden in a list under `ecohydro2d.options()`

```
# show the options
ecohydro2d.options()
```

```
## $NX
## [1] 12
##
## $NY
```

```

## [1] 1
##
## $NRBL
## [1] 2
##
## $IXR
## [1] 1 12
##
## $IYR
## [1] 1 1
##
## $DELR
## [1] 20 20 40 60 80 100 120 140 160 180 200 20
##
## $DELY
## [1] 100
##
## $ITIM
## [1] 1
##
## $bottom
## [1] -25
##
## $GWthreshold
## [1] 0
##
## $DELTcrit
## [1] 21
##
## $dslope_x
## [1] 0
##
## $dslope_y
## [1] 0
##
## $RES
## [1] 100

```

```
# you can now call different elements and reset them
```

```

# Initial groundwater heads
gw_in <- rep(-6,ecohydro2d.options()$NX)
# this is needed for Trees with deep roots
Zmean <- rep(600,ecohydro2d.options()$NX)

```

step 3 define the soils and vegetation

We can now define the vegetation and the soils, starting off with uniform grass.

```

# Define the vegetation series as being uniform grass
veg <- rep("Grass",ecohydro2d.options()$NX)
# Define what soil to use
soils <- "L Med Clay"

```

```
sp <- Soil_cpp("soils")
#
```

step 4 Run the model over the grid cells

Finally the model can be run via `big_fun`. I need to come up with a better name for this function. There is the ability to specify a separate Ks for the aquifer `aq_K` and also a separate specific yield via `aq_specy`.

```
result <- big_fun(N=nrow(Rain_uni),stype=soils,vtype=veg,
  aq_K = sp$K_s/100,
  aq_specy = sp$spec_y,
  Rain=Rain_uni, ETp=ETp_uni,
  stream=Stream0, gwheads = gw_in,
  Zmean = Zmean,today = today)
```

step 5 plotting the results

Now some plotting of the results to demonstrate how the function works. The output of `big_fun()` is large data frame with all the different variables. The Table below lists all the different columns.

```
names(result)
```

```
## [1] "id"           "s"           "qcap"        "Tgw"
## [5] "Tsoil"       "Ttotal"      "leakage"     "gwrech"
## [9] "smloss"      "static_stress" "surfoff"    "x"
## [13] "vtype"       "gwlevel"     "qrivlat"     "rain"
## [17] "stream"
```

```
kable(read.csv("figs/big_funColumnExplanation.csv"))
```

Name	Explanation
id	simple counter for location
s	relative soil moisture
qcap	capillary flow from groundwater into soil profile
Tgw	Transpiration from groundwater via deep roots
Tsoil	Transpiration from soil
Ttotal	Total transpiration (Evapotranspiration)
leakage	Leakage from soil profile to groundwater
gwrech	Recharge into groundwater = Leakage
smloss	loss of soil moisture for each time step
static_stress	static stress of the vegetation
surfoff	surface runoff
x	distance along the x grid from the river
vtype	vegetation type for each grid cell
gwlevel	groundwater level
qrivlat	lateral flow from river to groundwater
rain	Rainfall
stream	Streamflow

First some preparation of the data to make the plotting easier.


```
# use stackfun
Out <- stackfun(result, ecohydro2d.options())$NX)
head(Out) # creates a wide dataframe
```

```
##   days rain location vtype stream qrivlat Ts Tg Ttotal      s qcap
## 1    1  0.1      20 Grass      0      0 0 0      0 0.5048114    0
## 2    2  0.1      20 Grass      0      0 0 0      0 0.5096431    0
## 3    3  0.1      20 Grass      0      0 0 0      0 0.5144713    0
## 4    4  0.1      20 Grass      0      0 0 0      0 0.5192960    0
## 5    5  0.1      20 Grass      0      0 0 0      0 0.5241171    0
## 6    6  0.1      20 Grass      0      0 0 0      0 0.5289346    0
##   gwlevel gwrech leakage surfoff smloss stress
## 1      -6      0      0      1      1      1
## 2      -6      0      0      1      1      1
## 3      -6      0      0      1      1      1
## 4      -6      0      0      1      1      1
## 5      -6      0      0      1      1      1
## 6      -6      0      0      1      1      1
```

```
# put in dates rather than sequential numbers:
Out$Dates <- rep(Rain_uni$Dates[1:length(result[[1]])],
                ecohydro2d.options())$NX)
```

Now use `ggplot()` to create a figure by location

```
p <- ggplot(Out, aes(x=Dates, y=Ttotal)) + geom_line(aes(col="Total T"))
p <- p + facet_wrap(~location)
print(p)
```



2. using different vegetation at the sites