

Supplementary Information: Changes to original database

Abstract

This supplementary material file gives an overview of the changes that were made in the original data based on review of the original literature. Overall 36 data points were changed. The most common problem was a change in the sign of the forest change or the streamflow change.

This document list all the changes that were made to the original Zhang et al. [14] database after review of all the original papers.

A particular problem was that many catchments had the wrong sign for the change in forest cover. There are many catchments with reported positive change in cover and a large increase in flow. These were all checked and corrected if needed and a full list of all these changes is below:

- 76, Beaver Creek, the flow was corrected from 600% to 157% after review of the original publication [1].
- 124, D3, Amatya and Skaggs, 2008: The originally recorded 250% change by Zhang et al. [14] is clearly wrong. The paper says on page 7: Both of these outflow ratios (0.64 and 0.50) were higher than the calculated expected values of 0.55 for 2003 and 0.44 for 2005, respectively. So value should be $0.64/0.55 * 100 - 100$ or $0.5/0.44 * 100 - 100$: 16% or 13%. corrected to 16%
- 3, Baker Creek, Zhang and Wei, 2012. The original recorded 201.1% change by Zhang et al. [14] is also wrong. Original paper says on page 2031: Annual mean flow has been increased by 47.6%. corrected
- 67, April rd, which is incorrectly attributed to Ruprecht and Schofield (1991) in Zhang et al. [14]. This is actually from Ruprecht and Schofield (1989) and the original paper clearly indicates “clearfelling”. As a result the change in forest cover was changed to -100% rather than +100%.
- 210, March rd, 100, 147.6. Same problem as 67, Bari et al. (1996) clearly state that the catchment was cleared, so therefore the change in forest cover changed to -100%.
- 213, 214 and 215, Monda 1, 2 and 3. These catchments are tricky. The original paper [11] only reports on the control period and indicates that the catchments will be cleared. The later summary paper [13] shows the timeseries of the flow change, but does not report a single value, so the

values in the database must have been estimated from the timeseries. The further complication is that the treatment included clearing and reseedling and regrowth. This suggest that the records should be removed from the database, or only the first few years of the experiment used. In any case, if the values are kept, the sign of the change in forest cover needs to be changed to negative (Clearing).

- 230, Oleolega catchment. The paper describes a removal of forest up to 85%. changed Delta_F_perc to -85 from 90.
- 312, Yerraminup South. The original publication for this catchment is a Western Australian Water Authority report from 1987, which is hard to find, but we have added a copy in the “Papers” folder on github. In this report, in Table 2 on page 11, for the catchment a “Crown cover” decrease of 60% is given. Changed the sign of the change in forest cover: -60%.
- 72 Barratta, 100 Coachwood, 103 Corkwood, and 83 Bollygum, as cited by Cornish [8] and Cornish and Vertessy [9]. In the database from Zhang et al. [14], the forest change for all these catchments is positive. However, the paper highlights that these catchments were all logged and either naturally regenerated or were planted with a plantation species. So, similar to the the earlier mentioned Monda catchments, the reported change probably only refers to the first couple of years after clearing (before regrowth). In any case, the reported change in forest cover should be negative (clearing) rather than positive. Corrected for all three catchments.
- 78, Black Spur 1, the treatments and effects are only reported in a conference paper [10] and once again indicated clearing, meaning that the change in forest cover should be negative rather than positive (as reported in Zhang et al. [14]). Corrected. Similar to other paired watershed experiments, only the first couple of years can be linked to the effect as later regrowth cancels out part of the increase in flow.
- 104, Coshocton. Checking the original paper indicates that this is in fact a reduction in flow as a result of reforestation. Changed the sign of Delta_Q_f to be negative.
- 102, Cold Spring. Checking the original paper [12] indicates that this is in fact a reduction in flow as a result of reforestation. Changed the sign of Delta_Q_f to be negative.
- 85 Bosboukloof. This is essentially a duplicate of 184, but the cited paper analyses only 1 year of runoff after a major fire. In any case, the data should reflect a decrease in forest cover: changed the sign of Delta_f_perc to -80%.

- 259 Shackam Brook. There were a few issues with this catchment in the original database. The name was misspelled and it was incorrectly attributed to Brown et al. [6]. The original paper is the same as 102 [12]. Finally, the catchments were all reforestation as the title of the original report indicates and the reported streamflows are all decreases. Corrected Delta_Qf_perc to -20.7%.
- 95 Sage Brook. Similar to 259 and 102, originates from Schneider and Ayer [12]. Reforestation so Delta_Qf_perc corrected to -19.8%.
- 101 Coalburn. Original publication (Robinson, 1993) which is a symposium paper, is not available. The best summary of the research is in Birkinshaw et al. [3] which summaries 45 years of research in the Coalburn catchment. It was a reforestation experiment, and there was a decrease in the streamflow over the longer time period. Changed to -20.3%.

A further issue was the inclusion of the results of several catchments, for example from the study by Beck et al. [2], which had no significant change in flow. Despite this, the “average” change in flow was reported in the database. We don’t believe that this is correct and the results from such studies should be set to 0. A full list of changes is provided below:

- 97 Cibucio, 123 Culebrinas, 244 Portugues, 161 Grande de Loiza, 271 Tanama, 132 Fajardo, 89 Canovanas, 73, Bauta, 163 Grande de Patillas, 283 Valenciano, 181 Inabon, and 162 Grande de Manati. These are all catchments in Puerto Rico from the study from Beck et al. [2]. They should probably be removed from the database as the paper clearly indicates that there is no evidence of a change in flow due to reforestation. The values that are cited in the database should all be set to “not significant from 0”, so might be included as 0. Including them with positive or negative values is misleading. This study is a very detailed hydrological modelling study, but in the end finds no significant change in streamflow as a result of deforestation. Values for all 12 studied catchments set to 0 in the database.
- 188 Kimakia. and 254 Sambret. The data in the database from Zhang et al. [14] appear to originate from Bruijnzeel et al. [7] which gives 3 values for different lengths of studies. However, the values in the original study by Blackie [5] and Blackie [4] do not seem to add up to the same values, and the specific values are not mentioned in the actual papers. In addition, as Bruijnzeel et al. [7] mentions in the footnotes, the control for Kimakia is a bamboo catchment, while the control for Sambret is a tea plantation. Overall, this suggests that the data are probably not a clear deforestation/reforestation study and should be discarded from the analysis.
- 221 N. Creek, Babinda, Queensland. The original paper from this study highlights that the differences between the catchments were insignificant.

128 References

- 129 [1] Malchus B. Baker Jr. Changes in streamflow in an herbicide-treated
130 pinyon-juniper watershed in arizona. *Water Resources Research*, 20
131 (11):1639–1642, 1984. ISSN 0043-1397. doi: [https://doi.org/10.1029/](https://doi.org/10.1029/WR020i011p01639)
132 WR020i011p01639. URL [https://agupubs.onlinelibrary.wiley.com/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/WR020i011p01639)
133 doi/abs/10.1029/WR020i011p01639.
- 134 [2] H. E. Beck, L. A. Bruijnzeel, A. I. J. M. van Dijk, T. R. McVicar, F. N.
135 Scatena, and J. Schellekens. The impact of forest regeneration on stream-
136 flow in 12 mesoscale humid tropical catchments. *Hydrol. Earth Syst. Sci.*,
137 17(7):2613–2635, 2013. ISSN 1607-7938. doi: 10.5194/hess-17-2613-2013.
138 URL <https://hess.copernicus.org/articles/17/2613/2013/>. HESS.
- 139 [3] Stephen J. Birkinshaw, James C. Bathurst, and Mark Robinson. 45
140 years of non-stationary hydrology over a forest plantation growth cycle,
141 coalburn catchment, northern england. *Journal of Hydrology*, 519:559–
142 573, 2014. ISSN 0022-1694. doi: [https://doi.org/10.1016/j.jhydrol.2014.](https://doi.org/10.1016/j.jhydrol.2014.07.050)
143 07.050. URL [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0022169414005848)
144 S0022169414005848.
- 145 [4] JR Blackie. 2.2. 1 the water balance of the kericho catchments. *East African*
146 *Agricultural and Forestry Journal*, 43(sup1):55–84, 1979.
- 147 [5] JR Blackie. 3.2. 1 the water balance of the kimakia catchments. *East*
148 *African Agricultural and Forestry Journal*, 43(sup1):155–174, 1979.
- 149 [6] Alice E. Brown, Lu Zhang, Thomas A. McMahon, Andrew W.
150 Western, and Robert A. Vertessy. A review of paired catch-
151 ment studies for determining changes in water yield resulting from
152 alterations in vegetation. *Journal of Hydrology*, 310(1-4):28–61,
153 2005. URL [http://www.sciencedirect.com/science/article/B6V6C-](http://www.sciencedirect.com/science/article/B6V6C-4G05MM9-1/2/bbc5fc0e958a8f34bcb7c1cc7fa57b48)
154 4G05MM9-1/2/bbc5fc0e958a8f34bcb7c1cc7fa57b48.
- 155 [7] Leendert Adriaan Bruijnzeel et al. Hydrology of moist tropical forests
156 and effects of conversion: a state of knowledge review. *Hydrology of moist*
157 *tropical forests and effects of conversion: a state of knowledge review.*, 1990.
- 158 [8] P. M. Cornish. The effects of logging and forest regeneration
159 on water yields in a moist eucalypt forest in new south wales,
160 australia. *Journal of Hydrology*, 150(2-4):301–322, 1993. URL
161 [http://www.sciencedirect.com/science/article/B6V6C-487D3Y2-](http://www.sciencedirect.com/science/article/B6V6C-487D3Y2-9J/2/73c981ba76284d9d629f6b221d6fd6c6)
162 9J/2/73c981ba76284d9d629f6b221d6fd6c6.
- 163 [9] P. M. Cornish and R. A. Vertessy. Forest age-induced changes in evapo-
164 transpiration and water yield in a eucalypt forest. *Journal of Hydrology*,
165 242(1-2):43–63, 2001. URL [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/B6V6C-429910G-3/2/0158b1f89ff436f338a9e688a47f06c4)
166 article/B6V6C-429910G-3/2/0158b1f89ff436f338a9e688a47f06c4.

- 167 [10] M. D. A. Jayasuriya and P. J. O'Shaughnessy. *The Use of*
168 *Mathematical Models in Evaluating Forest Treatment Effects on*
169 *Streamflow*, pages 135–139. Hydrology and Water Resources Sym-
170 posium, 1988. doi: 10.3316/informit.692214289455295. URL
171 [https://search-informit-org.ezproxy.library.sydney.edu.](https://search-informit-org.ezproxy.library.sydney.edu.au/doi/10.3316/informit.692214289455295)
172 [au/doi/10.3316/informit.692214289455295](https://search-informit-org.ezproxy.library.sydney.edu.au/doi/10.3316/informit.692214289455295). doi: 10.3316/infor-
173 mit.692214289455295.
- 174 [11] P. J. O'Shaughnessy, K. J. Langford, H. P. Duncan, and R. J. Moran.
175 Catchment experiments in mountain ash forests at north maroondah. *Aus-*
176 *tralian Forestry*, 42(3):150–160, 1979. ISSN 0004-9158. doi: 10.1080/
177 00049158.1979.10674220. URL [https://doi.org/10.1080/00049158.](https://doi.org/10.1080/00049158.1979.10674220)
178 [1979.10674220](https://doi.org/10.1080/00049158.1979.10674220). doi: 10.1080/00049158.1979.10674220.
- 179 [12] William Joseph Schneider and Gordon Roundy Ayer. Effect of reforestation
180 on streamflow in central new york. Report 1602, 1961. URL [http://pubs.](http://pubs.er.usgs.gov/publication/wsp1602)
181 [er.usgs.gov/publication/wsp1602](http://pubs.er.usgs.gov/publication/wsp1602).
- 182 [13] Fred Watson, Rob Vertessy, Tom McMahon, Bruce Rhodes, and Ian
183 Watson. Improved methods to assess water yield changes from paired-
184 catchment studies: application to the maroondah catchments. *For-*
185 *est Ecology and Management*, 143(1):189–204, 2001. ISSN 0378-1127.
186 doi: [https://doi.org/10.1016/S0378-1127\(00\)00517-X](https://doi.org/10.1016/S0378-1127(00)00517-X). URL [https://www.](https://www.sciencedirect.com/science/article/pii/S037811270000517X)
187 [sciencedirect.com/science/article/pii/S037811270000517X](https://www.sciencedirect.com/science/article/pii/S037811270000517X).
- 188 [14] Mingfang Zhang, Ning Liu, Richard Harper, Qiang Li, Kuan Liu, Xiao-
189 hua Wei, Dingyuan Ning, Yiping Hou, and Shirong Liu. A global re-
190 view on hydrological responses to forest change across multiple spatial
191 scales: Importance of scale, climate, forest type and hydrological regime.
192 *Journal of Hydrology*, 546:44–59, 2017. ISSN 0022-1694. doi: [https://](https://doi.org/10.1016/j.jhydrol.2016.12.040)
193 doi.org/10.1016/j.jhydrol.2016.12.040. URL [http://www.sciencedirect.](http://www.sciencedirect.com/science/article/pii/S0022169416308307)
194 [com/science/article/pii/S0022169416308307](http://www.sciencedirect.com/science/article/pii/S0022169416308307).