

Factors determining how catchments respond to forest cover change. Re-analysing global data sets.

R. Willem Vervoort^{*,a,1}, Eliana Nervi^c, Jimena Alonso^d

^aARC Training Centre Data Analytics for Resources and the Environment

^bSchool of Life and Environmental Sciences, The University of Sydney, Sydney, NSW 2006, Australia

^cProject Manager, FPTA 357, Instituto Nacional de Investigacion Agropecuaria, INIA-Uruguay, Ruta 48 km 10, Rincon del Colorado, 90100 Canelones, Uruguay

^dInstitute of Fluid Mechanics and Environmental Engineering, School of Engineering, Universidad de la República, 11200 Montevideo, Uruguay

Abstract

Changes in forest cover cause changes in streamflow, however this change is different between deforestation and reforestation, and strongly affected by climate, with drier climates indicating larger changes in streamflow. Deforestation causes a 32% greater change in flow compared to reforestation. Area of the catchment only affects the change in streamflow after log transformation, given the wide variety in the data from small scale paired watershed studies. Smaller studies dominate the database with 42% of the data < 1 km² and 65% of the data < 10 km². Length of the study and initial year of the study did not affect the change in flow, in contrast to other reported studies. Despite these findings, overall explained variance (38%) is low due the quality of the inputs and additional unknown confounding factors. Three recent papers review and analyse large global datasets related to impacts of forest cover on streamflow. Using three different approaches, they all find a strong relationship between forestation/deforestation and streamflow. However, the past approaches in the literature are variable and can be substantially improved in statistical rigour, and indicate different confounding factors on the impact of forestation. The data for these three papers were reviewed, combined and re-analysed to answer the following new and older questions: 1) How is streamflow impacted by the change in forest cover as a function of catchment area; 2) how is this relationship conditioned by the length of the study, and climate; and 3) are there other possible variables that impact the observed change in streamflow? Generalised additive models were used to run flexible regressions including multiple variables.

1. Introduction

There has been an long and on-going discussion in the hydrological literature around the impact of forests on streamflow (Andréassian, 2004; Brown et al., 2013, 2005; Filoso et al., 2017; Jackson et al., 2005; Zhang et al., 2017). The historic work highlights a general consensus that if forest areas increase, streamflow decreases and vice-versa. The most dramatic result in relation to this, is

*Corresponding Author
Preprint submitted to *Journal of Hydrology*
Email addresses: willem.vervoort@sydney.edu.au (R. Willem Vervoort), February 4, 2022
eliananervi@gmail.com (Eliana Nervi), jalonso@fing.edu.uy (Jimena Alonso)

18 Figure 5 in Zhang et al. (2011) indicating (for Australian catchments) a 100%
19 decrease in streamflow for catchments with 100% forest cover. However, on the
20 other end of the spectrum, for three French catchments (Cosandey et al., 2005),
21 there was no change in streamflow characteristics in two of the catchments after
22 deforestation.

23 For the purpose of this paper, *watershed* and *catchment* are interchangeable
24 terms. Many of the US studies use *watershed*, while European and Australian
25 studies use *catchment*. In particular, we retained the term “paired watershed
26 studies” as this is the most common terminology, but further mostly use the
27 term catchment.

28 Several review papers have summarized the plethora of forestation and defor-
29 estation studies across the globe, in relation to paired watershed studies (Bosch
30 and Hewlett, 1982; Brown et al., 2005), related to reforestation in particular
31 (Filoso et al., 2017), and more generally (Jackson et al., 2005; Zhang et al.,
32 2017). These studies aim to generalize the individual findings and to identify
33 if there are global trends or relationships that can be developed. The most
34 recent reviews (Filoso et al., 2017; Zhang et al., 2017) developed an impressive
35 global database of catchment studies in relation to changes in streamflow due
36 to changes in forest cover. The Zhang et al. (2017) dataset, which covers over
37 312 studies, is described in terms of the change in streamflow as a result of
38 the change in forest cover, where studies related to both forestation (increase
39 in forest cover) and deforestation (decrease in forest cover) were included. In
40 contrast, the paper by Filoso et al. (2017) focused primarily on reforestation,
41 and covered an equally impressive database of 167 studies using a systematic
42 review. In this case the collected data is mostly coded as count data and only a
43 subset of 37 studies was analysed for actual water yield change. There is some
44 overlap between the two data sets, but there are also some studies unique to
45 both sets.

46 The conclusions of the first paper (Zhang et al., 2017) suggest that there is
47 a distinct difference in the change in flow as a result of forestation or deforesta-
48 tion between small watersheds (catchments), defined as $< 1000 \text{ km}^2$ and large
49 watersheds (catchments) $> 1000 \text{ km}^2$. While for small catchments there was no
50 real change in runoff with changes in cover, for large catchments there was a
51 clear trend showing a decrease in runoff with and increase in forest cover. Their
52 main conclusion was that the response in annual runoff to forest cover was scale
53 dependent and appeared to be more sensitive to forest cover change in water
54 limited catchments relative to energy limited catchments (Zhang et al., 2017).

55 The second study (Filoso et al., 2017) was a systematic review which clas-
56 sified the historical research and highlighted gaps in the spatial distribution,
57 the types of studies and the types of analysis. Their main conclusion was also
58 that reforestation decreases streamflow, but that there were many interacting
59 factors. For a subset of quantitative data (37) they showed a log-linear relation-
60 ship between decreasing catchment size and an increasing decline in streamflow.
61 In addition, they identified that studies with shorter periods of data collection
62 resulted in larger declines in streamflow.

63 A final summary paper that includes much of the same data as Zhang et

al. (2017) and Filoso et al. (2017) is Zhou et al. (2015), which has one author in common with Zhang et al. (2017). However, this paper aims to explain the variation in the data using the Fuh model, and in particular aims to link the variation in the observed data to variations in the exponent m in the model. A key observation is that in drier environments, the effects of deforestation are much greater than in wetter environments, which is also suggested by Figure 4 in Zhang et al. (2017).

Encouraged by the work presented by Zhang et al. (2017), Filoso et al. (2017) and Zhou et al. (2015) and the large database of studies presented by these authors, we believe more can be done to add to this important discussion. In this paper, the aim extend the analysis of the collected data and to expand and combine the data sets.

In particular, the main method in the work by Zhang et al. (2017) is a single covariate linear regression, and in Filoso et al. (2017) the focus is mainly on classification and there is again some single covariate linear regression. As Zhang et al. (2017) points out, a main assumption in their work is that the catchment size threshold at 1000 km² is a distinct separation between “small” and “large” catchments. However, the subset of 37 data points in Filoso et al. (2017) (their Figure 9) does not appear to support this, suggesting a continuum. And while the work Filoso et al. (2017) provides important insights in study types, analysis types, forest types and broad classification, there is limited quantification of actual impact, and focussed only on forest cover increase and did not deal with forest cover removal.

As a result the objective of this paper is to 1) enhance the data set from Zhang et al. (2017) with further catchments (such as from Filoso et al. (2017)) and spatial coordinates and 2) to analyse the possibility of non-linear and confounding partial effects of the different factors and variables in the data using generalised linear (GLM) and generalised additive models (GAM Wood (2006)).

Building on the analyses by Zhang et al. (2017) and Filoso et al. (2017), and combining their conclusions, the main hypothesis to test is that the change in streamflow is impacted by the change in forest cover. However, this change is potentially modulated by the area under consideration (affecting the length of the flowpaths Zhou et al. (2015)), the length of the study (c.f. Jackson et al. (2005); Filoso et al. (2017)) and the climate (as indicated by either E0/Pa or latitude and longitude Filoso et al. (2017); Zhou et al. (2015)).

However, there could be further confounding factors, which are eluded to by Filoso et al. (2017):

- the type of analysis, i.e. paired watershed studies, modelling, time series analysis etc.
- the age of the study, assuming that historical studies might not have had the ability to measure at the accuracy that currently is available to researchers, or that more careful historical attention to detail in field studies might have been lost more recently due to reductions in research investment.

Finally, this work aims to point to further research that can expand this area of work, based on the collected data, to better understand the impact of forest cover change on streamflow.

2. Methods

2.1. The original data sets

The starting point of this paper is the data base of studies which were included in Zhang et al. (2017) as supplementary material. The columns in this data set are the catchment number, the catchment name, the Area in km², the annual average precipitation (Pa) in mm, the forest type, hydrological regime, and climate type, the change in forest cover in % ($\Delta F\%$) and the change in streamflow in % ($\Delta Qf\%$), based on equation 1 in Zhang et al. (2017)), the precipitation data type, the assessment technique, and the source of the info, which is a citation. Several of these columns contain abbreviations to describe the different variables, which are summarised in Table 1.

Table 1 Summary of abbreviations of factors used in the Zhang et al. (2017) data set

Factor	Abbreviation	Definition
forest type	CF	coniferous forest
	BF	broadleaf forest
	MF	mixed forest
hydrological regime	RD	rain dominated
	SD	snow dominated
climate type	EL	energy limited
	WL	water limited
	EQ	equitant
precipitation data type	OB	observed
	SG	spatial gridded
	MD	modelled
assessment technique	PWE	paired watershed experiment
	QPW	quasi-paired watershed experiment
	HM	hydrological modelling
	EA	elasticity analysis
	SH	combined use of statistical methods and hydrographs

While Zhang et al. (2017) use the dryness index in their analysis, and calculate the variable climate type from this index, the potential or reference evapotranspiration was not originally included as part of the published data set. In this paper, we used only the dryness index and did not use the climate type as a variable (as they are interchangeable). We combined the tables for small catchments (< 1000 km²) and large catchments (>= 1000 km²) in our analysis.

2.2. Additional data collection

To enhance the existing data set, this study added additional variables and cross-checked the studies with the data set from Filoso et al. (2017). In particular, we focussed on the 37 data points included in the quantitative analysis in Filoso et al. (2017).

In addition, additional variables added were the latitude and longitude for the center of the catchment as an approximation of its spatial location. Using this information reference evapotranspiration (E_0) was extracted from the Global Aridity Index and Potential Evapo-Transpiration (ET_0) Climate Databasev2 (Trabucco and Zomer, 2018), if a value of E_0 was not available from the original papers. For large catchments, this value, similar to annual average rainfall, is only an approximation of the climate at the location.

Similar to Zhang et al. (2017), the “dryness index” was calculated from the reference evapotranspiration and the annual average rainfall (Pa) as:

$$D = \frac{E_0}{Pa} \quad (1)$$

The length of the study can be a variable influencing the change in flow (Filoso et al., 2017; e.g. Jackson et al., 2005), as for example, more mature plantations are thought to have smaller impacts on flow or regrowth might follow a “Kuczera curve” (Kuczera, 1987). Therefore, the length of the study calculate as the difference between the starting data and completion date of the different studies was extracted from the references provided by Zhang et al. (2017). The length of the study was already included in the data from Filoso et al. (2017), but these were checked against the original publications.

Several additional data points from catchment studies were extracted from Zhang et al. (2011), Zhao et al. (2010), Borg et al. (1988), Thornton et al. (2007), Zhou et al. (2010), Rodriguez et al. (2010), Ruprecht et al. (1991) and Peña-Arancibia et al. (2012), and these were checked against the existing studies to prevent overlap. In the citation column in the data set, in general the main reference for the calculated change in streamflow was used, because sometimes the original study did not provide the quantification of the change in streamflow (i.e. Table 6 in Zhang et al. (2011)). We also removed one data point from the analysis, which corresponds to catchment #1 (Amazon) in Zhang et al. (2017). This is because the cited reference (Roche, 1981) only relates to 1 and 1.5 ha paired catchment studies in French Guyana, and in which the actual change in forest cover is not recorded. Furthermore, the change in flow for catchment #76 was corrected from 600% to 157% after review of the original publication (Baker Jr., 1984). Finally, on review of all the data in Zhang et al. (2017) and Filoso et al. (2017), 29 potential duplicates were identified and flagged in the data.

The final column in the improved data set is a “notes” column, which is not further used in the analysis, but gives context to some of the data for future research and highlights some of the discrepancies that we found between the original papers and the data in the tables from Zhang et al. (2017).

172 2.3. Statistical modelling

173 To estimate how the change in streamflow is affected by the change in forest
174 cover while considering the effects of the other variables, we applied generalised
175 additive modelling (GAM) (Wood, 2006).

176 The general model tested is:

$$\Delta Qf\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + \sum X_i + \sum s(Z_i) + \varepsilon \quad (2)$$

177 Here X_i are factorial variables, while Z_i are continuous variables. The model
178 assumes no direct interactions and all variables are additive. The changes in
179 forest cover contain both positive (forestation) and negative values (deforesta-
180 tion). In Zhang et al. (2017), these changes were jointly analysed, assuming the
181 effect on the change in flow was linear and the effect if removing forest cover
182 was the same as an equivalent reforestation. However, the impact of an increase
183 in forest cover can be different from the same fractional decrease in forest cover.
184 Therefore all the change in forest cover data is converted to positive values, and
185 an additional column ($sign_{forestcover}$) is added that indicates whether it was a
186 forest cover increase or decrease.

187 A further assumption in the model is that all continuous variables Z_i (such
188 as annual precipitation (Pa)) can have a linear or non-linear relationship with
189 $\Delta Qf\%$. This means that a smooth function $s()$ is applied to the Z_i variables.
190 For the smoothing function we applied thin plate regression splines with an
191 additional shrinkage penalty which means the terms can be shrunk to 0 if not
192 significant (Wood, 2006).

193 For the model in equation 2, we initially only used the data from Zhang et
194 al. (2017) to make sure that the additional catchments added to the data set
195 did not influence the results. Subsequently the analysis was repeated and the
196 additionally identified catchments were added.

197 More generally the results were analysed to identify:

- 198 1. the significance of the different variables
- 199 2. the direction of the categorical or shape of the smooth variables

200 3. Results

201 3.1. Description of the data

202 The overall dataset contains 330 observations of changes in flow, which in-
203 cludes the newly identified data sets and after removing identified duplicate
204 data and lines with missing data. In contrast, the original dataset from Zhang
205 et al. (2017) contained 312 catchments and the Filoso et al. (2017) study used
206 37 catchments (Table S2 in Filoso et al. (2017)). The overall distribution of
207 changes in flow is highly skewed as is the distribution of changes in forest cover
208 and $Area\ km^2$. The values of changes in flow greater than 100% and smaller
209 than -100% clearly create long tails on the change in flow distribution. Note also




Figure 1: Overview of the distribution of the data set for five of the included variables.

the large number of studies with 100% forest cover reduction. Smaller catchments dominate the database with 42% of the data from catchments $< 1 \text{ km}^2$ and 65% of the data for catchments $< 10 \text{ km}^2$ (Figure 1).

In addition, this shows that for the data related to forest decreases, there is almost always a positive flow change (Figure 2). In other words, flow almost always increased. However, for increases in forest cover, this is not the case, and flow can both increase and decrease. However in both cases the variability in the reported change in flow increases with the increase in forest cover change.

3.2. The general relationship between change in forest cover and streamflow

Following Zhang et al. (2017), the first step is to investigate the percent change in flow as a linear effect of the percent change forestry and modulated by the direction of the change, either an increase in forest cover, or decrease in forest cover:

$$\Delta Qf\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + \varepsilon \quad (3)$$




Figure 2: Changes in flow as a function of increases and decreases in forest cover

Table 2: Summary results of the first regression model predicting change in streamflow from change in forest cover and accounting for the direction of the change. The first three rows relate to the model using the original data base from Zhang et al. (2017). The bottom three rows are the results of the model including the new data.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	8.65	5.56	1.56	0.12
DeltaF_perc_pos	0.45	0.09	5.26	0
Forest_Signincrease	-29.17	5.79	-5.04	0
(Intercept)	10.76	5.34	2.02	0.04
DeltaF_perc_pos	0.41	0.08	5.04	0
Forest_Signincrease	-38.03	5.26	-7.23	0

223 The overall variance explained in this model (equation (3)) is not high with
 224 an adjusted r^2 of 0.23, it generally supports the hypothesized relationship be-
 225 tween the change in forest cover and the change in flow. The model suggests
 226 that for every 1% change in forest cover, on the average, the flow changes 0.45%.
 227 However the change in flow is different for forest cover decreases compared to
 228 forest cover increases. In fact, forest cover increases decrease flow by 29% less
 229 than a similar decrease in forest cover causes flow to increase. So roughly speak-
 230 ing, a 1% forest cover increase on the average decreases flow by $(1 - 0.29) * 0.45\%$,




Figure 3: Residual plots for the first simple regression model indicating a slightly fat-tailed residual distribution

while a the percentage forest cover decrease will increase flow by 0.45%.

Of importance here is to highlight the residuals of this regression (equation (3) and Figure 3). These are approximately normal, although there is still significant skew on the upper and lower parts of the distribution (Figure 3). In other words, the distribution of the residuals is somewhat fat-tailed. We will discuss this later.

Including the data from some of the newly identified studies indicates that this mainly strengthens the difference between the forest cover increases and decreases (Table 2), and the result indicate a reduction in the mean decrease in flow as a result of forest cover change if the new data is included.

It is however it is clear from the lack of explaining power for the model, that there could be confounding factors, as alluded to in the methods. The obvious ones being catchment dryness and area (following Zhang et al. (2017)), which we will analyse later.

3.3. The effect of location on the globe

As indicated, an initial hypothesis relates to whether there is a strong spatial global gradient as captured by latitude and longitude. These data were added for the different studies, mostly by using the data reported by the authors, but in some cases approximating the location of the centre of the catchment using Google Maps. In the dataset, an additional column is added to indicate the source of the location data. As the global map (Figure 4) shows, the distribution of case study catchments covers multiple continents and shows some distinct




Figure 4: Distribution of included catchments across the globe based on reported or estimated latitude and longitude

253 clustering in parts of the world. Of interest is whether the spatial clustering
 254 also indicates a difference in response to forest cover change:

$$\Delta Qf\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + Latitude + Longitude + \varepsilon \quad (4)$$

Table 3: Results of the model including Latitude and Longitude including new data

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	10.99	5.75	1.91	0.06
DeltaF_perc_pos	0.42	0.08	5.13	0
Forest_Signincrease	-38.99	5.54	-7.04	0
Latitude	-0.01	0.09	-0.13	0.9
Longitude	0.01	0.03	0.26	0.79

255 There appears to be no significant gradient in either latitude or longitude
 256 (Table 3), suggesting that the distribution of the catchments across the globe
 257 has little influence on the overall result. The total explaining power of the model
 258 is still low with an adjusted r^2 of 0.23 suggesting further factors influencing the
 259 change in streamflow that are currently not included in the model.

260 3.4. Impact of climate

261 Climate, and in particular evapotranspiration can have a significant effect
 262 on the streamflow change as represented by the dryness index, which is also
 263 highlighted by both Zhang et al. (2017) and Jackson et al. (2005). Increased
 264 evapotranspiration could lead to drier catchments, unless balanced by rainfall

265 (such as possibly in the tropics). Initially, we tested models using annual average
 266 precipitation (Pa (mm)), but because of the interactions between precipitation,
 267 evapotranspiration and the dryness index, we concentrated on the dryness index
 268 as the key variable. Given that Latitude and Longitude were not significant, we
 269 dropped these from the model.

$$\Delta Qf\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + Dryness + \varepsilon \quad (5)$$

Table 4: Results of the model including the dryness index

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	8.83	6.3	1.4	0.16
DeltaF_perc_pos	0.4	0.08	4.88	0
Forest_Signincrease	-38.26	5.32	-7.19	0
Dryness	1.7	2.67	0.64	0.52

270 The results from this model (equation (5) and Table 4) interestingly indicate
 271 no impact of dryness modulating the change in streamflow as a function of the
 272 change in forest cover change, which is surprising in light of earlier reported
 273 results (Filoso et al., 2017; Zhang et al., 2017). In this case, the evidence is
 274 highly doubtful ($p = 0.52$). However, it is very well possible that there is a
 275 further interaction in the data with other variables or unknown variables that
 276 this simpler version of the model cannot identify. This is partly evidenced by
 277 the fact that the overall variance explained is still low, with an adjusted r^2 of
 278 0.22.

Table 5: catchments for which the dryness index > 4

Latitude	Longitude	Watershed name
34.67	-111.7	Beaver Creek, AZ #3-2
36.4	-120.4	Cantua
32.74	-111.5	Natural Drainages, Ariz., U.S.A, C
32.74	-111.5	Natural Drainages, Ariz., U.S.A, A
34.43	-112.3	White Spar, Ariz., U.S.A, B
-25.75	28.23	Queens river

279 There are also possible issues with the data, as a few of the catchments have
 280 Dryness values that are very large (> 4) and these values have high leverage
 281 in the data, affecting the residual distribution. These catchments are listed in
 282 Table 5.

283 3.5. Is there a distinct effect of area?

284 The major hypothesis to test is the effect of area on the change in flow,
 285 following the analysis by Zhang et al. (2017) and Filoso et al. (2017). Given
 286 the highly skewed distribution of the catchment areas (Figure 1), a log base 10
 287 transformation was applied to the variable *Area* (km^2).

$$\Delta Q f\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + \log_{10}(Area\ (km^2)) + Dryness + \varepsilon \quad (6)$$

Table 6: Results of the model including Area and the dryness index

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	15.37	7.24	2.12	0.03
DeltaF_perc_pos	0.3	0.1	2.94	0
Forest_Signincrease	-36.61	5.38	-6.8	0
Dryness	2.48	2.69	0.92	0.36
log10(Area_km2)	-3.03	1.67	-1.81	0.07

288 The results of this model (Equation (6)) indicate there is at least some
 289 evidence ($p = 0.07$) that there is a reduction in the effect of forest cover change
 290 on streamflow related to Area (km^2) (Table 6). In fact, the results suggests that
 291 for every additional 10 km^2 in catchment size the mean change in flow reduces by
 292 3%. Another interesting fact to note is that with the inclusion of Area (km^2) as
 293 a variable in the model, the effect of Dryness becomes slightly more important,
 294 possibly suggesting an interaction between Dryness and Area. Including this
 295 interaction in the model (Table 7) results in the increased evidence ($p = 0$) that
 296 Dryness affects the change in flow caused by changes in forest cover and that
 297 the effect of Area is only important ($p = 0.95$)) as an interaction with Dryness.

Table 7: Results of the model including an interaction between Area and the dryness index

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	11.95	7.43	1.61	0.11
DeltaF_perc_pos	0.29	0.1	2.85	0
Forest_Signincrease	-34.62	5.46	-6.34	0
Dryness	5.46	3.1	1.76	0.08
log10(Area_km2)	0.15	2.35	0.06	0.95
Dryness:log10(Area_km2)	-2.62	1.37	-1.91	0.06

298 3.6. Are some of the variables possibly non-linear?

299 The work by Filoso et al. (2017) and earlier by Jackson et al. (2005) has
300 indicated that the length of the study might influence the response. This links
301 to the idea from Kuczera (1987) that the effect of logging or deforestation or
302 reforestation reduces with the length of time post intervention (see also Jackson
303 et al. (2005)). In addition to adding *length* (being the difference between the
304 reported start date and end date of data collection in the specific study) as a
305 variable, three other continuous variables (*Dryness*, *Area*, *From*) were consid-
306 ered non-linear in this model. As a result a shrinkage smoothing spline (Wood,
307 2006) was applied to these variables. *From* represents the starting date of the
308 data collection.

$$\begin{aligned} \Delta Qf\% \sim \Delta\%forest\ cover_{positive} + sign_{forest\ cover} + \\ s(\log10(Area\ (km^2))) + s(length) + s(Dryness) + \\ s(From) + \varepsilon \end{aligned} \quad \begin{matrix} (7) \\ (8) \end{matrix}$$

Table 8: Statistical summary for the linear terms in the model with non-linear terms

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	13.99	5.68	2.46	0.01
DeltaF_perc_pos	0.33	0.1	3.51	0
Forest_Signincrease	-37.18	5.48	-6.78	0

Table 9: Statistical summary for the smooth terms in the model with non-linear terms

	edf	Ref.df	F	p-value
s(log10(Area_km2))	0.68	9	0.28	0.05
s(Dryness)	4.28	9	0.94	0.07
s(length)	0	9	0	0.55
s(From)	4.62	9	0.9	0.1

309 Including non-linearity (Equation (8)) increases the overall explaining power
310 of the model to an adjusted r^2 of 0.26 and deviance explained of 0.29, but cre-
311 ates a few changes in the significance of the variables (Table 9). For example,
312 the smoothed variable for Area (km²) (p = 4.62)) and Dryness (p = 9))still
313 provides reasonable explanation of changes in stream flow. However *length* does
314 not explain any of the variation. In contrast, *From*, which indicates the start
315 date of the study (and therefore the age of the study) has an effect (p = 0.1)).

316 However, it also increases the chance of over fitting, as the smoothing splines al-
317 low significant flexibility, which will be investigated later. Including interactions
318 into the smooths is possible, but the results are even more difficult to interpret
319 given the high flexibility of the two-dimensional smooth.

320 We now also include the remaining categorical variables (Precipitation data
321 type, Assessment technique, Forest type and Hydrological regime) i.e. Equation
322 (2).

Table 10: Statistical summary for the linear terms the full model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-29.23	18.18	-1.61	0.11
DeltaF_perc_pos	0.31	0.09	3.29	0
Forest_Signincrease	-23.48	6.84	-3.43	0
Precip_data_typeOB	-5.03	13.94	-0.36	0.72
Precip_data_typeSG	19.41	15.38	1.26	0.21
Assessment_techniqueEA,	19.56	45.02	0.43	0.66
HM				
Assessment_techniqueHM	33.57	12.36	2.72	0.01
Assessment_techniquePWE	52.79	11.72	4.5	0
Assessment_techniquePWE,	43.58	45.93	0.95	0.34
HM				
Assessment_techniqueQPW	39.98	21.36	1.87	0.06
Assessment_techniqueQPW,	31.36	27.12	1.16	0.25
EA				
Assessment_techniqueSH	42.63	12.41	3.44	0
Forest_typeCF	-2.33	8.23	-0.28	0.78
Forest_typeMF	-10.36	8	-1.3	0.2
Hydrological_regimeSD	6.99	10.25	0.68	0.5

Table 11: Statistical summary for the smooth terms for the full model

	edf	Ref.df	F	p-value
s(log10(Area_km2))	0	9	0	0.37
s(Dryness)	3.96	9	1.68	0
s(length)	0	9	0	0.36
s(From)	8.8	9	3.64	0

323 This model explains more of the variance, but the improvement is marginal
324 compared to the previous model with a adjusted r^2 of 0.34. This indicates that
325 the categorical variables explain a limited amount of the variance. However,
326 interesting to note from Table 10 that several of the assessment methods are




Figure 5: Visualisation of the smooth variables in the model

327 significant. In particular Paired Watersheds experiments (PWE), Hydrological
 328 modelling (HM) and Statistical techniques (SH) are strongly significant ($p <$
 329 0.05). In this case, $\log_{10}(\text{Area } (km^2))$ is no longer a significant predictor, the
 330 reasons for this will be discussed later.

331 Figure 5 highlights that the relationship between $\log_{10}(\text{Area } km^2)$ and the
 332 change in flow is essentially linear, not significant, and does not need to be
 333 smoothed (this is the reason why using penalized smooths following Wood (2006)
 334 is useful). It still has a negative slope, indicating that in larger catchments
 335 the impact of changes in forest cover on streamflow is less than for smaller
 336 catchments. Similarly, the *length* variable is not significant. However, both
 337 *From* and *Dryness* variables show strong non-linearity, but this does not show
 338 a clear trend due to the scatter and the distributions of the data. For example,
 339 in the case of *From* the relationship is clearly strongly influenced by the two
 340 data points before 1920. Similarly, *length* and *Dryness* have points with high
 341 leverage.

Table 12: Statistical summary of the smooth terms reducing dataset to studies with the study length shorter than 60 years, Dryness > 4 and studies commencing after 1920

	edf	Ref.df	F	p-value
s(Dryness)	3.21	9	3.22	0
s(log10(Area_km2))	0.46	9	0.11	0.13

	edf	Ref.df	F	p-value
s(length)	0	9	0	0.44
s(From)	4.06	9	1.12	0.05

The flexible nature of the splines means that the length variable captures some substantial variation in the data, but it is unclear what exactly is captured. The shape of the conditional response (Figure 5) also does not reflect the type of response highlighted in Filoso et al. (2017) and Jackson et al. (2005). One reason could be the few data points with very long data series, and very old studies (before 1920 essentially), and highly variable responses (Figure 5). Therefore it can be important to investigate what removing these factors has on the overall model and the significance of the variables. The next model therefore removes the following data: *Dryness* > 4, *length* > 60 years, *From* > 1920. This result in a reduction of the data set from 330 to 304 catchments.

The model has more explaining power with an adjusted r^2 of 0.34. This results in *Dryness* now shows a clear non-linear response with drier catchments having a greater impact of changes in forest cover on streamflow (Figure 6). However the other 3 variables are no longer significant due to the scatter in the data, but *length* and *From* show some strong local non-linear behaviour.

4. Discussion

4.1. Catchment size

Essentially, the analysis shows that there is no clear effect of catchment size (Figure 6), and, in contrast to Zhang et al. (2017), there is no evidence a distinct threshold in the size of the catchment that influences the change in the streamflow as a result of changes in forestry. If anything the scatter in the data (in the change in flow) is greater for the smaller catchments than for the larger catchments (Figure 7). In other words, the response to changes in forest cover is more consistent for larger catchments than it is for smaller catchments.

An explanation for the catchment size effect might be that large catchments have more storage and longer flow paths and therefore have more opportunity to buffer the effects of forest cover change (Navas et al., 2019). Therefore, specifically if the forest cover change is small relative to the catchment size, the effect of this change will be buffered.

There are two caveats on this explanation. The first is that there is a distinct trend in the data between Δ Forest cover and $\log_{10}(\text{Area (km}^2\text{)})$ (linear regression indicates an adjusted r^2 of 0.36 with a slope of -9.76) indicating that for every 10 km² increase in catchment size on the average the recorded forest cover change is approximately 10% lower. This is basically a result of the fact that large changes in forest cover in larger catchments are difficult to “implement” in an experiment.

This is also reflected in the second caveat most of the smaller catchments are “real observed data” using paired watershed studies, while for larger catchments, the analysis are mostly based on modelling approximations using either




Figure 6: Visualisation of the smooth variables in the model with reduced data for dryness and length




Figure 7: Overview of the data highlighting the dominance of small catchment studies which are fully forested or cleared and the scatter in the data

elasticity analysis (EA), Hydrological modelling (HM) or a combined use of statistical methods (SH) or quasi paired watershed analysis (QPW) (Figure 7). For larger catchments, these techniques all provide an approximation of the effect of forestry on streamflow rather than a direct comparison of catchments. This is a confounding factor that is not easily addressed in the regression modelling attempted here. Furthermore, the catchments analysed using EA, are concentrated in the drier end of the Dryness index scale compared to the other methods, with only the paired watershed experiment (PWE) assessment technique covering the full range of dryness indices.

In other words, the current data sets cannot resolve whether there actually is a non-linear catchment size \times forest cover effect, which then feeds into the buffering in larger catchments.

Apart from a difficulty of analysing complex confounding factors in the data, a general limitation of the type of analysis presented is that this work does not consider the spatial arrangement of the forest clearing in the catchments. While for fully or almost fully cleared smaller catchments this might not be an issue, it is perceivable that for larger catchments being partially cleared, a interaction between spatial location and clearing could be a factor in determining the change in streamflow. Clearing head water catchments on shallower soils might have a larger impact than clearing in downstream areas on deeper soils.

4.2. Model residuals

As pointed out earlier the residuals of the model diverge from the normal distribution for large positive and large negative residuals. These residuals are mainly associated with the small catchments from the paired watershed studies (Figure 7), which show very high variability. The final model removing the data with large values of Dryness and long study lengths has removed some of the spreading, mainly for the large negative residuals (Figure 8).

The reason why the regression model is better able to resolve the variance in the data for the negative residuals (generally related to increases in forest cover) compared the large positive residuals might link back to the issue of buffering and flow paths in the catchments. Small catchments that are stripped of most of the forest cover would provide little buffering, interception and infiltration, does leading to greater changes in flow. In contrast, revegetated catchments would have increased interception and buffering and therefore relatively smaller changes in flow. This also provides an explanation for the differences between forest cover removal and forest cover restoration (Figure 2).

4.3. The effect of assessment techniques with very small numbers of observations

Table 13: Distribution of assessment techniques in the data set

Assessment_technique	n
PWE	183
HM	57

Assessment_technique	n
SH	43
EA	34
QPW	7
QPW, EA	4
EA, HM	1
PWE, HM	1

One concern is that there are a few assessment techniques in the original dataset with a very low number of observations and this might skew the results of the analysis. This includes the category of Quasi paired watersheds and combinations of elasticity analysis and hydrological modelling (EA,HM) and paired watersheds and hydrological modelling (PWE,HM) (Table 13 and Figure 9).

Table 14: Statistical overview of the linear components of the model removing studies with limited observations in the assessment techniques

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-27.03	18.34	-1.47	0.14
DeltaF_perc_pos	0.29	0.1	2.96	0
Forest_Signincrease	-22.23	6.83	-3.25	0
Precip_data_typeOB	-11.91	13.85	-0.86	0.39
Precip_data_typeSG	18.07	15.65	1.15	0.25
Assessment_techniqueHM	36.48	12.3	2.97	0
Assessment_techniquePWE	56.52	12.33	4.58	0
Assessment_techniqueQPW	42.07	21.03	2	0.05
Assessment_techniqueSH	48.82	12.33	3.96	0
Forest_typeCF	-4.64	8.18	-0.57	0.57
Forest_typeMF	-5.22	8.08	-0.65	0.52
Hydrological_regimeSD	8.29	10.25	0.81	0.42

Table 15: Statistical overview of the smooth components of the model removing studies with limited observations in the assessment techniques

	edf	Ref.df	F	p-value
s(Dryness)	2.93	9	2.41	0
s(log10(Area_km2))	0.45	9	0.1	0.14
s(length)	0	9	0	0.4
s(From)	3.84	9	0.66	0.17




Figure 8: Residual plots for the final model indicating a small improvement in the residual distribution towards normal




Figure 9: Boxplot of the log base 10 of the catchment area (in km²) for the different assessment techniques, showing the dominance of small catchments in the paired watershed experiments

424 Concentrating only on the assessment techniques that have more than 10
 425 observations in the data set does not change much in the results (Table 14 and
 426 15). It strengthens the significance of the different assessment techniques and
 427 *Dryness* but generally results in the same interpretation. Overall this suggests
 428 that although those observations have some impact on the overall relationships,
 429 they do not strongly bias the outcomes.

430 However, the model results also clearly highlight that some of the assessment
 431 techniques (in particular paired watershed studies (PWE) and combined use
 432 of statistical methods and hydrographs (SH)), have a strong impact on the
 433 predicted change in flow. Particularly, relative to EA (elasticity approaches)
 434 all other assessment techniques have higher predicted changes in flow. In other
 435 words, there is a distinct difference in the way the change in flow is assessed,
 436 and the EA method (for example in Zhou et al. (2015)) appears to suggest a
 437 much smaller effect on the change in flow. However, as indicated earlier, the EA
 438 studies are all on the drier end of the *Dryness* spectrum, highlighting another
 439 unresolved interaction in the data.

440 4.4. The effect of climate

441 In drier catchments, changes in forest cover have greater impact on flow,
 442 which is similar to the observations in earlier studies (Filoso et al., 2017; Zhang
 443 et al., 2017; Zhou et al., 2015). This is most likely because in these catchments
 444 the overall flow is surface flow dominated and therefore the buffering that is
 445 afforded by the groundwater inputs is not as great. As the dataset currently
 446 does not include a separate variable for groundwater inputs (although this ef-
 447 fect is estimated in several of the studies), the effect again cannot be analysed
 448 separately.

449 Excluding the few catchments that have very high dryness values, clarified
 450 the *Dryness* trend, agreeing with earlier studies and showing a increase in the
 451 change in flow for drier catchments. However, this really only starts to have an
 452 effect for *Dryness* > 2.

453 4.5. Interactions

454 Generally this study did not consider interactions, but the above discussion
 455 suggest that there are possible several interactions. The relationships between
 456 forest cover change and *Area* (km^2) and between *Area* (km^2) and assessment
 457 technique have already been highlighted. However there are further unexplored
 458 interactions between the study length and some of the variables and *From* and
 459 the other variables.

460 A principle component analysis of the numeric data reveals some of these
 461 interactions (Figure 10), such as between *length* and *Dryness* and between *From*
 462 and *Area* (km^2). Including these interactions into the smooths of the models
 463 (data not shown) increases the explained variance slightly but does not funda-
 464 mentally change the significance of the different variables.




Figure 10: Biplot of the first two principle components using a principle component analysis on the numerical values of the

465 4.6. Further considerations

466 In contrast to Filoso et al. (2017), we did not identify the length of the
467 observation period as a significant variable in our final model. There are fur-
468 ther confounding factors in the data, which were also classified by Filoso et
469 al. (2017) and these might create biases in the data set that can impact the
470 overall assessment. For example, snow dominated hydrological regimes (SD),
471 are dominated by Coniferous Forests (CF), while the majority of the rain domi-
472 nated regimes are all broadleaf of mixed type forests (BF or MF). However, the
473 forest type classification is very coarse and does not fully capture possible phys-
474 iological differences that could affect evapotranspiration and therefore changes
475 in streamflow (Vervoort et al., 2021). This is not further investigated in this
476 study, but with more data available this might provide further opportunities for
477 investigations.

478 By making the updated the database of this study available, we hope that this
479 provides further incentive to investigate the impact of land cover change on
480 streamflow more generally.

481 5. Conclusions

482 More rigorous checking of data and statistical analysis results in both agree-
483 ment and disagreement with older studies. It demonstrates that analysis of
484 large databases of essentially “aggregated data” should be considered carefully.

485 Results of this statistical analysis need to be considered “conditional on the
486 data” Conditional on the data, the impact of forestry on streamflow is:

- 487 • Greater for forest clearing then for reforestation
- 488 • Possibly reduced for larger watersheds, but this is only in interaction with
489 other factors such as climate
- 490 • Increases for drier watersheds
- 491 • Sensitive to the assessment method used in the historical data

492 Stronger statements about the trends in the change in flow cannot be made
493 until more data or better data becomes available in this area. Furthermore, the
494 current study analyses a large global dataset. This analysis does not exclude
495 more local and regional effects that cannot be identified in the global data.

496 References

- 497 Andréassian, V., 2004. Waters and forests: From historical controversy to scien-
498 tific debate. *Journal of Hydrology* 291, 1–27. doi:[https://doi.org/10.1016/j.](https://doi.org/10.1016/j.jhydrol.2003.12.015)
499 [jhydrol.2003.12.015](https://doi.org/10.1016/j.jhydrol.2003.12.015)
- 500 Baker Jr., M.B., 1984. Changes in streamflow in an herbicide-treated pinyon-
501 juniper watershed in arizona. *Water Resources Research* 20, 1639–1642.
502 doi:<https://doi.org/10.1029/WR020i011p01639>

503 Borg, H., Bell, R.W., Loh, I.C., 1988. Streamflow and stream salinity in a
504 small water supply catchment in southwest western australia after reforesta-
505 tion. *Journal of Hydrology* 103, 323–333. doi:[https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-1694(88)90141-2)
506 1694(88)90141-2

507 Bosch, J.M., Hewlett, J.D., 1982. A review of catchment experiments to deter-
508 mine the effect of vegetation changes on water yield and evapotranspiration.
509 *Journal of Hydrology* 55, 3–23.

510 Brown, A.E., Western, A.W., McMahon, T.A., Zhang, L., 2013. Impact of forest
511 cover changes on annual streamflow and flow duration curves. *Journal of*
512 *Hydrology* 483, 39–50. doi:<http://dx.doi.org/10.1016/j.jhydrol.2012.12.031>

513 Brown, A.E., Zhang, L., McMahon, T.A., Western, A.W., Vertessy, R.A., 2005.
514 A review of paired catchment studies for determining changes in water yield
515 resulting from alterations in vegetation. *Journal of Hydrology* 310, 28–61.

516 Cosandey, C., Andréassian, V., Martin, C., Didon-Lescot, J.F., Lavabre, J.,
517 Folton, N., Mathys, N., Richard, D., 2005. The hydrological impact of the
518 mediterranean forest: A review of french research. *Journal of Hydrology*
519 301, 235–249. doi:<https://doi.org/10.1016/j.jhydrol.2004.06.040>

520 Filoso, S., Bezerra, M.O., Weiss, K.C.B., Palmer, M.A., 2017. Impacts of forest
521 restoration on water yield: A systematic review. *PLOS ONE* 12, e0183210.
522 doi:10.1371/journal.pone.0183210

523 Jackson, R.B., Jobbagy, E.G., Avissar, R., Roy, S.B., Barrett, D.J., Cook, C.W.,
524 Farley, K.A., Maitre, D.C. le, McCarl, B.A., Murray, B.C., 2005. Trading
525 water for carbon with biological carbon sequestration. *Science* 310, 1944–
526 1947. doi:10.1126/science.1119282

527 Kuczera, G., 1987. Prediction of water yield reductions following a bushfire
528 in ash-mixed species eucalypt forest. *Journal of Hydrology* 94, 215–236.
529 doi:10.1016/0022-1694(87)90054-0

530 Navas, R., Alonso, J., Gorgoglione, A., Vervoort, R.W., 2019. Identifying cli-
531 mate and human impact trends in streamflow: A case study in uruguay.
532 *Water* 11, 1433.

533 Peña-Arancibia, J.L., Dijk, A.I.J.M. van, Guerschman, J.P., Mulligan, M., Brui-
534 jnzeel, L.A., McVicar, T.R., 2012. Detecting changes in streamflow after
535 partial woodland clearing in two large catchments in the seasonal tropics.
536 *Journal of Hydrology* 416–417, 60–71. doi:[https://doi.org/10.1016/j.jhydrol.](https://doi.org/10.1016/j.jhydrol.2011.11.036)
537 2011.11.036

538 Roche, M., 1981. Watershed investigations for development of forest resources
539 of the amazon region in french guyana. *Tropical Agricultural Hydrology. J*
540 75–82.

541 Rodriguez, D.A., Tomasella, J., Linhares, C., 2010. Is the forest conversion to
542 pasture affecting the hydrological response of amazonian catchments? Sig-
543 nals in the ji-paraná basin. *Hydrological Processes* 24, 1254–1269. doi:<https://doi.org/10.1002/hyp.7586>

544 Ruprecht, J.K., Schofield, N.J., Crombie, D.S., Vertessy, R.A., Stoneman, G.L.,
545 1991. Early hydrological response to intense forest thinning in southwestern
546 australia. *Journal of Hydrology* 127, 261–277. doi:[https://doi.org/10.1016/](https://doi.org/10.1016/0022-1694(91)90118-2)
547 0022-1694(91)90118-2

549 Thornton, C.M., Cowie, B.A., Freebairn, D.M., Playford, C.L., 2007. The briga-
550 low catchment study: II*. Clearing brigalow (acacia harpophylla) for crop-
551 ping or pasture increases runoff. *Australian Journal of Soil Research* 45,
552 496–511. doi:doi:10.1071/SR07064

553 Trabucco, A., Zomer, R.J., 2018. Global aridity index and potential evapo-
554 transpiration (ET0) climate database v2. CGIAR consortium for spatial
555 information(CGIAR-CSI).

556 Vervoort, R.W., Dolk, M.M., Ogtrop, F.F. van, 2021. Climate change and
557 other trends in streamflow observations in australian forested catchments
558 since 1970. *Hydrological Processes* 35, e13999. doi:https://doi.org/10.1002/
559 hyp.13999

560 Wood, S., 2006. Generalized additive models: An introduction with r. CRC
561 Press, Boca Raton, FL.

562 Zhang, L., Zhao, F., Chen, Y., Dixon, R.N.M., 2011. Estimating effects of plan-
563 tation expansion and climate variability on streamflow for catchments in aus-
564 tralia. *Water Resources Research* 47, W12539. doi:10.1029/2011wr010711

565 Zhang, M., Liu, N., Harper, R., Li, Q., Liu, K., Wei, X., Ning, D., Hou, Y.,
566 Liu, S., 2017. A global review on hydrological responses to forest change
567 across multiple spatial scales: Importance of scale, climate, forest type and
568 hydrological regime. *Journal of Hydrology* 546, 44–59. doi:https://doi.org/
569 10.1016/j.jhydrol.2016.12.040

570 Zhao, F., Zhang, L., Xu, Z., Scott, D.F., 2010. Evaluation of methods for esti-
571 mating the effects of vegetation change and climate variability on streamflow.
572 *Water Resources Research* 46, W03505. doi:10.1029/2009wr007702

573 Zhou, G., Wei, X., Chen, X., Zhou, P., Liu, X., Xiao, Y., Sun, G., Scott,
574 D.F., Zhou, S., Han, L., Su, Y., 2015. Global pattern for the effect of
575 climate and land cover on water yield. *Nature Communications* 6, 5918.
576 doi:10.1038/ncomms6918

577 Zhou, G., Wei, X., Luo, Y., Zhang, M., Li, Y., Qiao, Y., Liu, H., Wang, C.,
578 2010. Forest recovery and river discharge at the regional scale of guangdong
579 province, china. *Water Resources Research* 46. doi:https://doi.org/10.1029/
580 2009WR008829