

Supplementary Information part 1: Changes to original database

R. Willem Vervoort¹, Eliana Nervi, Jimena Alonso

Abstract

This supplementary material file gives an overview of the changes that were made in the original data based on review of the original literature. Overall 36 data points were changed. The most common problem was a change in the sign of the forest change or the streamflow change.

1. Introduction

This supplementary material is related to ‘Generalizing the impact of forest cover on streamflow from experimental data: it is not that simple. Vervoort et al.’

This document list all the changes that were made to the original Zhang et al. [19] database after review of all the original papers.

2. Discussion of changes

Need to fix up references

A particular problem was that many catchments appeared to have the wrong sign for the change in forest cover. There are many catchments with reported positive change in cover and a large increase in flow. These were all checked and corrected if needed and a full list of all these changes is below:

- 76, Beaver Creek, the flow was corrected from 600% to 157% after review of the original publication [2].
- 124, D3, Amatya and Skaggs [1]: The originally recorded 250% change by Zhang et al. [19] is clearly wrong. The paper says on page 7: Both of these outflow ratios (0.64 and 0.50) were higher than the calculated expected values of 0.55 for 2003 and 0.44 for 2005, respectively. So the value should be $0.64/0.55 * 100 - 100$ or $0.5/0.44 * 100 - 100$: 16% or 13%. Corrected to 16%.
- 3, Baker Creek, Zhang and Wei [18]. The original recorded 201.1% change by Zhang et al. [19] also seems wrong. Original paper says on page 2031: Annual mean flow has been increased by 47.6%. corrected

*Corresponding author

Email addresses: willem.vervoort@sydney.edu.au (R. Willem Vervoort),
eliananervif@gmail.com (Eliana Nervi), jalonso@fing.edu.uy (Jimena Alonso)

Preprint submitted to Elsevier

October 7, 2022

- 67, April rd, which is incorrectly attributed to Ruprecht and Schofield [15] in Zhang et al. [19]. This is actually from Ruprecht and Schofield [14] and the original paper clearly indicates “clearfelling”. As a result the change in forest cover was changed to -100% rather than +100%.
- 210, March rd, 100, 147.6. Same problem as 67, Bari et al. [3] clearly state that the catchment was cleared, so therefor the change in forest cover changed to -100%.
- 213, 214 and 215, Monda 1, 2 and 3. These catchments are tricky. The original paper [13] only reports on the control period and indicates that the catchments will be cleared. The later summary paper [17] shows the timeseries of the flow change, but does not report a single value, so the values in the database must have been estimated from the timeseries. The further complication is that the treatment included clearing and reseedling and regrowth. This suggest that the records should be removed from the database, or only the first few years of the experiment used. In any case, if the values are kept, the sign of the change in forest cover needs to be changed to negative (Clearing).
- 230, Oleolega catchment. The paper describes a removal of forest up to 85%. changed Delta_F_perc to -85 from 90.
- 312, Yerraminup South. The original publication for this catchment is a Western Australian Water Authority report from 1987, which is hard to find, but we have added a copy in the “Papers” folder on github. In this report, in Table 2 on page 11, for the catchment a “Crown cover” decrease of 60% is given. Changed the sign of the change in forest cover: -60%.
- 72 Barratta, 100 Coachwood, 103 Corkwood, and 83 Bollygum, as cited by Cornish [10] and Cornish and Vertessy [11]. In the database from Zhang et al. [19], the forest change for all these catchments is positive. However, the paper highlights that these catchments were all logged and either naturally regenerated or were planted with a plantation species. So, similar to the the earlier mentioned Monda catchments, the reported change probably only refers to the first couple of years after clearing (before regrowth). In any case, the reported change in forest cover should be negative (clearing) rather than positive. Corrected for all three catchments.
- 78, Black Spur 1, the treatments and effects are only reported in a conference paper [12] and once again indicated clearing, meaning that the change in forest cover should be negative rather than positive (as reported in Zhang et al. [19]). Corrected. Similar to other paired watershed experiments, only the first couple of years can be linked to the effect as later regrowth cancels out part of the increase in flow.
- 104, Coshocton. Checking the original paper indicates that this is in fact a reduction in flow as a result of reforestation. Changed the sign of Delta_Q_f to be negative.
- 102, Cold Spring. Checking the original paper [16] indicates that this is in fact a reduction in flow as a result of reforestation. Changed the sign of Delta_Q_f to be negative.
- 85 Bosboukloof. This is essentially a duplicate of catchment 184, but

the cited paper analyses only 1 year of runoff after a major fire. In any case, the data should reflect a decrease in forest cover: changed the sign of Delta_f_perc to -80%.

- 259 Shackam Brook. There were a few issues with this catchment in the original database. The name was misspelled and it was incorrectly attributed to Brown et al. [8]. The original paper is the same as 102 [16]. Finally, the catchments were all reforestation as the title of the original report indicates and the reported streamflows are all decreases. Corrected Delta_Qf_perc to -20.7%.
- 95 Sage Brook. Similar to 259 and 102, originates from Schneider and Ayer [16]. Reforestation so Delta_Qf_perc corrected to -19.8%.
- 101 Coalburn. Original publication (Robinson, 1993) which is a symposium paper, can not be located, even after contacting some of the authors. The best summary of the research is in Birkinshaw et al. [5] which summarizes 45 years of research in the Coalburn catchment. It was a reforestation experiment, and there was a decrease in the streamflow over the longer time period. Changed to -20.3%.

A further issue was the inclusion of the results of several catchments, for example from the study by Beck et al. [4], which had no significant change in flow. Despite this, the “average” change in flow was reported in the database. We don’t believe that this is correct and the results from such studies should be set to 0. A full list of changes is provided below:

- 97 Cibucio, 123 Culebrinas, 244 Portugues, 161 Grande de Loiza, 271 Tanama, 132 Fajardo, 89 Canovanas, 73, Bauta, 163 Grande de Patillas, 283 Valenciano, 181 Inabon, and 162 Grande de Manati. These are all catchments in Puerto Rico from the study from Beck et al. [4]. They should probably be removed from the database as the paper clearly indicates that there is no evidence of a change in flow due to reforestation. The values that are cited in the database should all be set to “not significant from 0”, so might be included as 0. Including them with positive or negative values is misleading. This study is a very detailed hydrological modelling study, but in the end finds no significant change in streamflow as a result of deforestation. Values for all 12 studied catchments set to 0 in the database.
- 188 Kimakia. and 254 Sambret. The data in the database from Zhang et al. [19] appear to originate from Bruijnzeel et al. [9] which gives 3 values for different lengths of studies. However, the values in the original study by Blackie [7] and Blackie [6] do not seem to add up to the same values, and the specific values are not mentioned in the actual papers. In addition, as Bruijnzeel et al. [9] mentions in the footnotes, the control for Kimakia is a bamboo catchment, while the control for Sambret is a tea plantation. Overall, this suggests that the data are probably not a clear deforestation/reforestation study and should be discarded from the analysis.

- 221 N. Creek, Babinda, Queensland. The original paper from this study highlights that the differences between the catchments were insignificant.

References

- [1] Devendra M Amatya and Wayne R Skaggs. Effects of thinning on hydrology and water quality of a drained pine forest in coastal north carolina. In *21st Century Watershed Technology: Improving Water Quality and Environment Conference Proceedings, 29 March-3 April 2008, Concepcion, Chile*, page 62. American Society of Agricultural and Biological Engineers, 2008.
- [2] Malchus B. Baker Jr. Changes in streamflow in an herbicide-treated pinyon-juniper watershed in arizona. *Water Resources Research*, 20(11):1639–1642, 1984. ISSN 0043-1397. doi: <https://doi.org/10.1029/WR020i011p01639>. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/WR020i011p01639>.
- [3] M. A. Bari, N. Smith, J. K. Ruprecht, and B. W. Boyd. Changes in streamflow components following logging and regeneration in the southern forest of western australia. *Hydrological Processes*, 10(3):447–461, 1996. ISSN 0885-6087. doi: [https://doi.org/10.1002/\(SICI\)1099-1085\(199603\)10:3<447::AID-HYP431>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1099-1085(199603)10:3<447::AID-HYP431>3.0.CO;2-1). URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/%28SICI%291099-1085%28199603%2910%3A3%3C447%3A%3AAID-HYP431%3E3.0.CO%3B2-1>.
- [4] H. E. Beck, L. A. Bruijnzeel, A. I. J. M. van Dijk, T. R. McVicar, F. N. Scatena, and J. Schellekens. The impact of forest regeneration on streamflow in 12 mesoscale humid tropical catchments. *Hydrol. Earth Syst. Sci.*, 17(7):2613–2635, 2013. ISSN 1607-7938. doi: 10.5194/hess-17-2613-2013. URL <https://hess.copernicus.org/articles/17/2613/2013/>. HESS.
- [5] Stephen J. Birkinshaw, James C. Bathurst, and Mark Robinson. 45 years of non-stationary hydrology over a forest plantation growth cycle, coalburn catchment, northern england. *Journal of Hydrology*, 519:559–573, 2014. ISSN 0022-1694. doi: <https://doi.org/10.1016/j.jhydrol.2014.07.050>. URL <https://www.sciencedirect.com/science/article/pii/S0022169414005848>.
- [6] JR Blackie. 2.2. 1 the water balance of the kericho catchments. *East African Agricultural and Forestry Journal*, 43(sup1):55–84, 1979.
- [7] JR Blackie. 3.2. 1 the water balance of the kimakia catchments. *East African Agricultural and Forestry Journal*, 43(sup1):155–174, 1979.
- [8] Alice E. Brown, Lu Zhang, Thomas A. McMahon, Andrew W. Western, and Robert A. Vertessy. A review of paired catchment studies for determining changes in water yield resulting from

- alterations in vegetation. *Journal of Hydrology*, 310(1-4):28–61, 2005. URL <http://www.sciencedirect.com/science/article/B6V6C-4G05MM9-1/2/bbc5fc0e958a8f34bcb7c1cc7fa57b48>.
- [9] Leendert Adriaan Bruijnzeel et al. Hydrology of moist tropical forests and effects of conversion: a state of knowledge review. *Hydrology of moist tropical forests and effects of conversion: a state of knowledge review.*, 1990.
- [10] P. M. Cornish. The effects of logging and forest regeneration on water yields in a moist eucalypt forest in new south wales, australia. *Journal of Hydrology*, 150(2-4):301–322, 1993. URL <http://www.sciencedirect.com/science/article/B6V6C-487D3Y2-9J/2/73c981ba76284d9d629f6b221d6fd6c6>.
- [11] P. M. Cornish and R. A. Vertessy. Forest age-induced changes in evapotranspiration and water yield in a eucalypt forest. *Journal of Hydrology*, 242(1-2):43–63, 2001. URL <http://www.sciencedirect.com/science/article/B6V6C-429910G-3/2/0158b1f89ff436f338a9e688a47f06c4>.
- [12] M. D. A. Jayasuriya and P. J. O’Shaughnessy. *The Use of Mathematical Models in Evaluating Forest Treatment Effects on Streamflow*, pages 135–139. Hydrology and Water Resources Symposium, 1988. doi: 10.3316/informit.692214289455295. URL <https://search-informit-org.ezproxy.library.sydney.edu.au/doi/10.3316/informit.692214289455295>. doi: 10.3316/informit.692214289455295.
- [13] P. J. O’Shaughnessy, K. J. Langford, H. P. Duncan, and R. J. Moran. Catchment experiments in mountain ash forests at north maroonah. *Australian Forestry*, 42(3):150–160, 1979. ISSN 0004-9158. doi: 10.1080/00049158.1979.10674220. URL <https://doi.org/10.1080/00049158.1979.10674220>. doi: 10.1080/00049158.1979.10674220.
- [14] J. K. Ruprecht and N. J. Schofield. Analysis of streamflow generation following deforestation in southwest western australia. *Journal of Hydrology*, 105(1):1–17, 1989. ISSN 0022-1694. doi: [https://doi.org/10.1016/0022-1694\(89\)90093-0](https://doi.org/10.1016/0022-1694(89)90093-0). URL <https://www.sciencedirect.com/science/article/pii/0022169489900930>.
- [15] J. K. Ruprecht and N. J. Schofield. Effects of partial deforestation on hydrology and salinity in high salt storage landscapes. i. extensive block clearing. *Journal of Hydrology*, 129(1):19–38, 1991. ISSN 0022-1694. doi: [https://doi.org/10.1016/0022-1694\(91\)90042-G](https://doi.org/10.1016/0022-1694(91)90042-G). URL <https://www.sciencedirect.com/science/article/pii/002216949190042G>.
- [16] William Joseph Schneider and Gordon Roundy Ayer. Effect of reforestation on streamflow in central new york. Report 1602, 1961. URL <http://pubs.er.usgs.gov/publication/wsp1602>.

- 198 [17] Fred Watson, Rob Vertessy, Tom McMahon, Bruce Rhodes, and Ian
199 Watson. Improved methods to assess water yield changes from paired-
200 catchment studies: application to the maroondah catchments. *For-
201 est Ecology and Management*, 143(1):189–204, 2001. ISSN 0378-1127.
202 doi: [https://doi.org/10.1016/S0378-1127\(00\)00517-X](https://doi.org/10.1016/S0378-1127(00)00517-X). URL [https://www.
203 sciencedirect.com/science/article/pii/S037811270000517X](https://www.sciencedirect.com/science/article/pii/S037811270000517X).
- 204 [18] Mingfang Zhang and Xiaohua Wei. Contrasted hydrological responses to
205 forest harvesting in two large neighbouring watersheds in snow hydrol-
206 ogy dominant environment: implications for forest management and fu-
207 ture forest hydrology studies. *Hydrological Processes*, 28(26):6183–6195,
208 2014. ISSN 0885-6087. doi: <https://doi.org/10.1002/hyp.10107>. URL
209 <https://onlinelibrary.wiley.com/doi/abs/10.1002/hyp.10107>.
- 210 [19] Mingfang Zhang, Ning Liu, Richard Harper, Qiang Li, Kuan Liu, Xiao-
211 hua Wei, Dingyuan Ning, Yiping Hou, and Shirong Liu. A global re-
212 view on hydrological responses to forest change across multiple spatial
213 scales: Importance of scale, climate, forest type and hydrological regime.
214 *Journal of Hydrology*, 546:44–59, 2017. ISSN 0022-1694. doi: [https://
215 doi.org/10.1016/j.jhydrol.2016.12.040](https://doi.org/10.1016/j.jhydrol.2016.12.040). URL [http://www.sciencedirect.
216 com/science/article/pii/S0022169416308307](http://www.sciencedirect.com/science/article/pii/S0022169416308307).