

CSE 461: Programming Languages Concepts

Prof. G. Tan
Spring 2018

Homework 5: **Due on Mar 30th before class (12:20pm) in Canvas.**

We will allow late homework for up to only one day with 10% penalty this time, as I want to release a solution soon after the due date to help students prepare for the second midterm.

Submission: Please submit your homework via Canvas. It's okay if you submit a scanned version of your on-paper answers, but please make sure your scanned version is legible.

1. (5 points) For the lambda-calculus term $(\lambda x. \lambda y. x) (\lambda z. y)$.
 - (a) (2 point) Calculate its free variables using the FV function. Show the steps.
 - (b) (2 point) Use lambda calculus reduction to reduce the expression to a normal form. Begin by renaming bound variables and show every step.
 - (c) (1 point) Describe what would go wrong if you did not rename bound variables.
2. (3 points) Reduce the following lambda-calculus term to the normal form. Show all intermediate steps, with one beta reduction at a time. In the reduction, assume that you are supplied with extra rules that allow you to reduce the addition of two natural numbers into the corresponding results.

$$(\lambda f. \lambda x. f (f (f x))) (\lambda y. y + 2) 2$$

3. (3 points) The Algol-like program fragment

```
function f(x)
  return x+5
end;
function g(y)
  return 2-y
end;
f(g(2));
```

can be written as the following lambda expression:

$$\left((\lambda f. \lambda g. f (g\ 2)) (\lambda x. x + 5) \right) (\lambda y. 2 - y)$$

Reduce the expression to a normal form. In the reduction, assume that you are supplied with extra rules that allow you to reduce the addition or subtraction of two natural numbers into the corresponding results.