

UiO : **Department of Physics**
University of Oslo

Application of Supervised Machine Learning to the Search for New Physics in ATLAS data

A Study of Ordinary Dense, Parameterized
and Ensemble Networks and their Application
to High Energy Physics

William Hirst

May 29, 2023

1 Introduction & Motivation

2 The Implementation

3 Methods & Results

4 Conclusion & Outlook

1 Introduction & Motivation


2 The Implementation

3 Methods & Results

4 Conclusion & Outlook



Why apply machine learning to HEP problems?

- The standard model (SM) is successful, but not enough
 - Neutrino masses
 - Gravity
- Large amount of data
- Machine learning (ML)



How do we search for new physics?

- Compare theory with experiment
 - Experiment: Measured
 - Theory: Simulated
- Search regions
- Expected significance
 - $Z_{\text{exp}} \approx \frac{\text{signal}}{\sqrt{\text{background}}}$
- Difficult to separate \rightarrow ML



Aim

- 1 Signal → background
- 2 Use ML to separate signal from background
- 3 Measure performance in Z_{exp}
- 4 Study and compare

The Implementation

1 Introduction & Motivation


2 The Implementation

3 Methods & Results

4 Conclusion & Outlook



The Super Symmetry (SUSY) signal

- Chargino-neutralino production
- Free parameters → masses
- Nr-of-Events(Mass)



A summary of the applied methods

- Three neural network variants
 - Ordinary dense neural network
 - Ensemble networks/Local-Winner-Takes-All (LWTA)
 - Parameterized neural networks (PNN)
- One boosted decision tree method



Training strategy

- Classification
 - Background → 0
 - Signal → 1
- 80% training and 20% validation
- Early stopping criteria
 - Performance on validation set


1 Introduction & Motivation

2 The Implementation


3 Methods & Results

4 Conclusion & Outlook


“Ordinary” dense neural network



Compare one-mass approach to several-masses approach




One-mass-model (OMM)



Several-mass-model (SMM)


Parameterized neural network

- Long-term memory
- PNN → signal includes mass parameter in feature set
- Background assigned parameters randomly




Ensemble methods - Local-Winner-Takes-All (LWTA)

- Dropout
- Competing nodes - Units
- Encode information in pattern specific pathways





Channel-Out, Maxout and SCO





Visualization and study of sparse pathways

- Visualize activation and paths
- The bolder the line the more frequently the path is used.
- Color of lines
 - Pink: SM background
 - Blue: SUSY signal




Visualization and study of sparse pathways


- Visualize activation and paths
- The bolder the line the more frequently the path is used.
- Color of lines
 - Pink: SM background
 - Blue: SUSY signal



Comparing activation of Maxout with SCO




Maxout




SCO

Ensemble network architecture


MaxOut:



Comparing sensitivity of channel-out, SCO and maxout




Comparing methods on full signal grid



Comparing the methods to previous analysis

- Compare the expected limits to analysis made by ATLAS in 2021 [4]



Conclusion & Outlook

1 Introduction & Motivation

2 The Implementation

3 Methods & Results

4 Conclusion & Outlook

Key findings

- 1 Diverse signal → improve performance
- 2 SCO
- 3 PNN bias towards smaller masses
- 4 Maxout achieved balanced performance

The way forwards

- 1 More advanced analysis of ML output
- 2 LWTA promising (SCO)
- 3 Combine PNN and LWTA

References

-  Maximilien Brice.
‘Installing the ATLAS calorimeter. Vue centrale du détecteur ATLAS avec ses huit toroides entourant le calorimètre avant son déplacement au centre du détecteur’.
<https://cds.cern.ch/record/910381>
Figure on front page
-  Joao Pequenao.
‘Event Cross Section in a computer generated image of the ATLAS detector.’.
<https://cds.cern.ch/record/1096081>
Figure on slide 3
-  ATLAS Collaboration.
‘ATLAS event at 900 GeV - 6 May 2015 - Run 264034 lb 659 event 11526514’.
<https://cds.cern.ch/record/2015238>
Figure on slide 4
-  ATLAS Collaboration [4].
‘Search for chargino–neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector’.
<http://arxiv.org/abs/2106.01676>



William Hirst



**Application of Supervised Machine Learning
to the Search for New Physics in ATLAS data**

A Study of Ordinary Dense, Parameterized and Ensemble Networks and their Application to High Energy Physics