



Full Length Article

Alterations in T-helper cell type 1 and blood cell parameters in malaria-infected patients



Olarewaju Abdulkareem Babamale^{a,*}, Adam Olaitan Abdulkareem^b, Olufunke A. Opeyemi^a, Uade Samuel Ugbomoiko^a

^aParasitology Unit, Department of Zoology, University of Ilorin, Ilorin, Nigeria

^bPhysiology Unit, Department of Zoology, University of Ilorin, Ilorin, Nigeria

ARTICLE INFO

Article history:

Received 13 October 2016

Received in revised form 11 May 2017

Accepted 11 May 2017

Available online 23 May 2017

Keywords:

Cytokine

Thrombocytopenia

Parameters

Alterations

Prevalences

ABSTRACT

Malaria is a global public health disease. Haematological and cytokine alterations are the major sources of its pathological conditions. Therefore, blood and serum of patients attending health centres were screened to investigate the effects of *Plasmodium falciparum* on the T-helper cell type 1 and blood cell parameters using Enzyme linked immunosorbent assay and automatic hematology analyzer respectively. Approximately 55% of malaria-infected patients with average parasitaemia of 2523.64 parasite/ μ l of blood concurrently suffered anaemia, thrombocytopenia, leucopenia, microcytosis and hypochromasia. However, thrombocytopenia and leucopenia were age-specific and their prevalences in children within ≤ 10 years were higher. These disease conditions significantly vary with severity of malaria infection ($p < 0.05$). Blood parameters with the exception of RBC and MCHC were significantly lower in the infected patients ($p < 0.05$) with 12.9% and 41.2% reduction in haemoglobin and platelet counts respectively. A high plasma concentration of IL-10, IL-12, INF- γ and TNF- α , ratio of IL-10/TNF- α (1.86) and IL-10/INF- γ (1.55) were recorded among the malaria-infected groups. This study revealed that unregulated interaction of the parasite with host immune response has important consequences in disease progression and thus relevant for therapeutic and vaccine development.

© 2017 Mansoura University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Malaria is still a life-threatening parasitic disease in many countries worldwide, causing between 80 and 90% morbidity and mortality in sub-Saharan Africa [1,2]. In this region, *Plasmodium falciparum* and *P. vivax* are common human malaria parasites and the prognosis is severe [3]. The activities of their erythrocytic asexual stages in asymptomatic and symptomatic patients are important foci in the pathogenesis [4,5]. Infections in less immuned individuals, especially young children and pregnant women are always severe, resulting in anaemia, thrombocytopenia, leucopenia and other haematological disorders [4]. Most infected individuals are incapacitated for days due to recurrent clinical episodes attributed to the release of malaria pigment (hemozoin) and other toxic factors such as glycosylphosphatidylinositol (GPI) when the infected erythrocytes rupture [5].

Previous studies [6,7] showed that *P. falciparum*, upon invasion of red blood cell, interferes with rigidity and deformability

properties of the cell to permit the direct exchange of metabolites. Consequently, inflammatory cytokines (TNF- α , IL-1, IL-10 and IFN- γ), cascade coagulation and sequestration of parasitized RBCs increase, and thus the manifestations of malaria infection ensued [8–10].

However, a steady balance between pro- and anti-inflammatory responses is obligatory to elicit anti-parasitic activities and enhance parasite clearance; otherwise an uncontrolled production of pro-inflammatory cytokines, such as IL-6, TNF and IFN- γ may lead to severe malaria syndromes thereby suppressing T-cell response [11,12]. The resulting protective Th1-dependent immune responses to blood-stage malaria infection are mediated by IFN- γ and TNF- α to optimize nitric oxide production [13,14].

Report [15] has shown that infections due to *P. falciparum* result in progressive structural, biochemical, and mechanical modifications of the host cells that lead to life-threatening complications. Information on the mechanism of alteration in infected host cell required in predicting the severity of malaria is limited. This study therefore investigated the effect of *P. falciparum* on the type 1 T-helper cell and blood cell parameter amongst the malaria infected patients in Ilorin, Kwara state, Nigeria for better understanding of

* Corresponding author.

E-mail address: olas4nice2004@yahoo.co.uk (O.A. Babamale).

host-parasite interaction which can be exploited in the effective management of malaria infection.

2. Materials and methods

2.1. Study design, sample and data collection

The blood and serum of patients attending health centres in Ilorin, Kwara State (located at Lat. 8° 30'N and Long. 8° 50'N and Long. 4° 20'E and 4° 35'E), Nigeria were used for this study. Thick and thin blood films were prepared and stained with Giemsa and *P. falciparum* was identified under oil immersion microscope. The estimation of parasitaemia was done by counting the number of parasites per 200 white blood cells (WBC) on blood films. Complete blood cell counts was conducted using an automatic hematology analyzer (Pentra ABX) and peripheral blood smears were performed for routine differential blood cellular quantification.

In accordance with manufacturer's recommendations, the levels of TNF- α , IL-10 and IL-12 in the serum were determined using Enzyme linked immunosorbent assay, (ELISA) Kit procured from Abcam, Cambridge, United Kingdom. The test kits were sensitive detecting cytokines levels at less than 0.8 pg/ml and cross reactivity was not observed for any other proteins tested.

2.2. Ethical clearance and procedure for data collection

The study protocol was approved by University of Ilorin ethical review committee and the Medical and Information Management units of selected health centres in Ilorin, Kwara state. The approval detail of the study was contained in the letter with reference number KW/REF/GT0126.

2.3. Statistical analysis

Statistical analysis was performed with SPSS version 20; (SPS Inc., Chicago IL). Descriptive statistics were expressed as means and standard deviations while the student's t-test and Chi square analysis were applied to determine the differences in haematological parameters and cytokines levels. Comparison of associations between the haematological parameters and parasite density were assessed using the Pearson's linear regression. The level of statistical difference was set at $p < 0.05$.

3. Results

Of the total 223 out-patients (100 males and 123 females) with average age of 21.41 years enrolled in the study, 55.2% was infected with *Plasmodium falciparum* with average parasitaemia of 2523.64 parasite/ μ l of blood (Table 1). Approximately, 25%, 26% and 5% of malaria-infected individuals had low, moderate and severe infection respectively. Similarly, 55.2% and 41.7% of infected patient were anaemic and thrombocytopenic respectively. Table 2 shows that the prevalences of haematological disorder among the malaria-infected patients was not sex related ($P > 0.05$) but thrombocytopenia and leucopenia were age-specific with high prevalences in children within age-group of ≤ 10 years. However, these disease conditions significantly vary with degree of malaria infection ($p < 0.05$). For instance, 12.9% and 41.2% reduction were observed in haemoglobin and platelet counts among infected participants.

Table 3 revealed that lower values of red blood cell, packed cell volume, haemoglobin and differential counts of neutrophil, platelets and white blood cells were recorded among malaria patients. High plasma concentrations of IL-10, IL-12, INF- γ and TNF- α , also ratio of IL-10/TNF- α (1.86) and IL-10/INF- γ (1.55) were obtained

Table 1

Demographical and clinical characteristics of study population.

	N (%)
Sex (Male/Female)	100/123
Mean Age + Std.dev.	21.41 \pm 10.27
Number infected with malaria	123(55.2)
Mean intensity + Std.dev	2523.64 \pm 4.47
Degree of malaria infection	
No Infection	100(44.8)
Low	55(24.7)
Moderate	57(25.6)
Severe	11(4.9)
Anaemia	
Malaria patient	123(55.2)
Uninfected patient	7(3.1)
Thrombocytopenia	
Malaria patient	93(41.7)
Uninfected patient	21(9.4)

among the infected patients (Table 3). Further analysis indicated that haematological indices significantly decreased ($p < 0.05$) whereas immunological indices increased with increasing parasitaemia (Table 4).

4. Discussion

Pronounced haematological and cytokine alterations are trademarks in malaria infection and other infectious diseases [4,13,14]. The current study confirms that anaemia, thrombocytopenia, hypochromasia and leucopenia were common disorders in malaria infected patients. This is similar to reports in many malaria-endemic settings worldwide [16–19]. However, the relative severity of the haematological disorders varied, similar in infected male and female but higher in young children. This observed age and sex relative pattern of morbidity is a common characteristic in endemic areas in Africa [20,21]. The severity of these disorders is observed to be increasing with parasite density in infected individuals except in the case of hypochromasia which is more prevalent among individual with low parasite density. This observation is concordant with reports of Erhart et al. [22] and Saravu et al. [23].

Anaemia is a common clinical indicator of severe malaria in children [24,25], a phenomenon that follows inhibition of erythropoiesis and massive destruction of both parasitized and non-parasitized red blood cells. The low mean haemoglobin concentration (Hb) and packed cell volume (PCV) observed in our infected patient indicate anaemic condition. This is similar to many reports [26–28]. The reduction in Hb and PCV increases with the intensity of infection in malaria [29]. Thus, individual with severe malaria are more susceptible to severe anaemia.

Thrombocytopenia and platelet dysfunction are another important pathological conditions observed in 76.4% malaria infected group in this study. This agrees with previous findings [30–33]. Other studies [34,35] reported that 95% of falciparum malaria patient had thrombocytopenia. The mechanism of this disease condition has been well described [36–38]. The authors postulated that immune mediated lysis, sequestration in the spleen and a dyspoietic process in the bone marrow diminish platelet production which may probably lead to reduced platelets subpopulation in the circulation. However, the pathogenic mechanisms by which platelets mediate disease severity in patients with falciparum malaria remain to be delineated. Previous study [39] revealed that platelets in patients with *P. falciparum* expressed Toll-like receptors (TLRs), which release prepackaged inflammatory mediators such as Nitric oxide (NO), a key mediator of platelet homeostasis. A decreased bioavailability of NO in patients with severe malaria may contribute to increased platelet activation and consumption

Table 2

Prevalence of haematological disorders with respect to age, sex and infection status of the malaria infected participants.

Variables	No. examined	Haematological conditions				
		Anaemia (%)	Thrombocytopenia (%)	Leucopenia (%)	Microcytosis (%)	Hypochromasia (%)
Overall	123	100(81.3)	94(76.4)	93(75.6)	26 (21.1)	56 (45.5)
Sex						
Male	55	44 (80.0)	42 (76.4)	44(80.0)	12 (21.8)	22(40.0)
Female	68	56 (82.4)	52 (76.4)	49(72.1)	14 (20.6)	34(50.0)
P. value		0.739	0.350	0.308	0.818	0.517
Age group (years)						
≤10	23	21 (91.3)	20 (86.9)	20(86.9)	4 (17.4)	11(47.8)
11–20	25	19 (76.0)	16 (64.0)	19 (76.0)	7 (28.0)	10 (40.0)
21–30	59	49 (83.1)	46 (77.9)	14(23.7)	11 (18.6)	27 (45.8)
≥31	16	11 (68.8)	12 (75.0)	7(43.8)	4 (25.0)	8 (50.0)
P. value		0.289	0.005	0.028	0.791	0.531
Malaria infection						
Low	55	38 (69.1)	42 (76.4)	37(67.3)	9 (16.4)	29 (52.7)
Moderate	57	54 (94.7)	44 (77.2)	46(80.7)	16 (28.1)	23(40.4)
Severe	11	8 (72.7)	8 (72.7)	10(90.9)	1 (9.1)	4 (36.4)
P. value		0.002	0.002	0.018	0.174	0.047

Table 3

Comparison of haematological and immunological indices among infected and uninfected individuals in the study populations.

	Uninfected patients		Infected patients		P. value
	Mean ± Std.dev.	95% CI	Mean ± Std.dev.	95% CI	
<i>Hematological indices</i>					
RBC ($10^3/\mu\text{l}$)	3.68 ± 0.88	3.17–3.79	3.46 ± 0.63	3.55–3.77	0.724
Haemoglobin (g/dl)	14.44 ± 0.63	10.22–15.66	9.09 ± 1.54	6.82–9.37	<0.001
PCV (%)	33.96 ± 4.92	32.22–35.71	26.45 ± 4.07	25.73–27.18	<0.001
MCHC (g/dl)	32.53 ± 2.55	31.63–33.44	32.17 ± 5.37	31.22–33.13	0.712
MCV (pg)	89.83 ± 15.20	84.44–95.22	71.90 ± 8.03	70.46–73.33	<0.001
MCH (pg)	31.01 ± 4.66	29.36–32.66	26.81 ± 2.85	26.29–27.31	<0.001
WBC ($10^3/\mu\text{l}$)	3.82 ± 0.69	36.09–41.75	3.21 ± 0.85	31.18–34.13	0.007
Neutrophil ($10^3/\mu\text{l}$)	2.03 ± 0.42	19.11–22.77	1.21 ± 0.57	11.41–13.48	<0.001
Basophil ($10^3/\mu\text{l}$)	0.02 ± 0.003	22.25–23.18	0.04 ± 0.02	38.63–44.88	<0.001
Eosinophil ($10^3/\mu\text{l}$)	1.43 ± 0.18	14.66–15.56	3.26 ± 0.51	30.33–35.15	<0.001
Platelet count ($10^3/\mu\text{l}$)	80.27 ± 10.07	76.70–83.84	47.16 ± 26.31	42.46–51.85	<0.001
<i>Immunological indices</i>					
INF- γ (pg/ml)	41.43 ± 9.81	39.27–4.32	51.04 ± 8.57	43.63–54.74	<0.001
TNF- α (pg/ml)	27.65 ± 8.28	26.10–2.91	42.37 ± 5.52	40.18–44.35	<0.001
IL-10 (pg/ml)	25.18 ± 13.18	27.32–3.10	79.01 ± 10.07	20.19–30.17	0.278
IL-12 (pg/ml)	10.82 ± 10.13	6.07–11.21	23.11 ± 0.20	20.02–20.70	0.282
IL-10/TNF- α ratio	0.91 ± 1.52	0.32–0.96	1.86 ± 1.31	0.31–2.12	0.001
IL-10/INF- γ ratio	0.61 ± 0.84	0.47–0.80	1.55 ± 0.26	0.61–1.98	0.001

Table 4

Variation of haematological and immunological indices with intensity of infection.

	Intensity of infection			P. value
	Low	Moderate	Severe	
<i>Haematological indices</i>				
RBC ($10^3/\mu\text{l}$)	3.77 ± 0.64	3.63 ± 0.63	3.31 ± 0.45	0.208
Haemoglobin (g/dl)	9.46 ± 1.49	8.98 ± 1.44	7.7 ± 1.49	<0.001
PCV (%)	27.78 ± 4.47	25.74 ± 5.19	25.30 ± 2.99	<0.001
MCHC (g/dl)	31.59 ± 6.27	32.32 ± 4.70	34.37 ± 2.93	0.416
MCV	72.01 ± 5.62	71.05 ± 9.16	75.70 ± 11.28	<0.001
MCH (pg)	27.30 ± 3.20	26.66 ± 2.31	24.96 ± 2.80	<0.001
WBC ($10^3/\mu\text{l}$)	3.45 ± 0.85	2.99 ± 0.86	3.21 ± 0.48	<0.001
Neutrophil ($10^3/\mu\text{l}$)	1.12 ± 0.53	1.22 ± 0.58	1.67 ± 0.46	<0.001
Basophil ($10^3/\mu\text{l}$)	0.05 ± 0.001	0.04 ± 0.001	0.01 ± 0.002	<0.001
Eosinophil ($10^3/\mu\text{l}$)	3.50 ± 1.33	3.41 ± 1.62	1.52 ± 0.62	<0.001
Platelet count ($10^3/\mu\text{l}$)	51.00 ± 19.70	47.18 ± 26.42	46.36 ± 27.65	<0.001
<i>Immunological indices</i>				
INF- α (pg/mL)	43.5 ± 15.07	49.04 ± 19.31	52.92 ± 12.22	<0.001
TNF- γ (pg/mL)	40.3 ± 11.28	42.35 ± 17.63	42.71 ± 18.84	<0.001
IL-10 (pg/mL)	48.93 ± 17.39	69.37 ± 13.90	80.60 ± 23.18	0.008
IL-12 (pg/mL)	20.5 ± 21.08	30.81 ± 14.93	27.61 ± 13.09	<0.001
IL-10/TNF- α	1.21 ± 0.05	1.64 ± 0.56	1.89 ± 0.92	0.061
IL-10/INF- γ	1.12 ± 0.23	1.41 ± 0.24	1.52 ± 0.73	0.024

[39–41]. The trend of decreasing platelets with increasing levels of parasitaemia observed in this study has been previously noted for malaria infection due to *P. falciparum* [22,23].

We further observed low value of differential WBC counts among infected malaria patient as compared to uninfected ones. This finding is in direct contrast with report of Ladhani et al. [42] but consistent with many studies [30,22,43]. The association of *P. falciparum* malaria to low WBC count has been suggested to be attributable to sequestration and accelerated destruction of leukocytes [44]. Further studies have shown that the production of proinflammatory cytokines by parasite immunogenic antigen (Glycosylphosphatidylinositol) increase phagocytosis of RBC and WBC, thus causing TNF- α , causing altered hemopoiesis [45,46].

Cytokines are considered to act as double-edged swords in the pathogenesis of malaria, especially in *P. falciparum* infections [47,48]. The high levels of innate immune response in malaria infection is driven by malarial pigment (hemozoin, Hz), and glycosylphosphatidylinositols (GPIs) in the circulating monocytes and neutrophils, and resident macrophages [49,50,44]. In this study, plasma concentrations of TNF- α , IFN- γ , IL-10 and IL-12 increased with increasing parasite density and it is in accordance with the report of Mugweru and his colleagues [51]. However, T-helper cell type 1 cytokines is essential to achieve elimination of infection with less damage to the host. Walther et al. [52] noted that expression of these pro-inflammatory cytokines may have far reaching consequences on disease outcome such as fever, paroxysms, anaemia, cerebral malaria and many other systemic infection symptoms [51,53–55]. It has been postulated that induction of TNF- α delocalizes ferroportin in the reticuloendothelial system that mediates stimulation of macrophage and intestinal iron absorption. This, however, decreases iron absorption and enhances the release of hepcidin to the circulation, which its increased serum level is often associated with severe malaria anaemia [56,57]. Development of Th1 response can be antagonized indirectly by IL-10, which inhibits the production of proinflammatory cytokines. IL-10 induces B-cell proliferation. The inhibition of interferon and TNF- α secretion by IL-10 synthesis has been reported to be important in counteracting the pathological role of macrophages in cerebral malaria [58]. This accounted for high level of this cytokine and its relative smaller ratio with TNF- α in the infected group and it is consistent with many other studies [59,12]. Much consistent evidence based on experimental models and human *P. falciparum* infection suggests increased IL-10 levels may favor parasite multiplication by inhibiting parasite-killing effector mechanisms in humans and mice [60,61].

In conclusion, outcome of this study underscores haematological alterations as the hallmark pathological conditions of malaria infection and their significance in the prediction of severe infection. It also revealed that unregulated interaction of the parasite with host immune response can have important consequences for disease progression. Therefore, it becomes pertinent to incorporate findings of this study in the on-going malaria vaccine and effective therapeutic development.

Acknowledgments

The authors are grateful to the medical directors of children specialist hospital and University of Ilorin Health Centre for their supports. We equally appreciate our participants and our numerous project students (Akande, OA; Afolayan, MA; Aderibigbe, EK; Shuaib, AB) for their supports during sample collection and data analysis. We are indeed very grateful to you all.

References

- [1] Ikekpeazu EJ, Neboh EE, Maduka IC, Nwagbara JJ, Nwobodo MW. Type-2 diabetes mellitus and malaria parasitemia: effect on liver function tests. *Asian J Med Sci* 2010;2:214–7.
- [2] Ogbodo SO, Okeke AC, Obu HA, Shu EN, Chukwurah EF. Nutritional status of parasitemic children from malaria endemic rural communities in eastern Nigeria. *Curr Pediatric Res* 2010;14:131–5.
- [3] World Health Organisation. World malaria Report 2015. <www.who.int/malaria/publications/world-malaria-report>; 2015.
- [4] Lima-Junior JC, Rodrigues-da-Silva RN, Pereira VA, Storer FL, Perceda-Silva DS, Fabris DL, Santos F, Banic DM, de Oliveira-Ferreira J. Cells and mediators of inflammation (C-reactive protein, nitric oxide, platelets and neutrophils) in the acute and convalescent phases of uncomplicated *Plasmodium vivax* and *Plasmodium falciparum* infection. *Mem Inst Oswaldo Cruz Rio de Janeiro* 2012;107:1035–41.
- [5] Mendis K, Sina BJ, Marchesini P, Carter R. The neglected burden of *Plasmodium vivax* malaria. *Am J Trop Med Hyg* 2001;64:97–106.
- [6] Glenister FK, Coppel RL, Cowman AF, Mohandas N, Cooke BM. Contribution of parasite proteins to altered mechanical properties of malaria-infected red blood cells. *Blood* 2002;99:1060–3.
- [7] Shelby JP, White J, Ganesan K, Rathod PK, Chiu DT. A microfluidic model for single-cell capillary obstruction by *Plasmodium falciparum*-infected erythrocytes. *Proc Natl Acad Sci USA* 2003;100:14618–22.
- [8] Dorothee FVC, Andrew JM, Albin F, Irène JV, Marie-Christine A, Giovanna C, Thierry F, Georges EG. Platelet micro particles: a new player in malaria parasite cytoadherence to human brain endothelium. *FASEB J* 2009;23:3449–58.
- [9] Jayakumar SP, Demba S, Geoffrey S, Tamas N, Julie MM. Malaria-induced murine pregnancy failure: Distinct roles for IFN- γ and TNF. *J Immunol* 2009;183:5342–9.
- [10] Srabasti JC, Katie RH, Alister GC. Host response to cytoadherence in *Plasmodium falciparum*. *Biochem Soc Trans* 2008;36:221–8.
- [11] Erdman LK, Finney CA, Liles WC, Kain KC. Inflammatory pathways in malaria infection: TLRs share the stage with other components of innate immunity. *Mol Biochem Parasitol* 2008;162(2):105–11.
- [12] Gonçalves RM, Scopel KKG, Bastos MS, Ferreira MU. Cytokine balance in human malaria: Does *Plasmodium vivax* elicit more inflammatory responses than *Plasmodium falciparum*? *PLoS One* 2012;7:e44394.
- [13] Ing R, Segura M, Thawani N, Tam M, Stevenson MM. Interaction of mouse dendritic cells and malaria-infected erythrocytes: uptake, maturation, and antigen presentation. *J Immunol* 2006;176:441–50.
- [14] Artavanis-Tsakonas K, Riley EM. Innate immune response to malaria: Rapid induction of IFN-gamma from human NK cells by live *Plasmodium falciparum*-infected erythrocytes. *J Immunol* 2002;169:2956–63.
- [15] Cox-Singh J, Jessie H, Sebastian BL, Paul CD, Mohammad Z, Patricia C, Kum TW, Patricia A, Sherif RZ, Balbir S, Sanjeev K. Severe malaria – a case of fatal *Plasmodium knowlesi* infection with post-mortem findings: a case report. *Malar J* 2010;9:10.
- [16] Leowattana W, Tangpukdee N, Thar SK, Nakasiri S, Srivilairit S, Kano S, Wilairatana P, Krudsood S. Changes in platelet count in uncomplicated and severe falciparum malaria. *Southeast Asian J Trop Med Public Health* 2010;41:1035–41.
- [17] Lacerda MVG, Mourão MPG, Coelho HCC, Santos JB. Thrombocytopenia in malaria: who cares? *Mem Inst Oswaldo Cruz* 2011;106:52–63.
- [18] George IO, Ewelike-Ezeani CS. Haematological changes in children with malaria infection in Nigeria. *J Med Med Sci* 2011;2:768–71.
- [19] Bakhubaira S. Hematological parameters in severe complicated *Plasmodium falciparum* malaria among adults in Aden. *Turkish J Hematol* 2013;30:394–9.
- [20] Nmadu PM, Peter E, Alexander P, Koggie AZ, Maikenti JL. The prevalence of malaria in children between the Ages 2–15 visiting Gwarinpa General Hospital life-camp, Abuja, Nigeria. *J Health Sci* 2015;5:47–51.
- [21] Babamale OA, Ugboimoiko US. Status of malaria infection in peri-urban community of North Central Region of Nigeria. *J Bacteriol Parasitol* 2016;7:256.
- [22] Erhart LM, Yingyuen K, Chuanak N, Buathong N, Laoboonchai A, Miller RS, et al. Hematologic and clinical indices of malaria in a semi-immune population of western Thailand. *Am J Trop Med Hyg* 2004;70:8–14.
- [23] Saravu K, Docherla M, Vasudev A, Shastri BA. Thrombocytopenia in vivax and falciparum malaria: An observational study of 131 patients in Karnataka, India. *Ann Trop Med Parasitol* 2011;105:593–8.
- [24] Breman JG, Egan A, Keusch GT. The intolerable burden of malaria: a new look at the numbers. *Am J Trop Med Hygiene* 2001;64:4–7.
- [25] WHO. WHO world malaria report 2005. Geneva: World Health Organization/ United Nations Children's Fund; 2005. <http://rbm.who.int/wmr2005/>.
- [26] Maina RN, Walsh D, Gaddy C, Hongo G, Waitumbi J, Otieno L, Jones D, Ogutu RB. Impact of *Plasmodium falciparum* infection on haematological parameters in children living in Western Kenya. *Malar J* 2010;9:54. doi: <http://dx.doi.org/10.1186/1475-2875-9-S3-S4>.
- [27] Chandra S, Chandra H. Role of haematological parameters as an indicator of acute malarial infection in Uttarakhand State of India. *Mediterr J Hematol Infect Dis* 2013;5:e2013009.
- [28] Mohamed A, Bushra S, Gamal A, Ahmed M, Ali A. Parasitaemia and Its Relation to Hematological Parameters and Liver Function among Patients Malaria in Abs, Hajjah, Northwest Yemen. *Interdiscip Perspect Infect Dis* 2016;5. doi: <http://dx.doi.org/10.1155/2016/595439>.
- [29] Ekvall H. Malaria and anemia. *Curr Opin Hematol* 2003;10:108–14.
- [30] Lee HK, Lim J, Kim M, Lee S, Oh EJ, Lee J, Oh J, Kim Y, Han K, Lee EJ, Kang CS, Kim BK. Immunological alterations associated with *Plasmodium vivax* malaria in South Korea. *Ann Trop Med Parasitol* 2001;95:31–9.
- [31] Casals-Pascual C, Kai O, Newton CR, Peshu N, Roberts DJ. Thrombocytopenia in falciparum malaria is associated with high concentrations of IL-10. *Am J Trop Med Hyg* 2006;75:434–6.

- [32] Oh MD, Shia H, Shin Kim U, Lee S, Kim N, et al. Clinical features of vivax malaria. *Am J Trop Med Hyg* 2001;65:143–6.
- [33] Taha K, El-Dein Z, Idrees M, Makboul G, Ghassan B. Hematological changes in malaria: relation to plasmodium species. *Kuwait Med J* 2007;39:262–7.
- [34] Garba N, Danladi SB, Muhammad A. Determination of some haematological parameters in malaria infected subjects in Usmanu Danfodiyo University Teaching Hospital (Uduth) Sokoto, Nigeria. *Bayero J Pure Appl Sci* 2015;8:80–3.
- [35] Pavithran K. Haematological changes in malaria. *Southeast Asian J Trop Med Public Health* 2005;19:25–33.
- [36] Abdul Rauf M, Afsar S. Thrombocytopenia in hospitalized malaria patients. *Pak J Med Sci* 2006;22:141–3.
- [37] Crabbe G, Von Poucke M, Cantinieux B. Artefactually-normal automated platelet counts due to malaria infected RBC. *Clin Lab Haematol* 2002;24:179–82.
- [38] Malik AM, Zaffar N, Ali N, Malik AM, Khan R. Hematological findings and endemicity of malaria In Gadap Region. *J Coll Physicians Surg* 2010;20:112–6.
- [39] Faseela TS, Ronald AR, Anita KB, Chaithra SM, Yashwanth R. Diagnostic value of platelet count in malaria. *J Clin and Diagn Res* 2011;5:464–6.
- [40] Morrell CN, Aggrey AA, Chapman LM, Modjeski KL. Emerging roles for platelets as immune and inflammatory cells. *Blood* 2013;123:2759–67.
- [41] Anstey NM, Weinberg JB, Hassanali MY, Mwaikambo ED, Manyenga D, Misukonis MA, et al. Nitric oxide in Tanzanian children with malaria: Inverse relationship between malaria severity and nitric oxide production/nitric oxide synthase type 2 expression. *J Exp Med* 1996;184:557–67.
- [42] Yeo TW, Lampah DA, Gitawati R, Tjitra E, Kenangalem E, McNeil YR, et al. Impaired nitric oxide bioavailability and L-arginine reversible endothelial dysfunction in adults with falciparum malaria. *J Exp Med* 2007;204:2693–704.
- [43] Sumbele IUN, Nkuo-Akenji T, Samje M, Ndzeize T, Ngwa EM, Titanji VPK. Haematological changes and recovery associated with treated and untreated *P. falciparum* infection in children in the Mount Cameroon Region. *J Clin Med Res* 2010;2:143–51.
- [44] Ladhani S, Lowe B, Cole AO, Kowuondo K, Newton CR. Changes in white blood cells and platelets in children with falciparum malaria: relationship to disease outcome. *Br J Haematol* 2002;119:839–47.
- [45] McKenzie FE, Prudhomme WA, Magill AJ, et al. White blood cell counts and malaria. *J Infect Dis* 2005;192:323–30.
- [46] Tangpukdee N, Yew HS, Krudsood S, et al. Dynamic changes in white blood cell counts in uncomplicated *Plasmodium falciparum* and *P. vivax* malaria. *Parasitol Int* 2008;57:490–4.
- [47] Miller AR, Suttles J, Stout RD. Cytokine priming reduces dependence on TNF-R2 for TNF- α -mediated induction of macrophage nitric oxide generation. *J Interferon Cytokine Res* 1996;16:1055–63.
- [48] Jason J, Archibald LK, Nwanyanwu OC, Bell M, Buchanan I, Larned J, Kazembe PN, Dobbie H, Parekh B, Byrd MG, Eick A, Han A, Jarvis WR. Cytokines and malaria parasitemia. *Clin Immunol* 2001;100:208–18.
- [49] Angulo I, Fresno M. Cytokines in the pathogenesis of and protection against malaria. *Clin Diagn Lab Immunol* 2002;9:1145–52.
- [50] Gimenez F, Barraud de Lagerie S, Fernandez C, Pino P, Mazier D. Tumor necrosis factor alpha in the pathogenesis of cerebral malaria. *Cell Mol Life Sci* 2003;60:1623–35.
- [51] Tchinda VH, Tadem AD, Tako EA, Tene G, Fogako J, Nyonglema P, Sama G, Zhou A, Leke RG. Severe malaria in Cameroonian children: Correlation between plasma levels of three soluble inducible adhesion molecules and TNF- α . *Acta Trop* 2007;102:20–8.
- [52] Waihenya J, Kimani R, Ng'ang'a Z, Matiru V, Muregi Francis W. Cytokine levels associated with experimental malaria pathology during *Plasmodium Berghei* ANKA infection in a mouse model. *Clin Immunol Immunopathol Res* 2013;5:1–8.
- [53] Walther M, Thompson FM, Dunachie S, Keating S, Todryk S, Berthoud T. Safety, immunogenicity and efficacy of prime-boost immunization with recombinant pox viruses FP9 and modified vaccinia virus Ankara encoding the full length *Plasmodium falciparum* circumsporozoite protein. *Infect Immun* 2006;74:2706–16.
- [54] Karunaweera ND, Wijesekera SK, Wanasekera D, Mendis KN, Carter R. The paroxysm of *Plasmodium vivax* malaria. *Trends Parasitol* 2003;19:188–93.
- [55] Armah H, Wired EK, Dodoo AK, Adjei AA, Tettey Y, Gyasi R. Cytokines and adhesion molecules expression in the brain in human cerebral malaria. *Int J Environ Res Public Health* 2005;2:123–31.
- [56] Nweneka CV, Doherty CP, Cox S, Prentice A. Iron delocalization in the pathogenesis of malarial anaemia. *Trans R Soc Trop Med Hyg* 2010;104:175–84.
- [57] de Mast Q, Syafruddin D, Keijmel S, Riekerink TO, Deky O, Asih PB, Swinkels DW, van der Ven AJ. Increased serum hepcidin and alterations in blood iron parameters associated with asymptomatic *P. falciparum* and *P. vivax* malaria. *Haematologica* 2010;95:1068–74.
- [58] Lucia M, Salvatore M. The immune response to *Plasmodium falciparum* malaria. *Lancet Infect Dis* 2002;2:472–8.
- [59] Bueno LL, Moraes CG, Araújo FF, Gomes JA, Corrêa-Oliveira R, Soares IS, et al. *Plasmodium vivax*: induction of CD4+CD25+FoxP3+ Regulatory T Cells during infection are directly associated with level of circulating parasites. *PLoS One* 2010;5:e9623.
- [60] Hugosson E, Montgomery SM, Premji Z, Troye-Blomberg M, Troye-Blomberg A, Björkman A. Higher IL-10 levels are associated with less effective clearance of *Plasmodium falciparum* parasites. *Parasite Immunol* 2004;26:111–7.
- [61] Couper KN, Blount DG, Wilson MS, Hafalla JC, Belkaid Y. IL-10 from CD4+CD25-Foxp3-CD127- adaptive regulatory T cells modulates parasite clearance and pathology during malaria infection. *PLoS Pathog* 2008;4:e1000004.