

Partial Traces on Additive Categories

Naohiko Hoshino^{1,2}

*Research Institute for Mathematical Sciences
Kyoto University
Kyoto, Japan*

Abstract

In this paper, we study partial traces on additive categories. Haghverdi and Scott introduced partially traced symmetric monoidal categories generalizing traced symmetric monoidal categories given by Joyal, Street and Verity. The original example of a partial trace is given in terms of the execution formula on the category of vector spaces and linear functions. Malherbe, Scott and Selinger gave another example of a partial trace on the category of vector spaces, and they observed that we can define these two partial traces on arbitrary additive categories. A natural question is: what kind of partial traces does the category of vector spaces have? We give a (partial) answer to this question. Our main result is: every abelian category has a largest partial trace. Here, “largest” means that every partial trace on the abelian category is obtained by restricting the domain of the largest partial trace. As a corollary, we show that the partial trace given by Malherbe, Scott and Selinger is the largest partial trace on the category of vector spaces.

Keywords: partial trace, execution formula, kernel-image trace, Geometry of Interaction, additive category, abelian category.

1 Introduction

In computer science, traced monoidal categories [9] and their variants are fundamental algebraic structures in the study of categorical semantics for cyclic computation. For example, a trace operator on a cartesian category is a parametrized fixed point operator [7], and a trace operator on a cocartesian category models iteration. Traced monoidal categories also appear in a categorical framework for Girard’s Geometry of Interaction (GoI) [4,1] where trace operators capture interactive communication between automata.

In [5], Haghverdi and Scott introduced partially traced symmetric monoidal categories generalizing the notion of trace in order to give a categorical framework for the original GoI based on vector spaces. Their generalization provides many examples of partially traced symmetric monoidal categories that have not emerged

¹ The author is supported by the JSPS-INRIA Bilateral Joint Research Project CRECOGI and by JSPS KAKENHI Grant Number JP26730004

² Email: naophiko@kyoto-u.ac.jp

in studies of categorical semantics for lambda calculi. Typically, we need to consider partial traces when the underlying symmetric monoidal category can only model converging iteration process. For example, Haghverdi and Scott showed that the original execution formula

$$\text{Ex} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = A + B(I - D)^{-1}C \quad (1)$$

given in [4] is a partial trace on the symmetric monoidal category of vector spaces with the biproducts as the monoidal products, and they constructed a denotational semantics for linear logic using the execution formula. Roughly speaking, in their work, the execution formula computes the adjacent matrix associated to the normal form of a given proof Π of linear logic. The block matrices A, B, C and D in (1) are adjacent matrices corresponding to certain fragments of Π , and when D is nilpotent, (1) is equal to $A + \sum_{n \in \mathbb{N}} BD^n C$, which is the adjacent matrix associated to the normal form of Π . Since the inverse $(I - D)^{-1}$ may not be defined, the execution formula is partially defined. In [13], Malherbe, Scott and Selinger observed that we can generalize construction of the partial trace by Haghverdi and Scott to any additive category, and they also showed that every additive category has another partial trace called the kernel-image trace. A question about these observations is: what kind of partial traces does an additive (or an abelian) category have?

The goal of this paper is to give an answer to this question. We study partial traces on additive categories and abelian categories, and we prove the following results.

- In every additive category, the partial trace given by Haghverdi and Scott is the largest partial trace that satisfies strong naturality. (Theorem 3.3)
- Every abelian category \mathcal{C} has a largest partial trace that is given by restricting the canonical total trace on the category consisting of objects in \mathcal{C} and relations between them. We also show that the largest partial trace is a natural extension of the kernel-image trace on \mathcal{C} . (Theorem 4.5 and Theorem 4.6)
- If an abelian category \mathcal{C} is semisimple, then the largest partial trace on \mathcal{C} coincides with the kernel-image trace. (Corollary 4.8)

The second result means that any abelian category has essentially one partial trace; all partial traces on an abelian category are just restrictions of the largest partial trace. It follows from our results that the category of finite dimensional vector spaces equipped with the kernel-image trace is a traced symmetric monoidal subcategory of the category of finite dimensional vector spaces and relations between them. This is an answer to the following question posed in [13]:

“One question that we did not answer is whether specific partially traced categories can be embedded in totally traced categories in a natural way. For example, the category of finite dimensional vector spaces, with the biproduct \oplus as the tensor, can be equipped with a natural partial trace in several ways. By our proof, it

follows that it can be faithfully embedded in a totally traced category. However, we do not know any concrete natural description of such a totally traced category (i.e., other than the free one constructed in our proof)."

Our motivation to study (partial) trace comes from importance of (partial) traces in categorical semantics for recursive computation and GoI, and we are interested in partial traces on additive categories and abelian categories because of use of C^* -algebra in the first GoI. However, our results are just answers to purely technical questions on additive categories and abelian categories, and at this point, we do not have any practical application of our results to these research areas. To find practical application of our results is a future work.

The structure of this paper is as follows. In Section 2, we recall the notions of partially traced symmetric monoidal category, additive category and abelian category. We also give some examples and basic properties of these categories. In Section 3, we give examples of partial traces on additive categories and show that the partial trace given by Haghverdi and Scott is the largest partial trace among partial traces satisfying strong naturality. In Section 4, we show that every abelian category has a largest partial trace and when the abelian category is semisimple, the kernel-image trace is the largest partial trace. For proofs of some statements, see [8].

2 Categorical preliminaries

2.1 Symmetric monoidal category and symmetric monoidal functor

A *symmetric monoidal category* consists of a category \mathcal{C} equipped with a unit object I and a monoidal product $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and natural isomorphisms $\lambda_X: I \otimes X \cong X$, $\rho_X: X \otimes I \cong X$, $\alpha_{X,Y,Z}: (X \otimes Y) \otimes Z \cong X \otimes (Y \otimes Z)$ and $\sigma_{X,Y}: X \otimes Y \cong Y \otimes X$ subject to some coherence conditions (see [12,2] for example). We simply write $(\mathcal{C}, I, \otimes)$ or \mathcal{C} for symmetric monoidal categories when we can infer other data from the context. In this paper, we often regard a cartesian category \mathcal{C} as a symmetric monoidal category $(\mathcal{C}, 1, \times)$ where 1 is the terminal object and $X \times Y$ is the product of X and Y .

Let $(\mathcal{C}, I, \otimes)$ and (\mathcal{D}, J, \odot) be symmetric monoidal categories. A *symmetric monoidal functor* (F, n, m) from $(\mathcal{C}, I, \otimes)$ to (\mathcal{D}, J, \odot) consists of a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ with an arrow $n: J \rightarrow FI$ and a natural transformation $m_{X,Y}: FX \odot FY \rightarrow F(X \otimes Y)$ subject to some coherence conditions (see [12,2] for example). A symmetric monoidal functor (F, n, m) is *strong* when n and $m_{X,Y}$ are isomorphisms. We write $F: \mathcal{C} \rightarrow \mathcal{D}$ to denote a (strong) symmetric monoidal functor when we can infer n and $m_{X,Y}$ from the context.

2.2 Partial trace

We recall the definition of a partial trace introduced by Haghverdi and Scott in [5]. We prepare several notations. Given partially defined expressions \mathcal{E} and \mathcal{F} , we write $\mathcal{E} \preceq \mathcal{F}$ when \mathcal{E} is not defined or both \mathcal{E} and \mathcal{F} are defined and they are

the same. For example, we have $\sum_{n \in \mathbb{N}} x^n \preceq \frac{1}{1-x}$ for all $x \in [0, \infty)$ because the left hand side is defined if and only if $0 \leq x < 1$, and we have $\sum_{n \in \mathbb{N}} x^n = \frac{1}{1-x}$ for all $0 \leq x < 1$. We write $\mathcal{E} \simeq \mathcal{F}$ when both $\mathcal{E} \preceq \mathcal{F}$ and $\mathcal{F} \preceq \mathcal{E}$ are true.

Definition 2.1 Let $(\mathcal{C}, I, \otimes, \lambda, \rho, \alpha, \sigma)$ be a symmetric monoidal category. A *partial trace* \mathbf{tr} on \mathcal{C} is a family of partial functions

$$\mathbf{tr}_{X,Y}^Z: \mathcal{C}(X \otimes Z, Y \otimes Z) \rightarrow \mathcal{C}(X, Y) \quad (X, Y, Z \in \mathcal{C})$$

subject to the following conditions:

- (Naturality) For all $f: X \otimes Z \rightarrow Y \otimes Z$, $g: U \rightarrow X$ and $h: Y \rightarrow V$,

$$g \circ \mathbf{tr}_{X,Y}^Z(f) \circ h \preceq \mathbf{tr}_{U,V}^Z((g \otimes Z) \circ f \circ (h \otimes Z)).$$

- (Sliding) For all $f: X \otimes W \rightarrow Y \otimes Z$ and $g: Z \rightarrow W$,

$$\mathbf{tr}_{X,Y}^W(f \circ (Y \otimes g)) \simeq \mathbf{tr}_{X,Y}^Z((X \otimes g) \circ f).$$

- (Superposing) For all $f: X \otimes Z \rightarrow Y \otimes Z$,

$$W \otimes \mathbf{tr}_{X,Y}^Z(f) \preceq \mathbf{tr}_{W \otimes X, W \otimes Y}^Z(\alpha_{W,X,Z} \circ (W \otimes f) \circ \alpha_{W,Y,Z}^{-1}).$$


- (Vanishing I) For all $f: X \otimes I \rightarrow Y \otimes I$, $\mathbf{tr}_{X,Y}^I(f)$ is defined and is equal to $\rho_X^{-1} \circ f \circ \rho_Y$.
- (Vanishing II) For all $f: (X \otimes W) \otimes Z \rightarrow (Y \otimes W) \otimes Z$, if $\mathbf{tr}_{X \otimes W, Y \otimes W}^Z(f)$ is defined, then

$$\mathbf{tr}_{X,Y}^W(\mathbf{tr}_{X \otimes W, Y \otimes W}^Z(f)) \simeq \mathbf{tr}_{X,Y}^{W \otimes Z}(\alpha_{X,W,Z}^{-1} \circ f \circ \alpha_{Y,W,Z}).$$

- (Yanking) For all $X \in \mathcal{C}$, $\mathbf{tr}_{X,X}^X(\sigma_{X,X})$ is defined and is equal to id_X .

A *partially traced symmetric monoidal category* is a symmetric monoidal category equipped with a partial trace. A *total trace* is a partial trace that is totally defined, and a *totally traced symmetric monoidal category* is a symmetric monoidal category equipped with a total trace. We note that the notion of totally traced symmetric monoidal category coincides with the notion of *traced (symmetric) monoidal category* given in [9].

We often present $\mathbf{tr}_{X,Y}^Z(f): X \rightarrow Y$ as the following feedback loop:



For many totally traced symmetric monoidal categories in computer science, this feedback loop nicely fits in our intuition. As an example, let $(\mathbf{Pfn}, \emptyset, \oplus)$ be the symmetric monoidal category of sets and partial functions whose monoidal product


is the disjoint sum

$$X \oplus Y = \{(0, x) \mid x \in X\} \cup \{(1, y) \mid y \in Y\}.$$


The symmetric monoidal category $(\mathbf{Pfn}, \emptyset, \oplus)$ has a total trace **iter**: for a partial function $f: X \oplus Z \rightarrow Y \oplus Z$, the trace $\mathbf{iter}_{X,Y}^Z(f): X \rightarrow Y$ is given by

$$\mathbf{iter}_{X,Y}^Z(f)(x) = y \iff \text{there is a finite sequence } z_1, \dots, z_n \in Z \text{ such that} \\ f(0, x) = (1, z_1) \text{ and } f(1, z_1) = (1, z_2) \text{ and } \dots \text{ and } f(1, z_n) = (0, y).$$

The definition of $\mathbf{iter}_{X,Y}^Z(f)(x)$ means that for $x \in X$, the partial trace $\mathbf{iter}_{X,Y}^Z(f)(x)$ is defined and is equal to $y \in Y$ if and only if there is a finitely many loops like the following dotted line:



This diagrammatic presentation helps us to convince that **iter** is a total trace. For example, the sliding axiom holds because the difference between the following two diagrams



is just how we arrange boxes and wires, and the arrangement has nothing to do with how data flow along wires.

We can construct various partial traces by applying the following theorem to already known partial traces.

Theorem 2.2 ([13, Proposition 3.20]) *Let $(\mathcal{C}, I, \otimes)$ and (\mathcal{D}, J, \odot) be symmetric monoidal categories, and let \mathbf{tr} be a partial trace on \mathcal{D} . If there is a faithful strong symmetric monoidal functor $(F, n, m): (\mathcal{C}, I, \otimes) \rightarrow (\mathcal{D}, J, \odot)$, then \mathcal{C} has a partial trace \mathbf{tr} given by*

$$\mathbf{tr}_{X,Y}^Z(f) = \begin{cases} g: X \rightarrow Y, & \text{if } \mathbf{tr}_{FX,FY}^{FZ}(m_{X,Z} \circ Ff \circ m_{Y,Z}^{-1}) \text{ is defined and equals } Fg, \\ \text{undefined,} & \text{otherwise.} \end{cases}$$

This is well-defined since F is faithful.

Example 2.3 Let $(\mathbf{Set}, \emptyset, \oplus)$ be the symmetric monoidal category of sets and functions whose monoidal product is given by the disjoint sum of sets. Since there is a trivial inclusion functor $J: \mathbf{Set} \rightarrow \mathbf{Pfn}$, we have a partial trace on \mathbf{Set} that is given by pulling back **iter** along J . Concretely, the partial trace $\mathbf{iter}_{X,Y}^Z$ on \mathbf{Set} is given

by

$$\mathbf{iter}_{X,Y}^Z(f: X \oplus Z \rightarrow Y \oplus Z) = \begin{cases} \mathbf{iter}_{X,Y}^Z(f), & \text{if } \mathbf{iter}_{X,Y}^Z(f) \text{ is totally defined,} \\ \text{undefined,} & \text{otherwise.} \end{cases}$$

Example 2.4 Let $(\mathbf{Set}, \{0\}, \times)$ be the symmetric monoidal category of sets and functions whose monoidal product is given by the cartesian product of sets, and let $(\mathbf{Rel}, \{0\}, \times)$ be the symmetric monoidal category of sets and relations whose monoidal product is also given by the cartesian product of sets. The symmetric monoidal category $(\mathbf{Rel}, \{0\}, \times)$ has a total trace given by

$$\mathbf{tr}_{X,Y}^Z(r) = \{(x, y) \in X \times Y \mid ((x, z), (y, z)) \in r \text{ for some } z \in Z\}.$$

By pulling back the total trace \mathbf{tr} along the obvious inclusion functor $\mathbf{Set} \rightarrow \mathbf{Rel}$, we obtain a partial functor \mathbf{tr} on \mathbf{Set} . To be concrete, given a function $f: X \times Z \rightarrow Y \times Z$, the partial trace $\mathbf{tr}_{X,Y}^Z(f)$ is defined and is equal to a function $g: X \rightarrow Y$ if and only if

$$\{(x, y) \in X \times Y \mid f(x, z) = (y, z) \text{ for some } z \in Z\}$$

is the graph relation of g . In the next section, we will generalize this construction to regular categories.

The next example is about existence of the least partial trace.

Example 2.5 For partial traces \mathbf{tr}_1 and \mathbf{tr}_2 on a symmetric monoidal category $(\mathcal{C}, I, \otimes)$, we write $\mathbf{tr}_1 \leq \mathbf{tr}_2$ when we have $(\mathbf{tr}_1)_{X,Y}^Z(f) \preceq (\mathbf{tr}_2)_{X,Y}^Z(f)$ for all arrows $f: X \otimes Z \rightarrow Y \otimes Z$ in \mathcal{C} . This is a partial order on the set of all partial traces on \mathcal{C} . With respect to this partial order, every non-empty family of partial traces $\{\mathbf{tr}_\lambda\}_{\lambda \in \Lambda}$ on \mathcal{C} has a greatest lower bound \mathbf{tr} given by

$$\begin{aligned} \mathbf{tr}_{X,Y}^Z(f) \text{ is defined and equals } g &\iff \\ \text{for all } \lambda \in \Lambda, (\mathbf{tr}_\lambda)_{X,Y}^Z(f) \text{ is defined and equals } g. \end{aligned}$$

In particular, if \mathcal{C} has a partial trace, then the meet of all partial traces on \mathcal{C} exists and is the least partial trace on \mathcal{C} . We note that the least partial trace on \mathcal{C} is not the everywhere undefined operator because every partial trace must be defined at $\sigma_{X,X}$. It is not easy to find concrete description of the least partial trace.

2.3 Regular category

In this paper, a partial trace on a regular category plays an important role. In a regular category \mathcal{C} , we can consider relations between objects in \mathcal{C} , and we can define a category $\mathbf{Rel}(\mathcal{C})$ consisting of objects in \mathcal{C} and relations between them. It is well-known that $\mathbf{Rel}(\mathcal{C})$ has the structure of a totally traced symmetric monoidal category. Then we can define a partial trace on \mathcal{C} by pulling back the total trace on $\mathbf{Rel}(\mathcal{C})$. Below, we precisely describe this construction. We first recall the definition of a regular category.

Definition 2.6 A *regular category* is a category with finite limits such that

- every arrow $f: X \rightarrow Y$ has a factorization $f = e \circ m$ with a monomorphism $m: Z \rightarrow Y$ and a regular epimorphism $e: X \rightarrow Z$,
- the pullback of a regular epimorphism along any arrow is a regular epimorphism.

Here, an epimorphism $e: X \rightarrow Y$ is *regular* if and only if e is a coequalizer of a pair of arrows $u, v: Z \rightrightarrows X$. Below, we write $e: X \twoheadrightarrow Y$ when e is a regular epimorphism from X to Y , and we write $m: X \rightarrowtail Y$ when m is a monomorphism from X to Y .

For example, the category **Set** of sets and function is a regular category where the image factorization of a function $f: X \rightarrow Y$ from a set X to a set Y is:

$$X \xrightarrow{f} \{y \in Y \mid y = f(x) \text{ for some } x \in X\} \hookrightarrow Y.$$

It is known that regular-epi/mono factorization is unique up-to isomorphism. This means that if an arrow $f: X \rightarrow Y$ in a category \mathcal{C} has two regular-epi/mono factorizations

$$f = X \xrightarrow{e} Z \rightarrowtail Y, \quad f = X \xrightarrow{e'} Z' \rightarrowtail Y,$$

then there is an isomorphism $\theta: Z \rightarrow Z'$ such that $e \circ \theta = e'$ and $\theta \circ m' = m$.

When \mathcal{C} is a regular category, regular-epi/mono factorization enables us to define a totally traced symmetric monoidal category **Rel**(\mathcal{C}): objects in **Rel**(\mathcal{C}) are objects in \mathcal{C} , and arrows from X to Y are (equivalence classes of) subobjects $r: R \rightarrowtail X \times Y$. This is a generalization of the category **Rel** of sets and relations. In fact, we have **Rel**(**Set**) \cong **Rel**. The identity on X is the diagonal arrow $\delta_X: X \rightarrowtail X \times X$, and the composition $r * s: T \rightarrowtail X \times Z$ of relations $r = \langle r_1, r_2 \rangle: R \rightarrowtail X \times Y$ and $s = \langle s_1, s_2 \rangle: S \rightarrowtail Y \times Z$ is defined to be the mono-part of the regular-epi/mono factorization of $\langle p_1 \circ r_1, p_2 \circ s_2 \rangle$:

$$\begin{array}{ccc} & \langle p_1 \circ r_1, p_2 \circ s_2 \rangle & \\ & \curvearrowright & \\ U & \xrightarrow{\quad} T \rightarrowtail X \times Z & \end{array} \quad \text{where} \quad \begin{array}{ccc} U & \xrightarrow{p_2} & S \\ p_1 \downarrow & \text{p.b} & \downarrow s_1 \\ R & \xrightarrow{r_2} & Y \end{array}.$$

Here, $\langle -, - \rangle$ is the tuple of arrows. Well-definedness of the composition of arrows in **Rel**(\mathcal{C}) follows from uniqueness of regular-epi/mono factorization. The unit object of **Rel**(\mathcal{C}) is the terminal object $1 \in \mathcal{C}$, and the monoidal product of objects X and Y in **Rel**(\mathcal{C}) is the product $X \times Y$. The monoidal product of relations $r: R \rightarrowtail X \times Y$ and $s: S \rightarrowtail Y \times Z$ is given by

$$r \otimes s = R \times S \xrightarrow{r \times s} (X \times Y) \times (Y \times Z) \xrightarrow{\cong} (X \times Z) \times (Y \times Y).$$

For a relation $r: R \rightarrowtail (X \times Z) \times (Y \times Z)$, the trace $\mathbf{tr}_{X,Y}^Z(r)$ is the mono-part of the regular-epi/mono factorization of the vertical arrow from P to $X \times Y$ in the

following diagram:

$$\begin{array}{ccccc}
 P & \xlongequal{\quad} & P & \xrightarrow{\quad} & Z \\
 \downarrow & & \downarrow & \text{p.b.} & \downarrow \delta_Z \\
 S & & R & \xrightarrow{r \circ (\pi'_{X,Z} \times \pi'_{Y,Z})} & Z \times Z \\
 \text{tr}_{X,Y}^Z(r) \downarrow & & \downarrow r \circ (\pi_{X,Z} \times \pi_{Y,Z}) & & \\
 X \times Y & \xlongequal{\quad} & X \times Y & &
 \end{array}$$

To clarify intuition of $\text{tr}_{X,Y}^Z(r)$, we concretely describe the above diagram assuming that \mathcal{C} is the category of sets and functions. First, the set R is the graph of r :

$$R = \{((x, z), (y, z')) \in (X \times Z) \times (Y \times Z) \mid ((x, z), (y, z')) \in r\},$$

and two arrows going out from R are projections. The set P is a subset of R given by

$$P = \{((x, z), (y, z)) \in (X \times Z) \times (Y \times Z) \mid ((x, z), (y, z)) \in r\}.$$

Therefore, the image S of the arrow from P to $X \times Y$ is

$$\{(x, y) \in X \times Y \mid ((x, z), (y, z)) \in r \text{ for some } z \in Z\}.$$

We note that we can derive this total trace from the compact closed structure of $\mathbf{Rel}(\mathbf{Set})$, and this is true for arbitrary regular category: the total trace $\text{tr}_{X,Y}^Z$ on $\mathbf{Rel}(\mathcal{C})$ is derived from a *compact closed structure* of $\mathbf{Rel}(\mathcal{C})$ and is called the *canonical trace* in [9]. (We can check that $\mathbf{Rel}(\mathcal{C})$ is a compact closed category by following the idea of the proof of \mathbf{Rel} being compact closed.) For relationship between compact closed categories and total traces, see [9,14]. It is tedious but doable to directly check that tr is a total trace on $\mathbf{Rel}(\mathcal{C})$.

Now, we can apply Theorem 2.2 to the totally traced symmetric monoidal category $(\mathbf{Rel}(\mathcal{C}), 1, \times, \text{tr})$.

Example 2.7 Let \mathcal{C} be a regular category, and let $H_{\mathcal{C}}$ be a faithful strong symmetric monoidal functor $H_{\mathcal{C}}: \mathcal{C} \rightarrow \mathbf{Rel}(\mathcal{C})$ given by

$$H_{\mathcal{C}}(X) = X, \quad H_{\mathcal{C}}(f: X \rightarrow Y) = \langle \text{id}_X, f \rangle: X \rightarrow X \times Y.$$

As we have observed, $\mathbf{Rel}(\mathcal{C})$ has a total trace. Therefore, it follows from Theorem 2.2 that $H_{\mathcal{C}}$ gives rise to a partial trace on \mathcal{C} . We write $\text{rel}_{\mathcal{C}}$ for this partial trace on \mathcal{C} . Concretely, for an arrow $f: X \times Z \rightarrow Y \times Z$ in \mathcal{C} , the partial trace $(\text{rel}_{\mathcal{C}})_{X,Y}^Z(f)$ is defined if and only if there exists $g: X \rightarrow Y$ in \mathcal{C} such that the mono-part of the regular-epi/mono factorization of $p \circ \langle \pi_{X,Z}, f \circ \pi_{Y,Z} \rangle$ is equal to $\langle \text{id}_X, g \rangle$:

$$P \xrightarrow{p} X \times Z \xrightarrow{\langle \pi_{X,Z}, f \circ \pi_{Y,Z} \rangle} X \times Y = P \twoheadrightarrow X \xrightarrow{\langle \text{id}_X, g \rangle} X \times Y$$

where $p: P \rightarrow X \times Z$ is the pullback of δ_Z along $\langle \pi'_{X,Z}, f \circ \pi'_{Y,Z} \rangle$:

$$\begin{array}{ccc} P & \xrightarrow{\quad} & Z \\ p \downarrow & \text{p.b} & \downarrow \delta_Z \\ X \times Z & \xrightarrow{\langle \pi'_{X,Z}, f \circ \pi'_{Y,Z} \rangle} & Z \times Z \end{array} ,$$

and when $(\mathbf{rel}_{\mathcal{C}})_{X,Y}^Z(f)$ is defined, $(\mathbf{rel}_{\mathcal{C}})_{X,Y}^Z(f)$ is equal to g .

2.4 Additive categories and abelian categories

We recall the notions of an additive category and an abelian category, and we give their basic properties. For more details, see [12,3].

Definition 2.8 A *pre-additive category* \mathcal{C} is a category such that every hom-set has the structure of an abelian group and the composition is bilinear. We write $0_{X,Y}: X \rightarrow Y$ for the unit of $\mathcal{C}(X, Y)$ and $f + g$ for the addition of $f, g: X \rightarrow Y$. An *additive category* is a pre-additive category that has finite coproducts.

The category **Ab** of abelian groups and homomorphisms is an additive category. When R is a ring, the category **Mod** $_R$ of left R -modules and homomorphisms is also an additive category. In particular, the category **Vect** $_K$ of vector spaces over a field K and linear maps is an additive category. We denote the subcategory of **Vect** $_K$ consisting of finite dimensional vector spaces by **fdVect** $_K$, which is an additive category.

When \mathcal{C} is an additive category, \mathcal{C} has finite products and those coincide with finite coproducts: the initial object 0 is the terminal object, and the coproduct $X \oplus Y$ with

$$\pi_{X,Y} = [\text{id}_X, 0_{Y,X}]: X \oplus Y \rightarrow X, \quad \pi'_{X,Y} = [0_{X,Y}, \text{id}_Y]: X \oplus Y \rightarrow Y \quad (2)$$

is the product of X and Y where $[-, -]$ is the cotuple of arrows. The diagonal arrow $\delta_X: X \rightarrow X \oplus X$ is given by $\iota_{X,X} + \iota'_{X,X}$ where $\iota_{X,Y}: X \rightarrow X \oplus Y$ and $\iota'_{X,Y}: Y \rightarrow X \oplus Y$ are injections. By the definition of projections, we have

$$\iota_{X,Y} \circ \pi_{X,Y} = \text{id}_X \quad \iota'_{X,Y} \circ \pi'_{X,Y} = \text{id}_Y \quad \iota_{X,Y} \circ \pi'_{X,Y} = 0_{X,Y} \quad \iota'_{X,Y} \circ \pi_{X,Y} = 0_{Y,X}$$

and $\pi_{X,Y} \circ \iota_{X,Y} + \pi'_{X,Y} \circ \iota'_{X,Y} = \text{id}_{X \oplus Y}$. The codiagonal arrow $\gamma_X: X \oplus X \rightarrow X$ is equal to $\pi_{X,X} + \pi'_{X,X}$. The addition $f + g: X \rightarrow Y$ is equal to $\delta_X \circ (f \oplus g) \circ \gamma_Y$. It follows from coincidence of finite coproducts and finite products that every arrow $f: X \oplus Y \rightarrow Z \oplus W$ in \mathcal{C} is uniquely decomposed into a matrix of arrows

$$\begin{pmatrix} f_{XZ}: X \rightarrow Z & f_{XW}: X \rightarrow W \\ f_{YZ}: Y \rightarrow Z & f_{YW}: Y \rightarrow W \end{pmatrix} = \begin{pmatrix} \iota_{X,Y} \circ f \circ \pi_{Z,W} & \iota_{X,Y} \circ f \circ \pi'_{Z,W} \\ \iota'_{X,Y} \circ f \circ \pi_{Z,W} & \iota'_{X,Y} \circ f \circ \pi'_{Z,W} \end{pmatrix}$$

called the matrix decomposition of f . This decomposition is unique because we can recover f as follows:

$$f = \pi_{X,Y} \circ f_{XZ} \circ \iota_{Z,W} + \pi_{X,Y} \circ f_{XW} \circ \iota'_{Z,W} + \pi'_{X,Y} \circ f_{YZ} \circ \iota_{Z,W} + \pi'_{X,Y} \circ f_{YW} \circ \iota'_{Z,W}.$$

The composition of arrows in \mathcal{C} is compatible with matrix multiplication, namely, the matrix decomposition of $f \circ g: X \oplus Y \rightarrow Z \oplus W \rightarrow U \oplus V$ is equal to

$$\begin{pmatrix} f_{XZ} & f_{XW} \\ f_{YZ} & f_{YW} \end{pmatrix} \circ \begin{pmatrix} g_{ZU} & g_{ZV} \\ g_{WU} & g_{WV} \end{pmatrix} = \begin{pmatrix} f_{XZ} \circ g_{ZU} + f_{XW} \circ g_{WU} & f_{XZ} \circ g_{ZV} + f_{XW} \circ g_{WV} \\ f_{YZ} \circ g_{ZU} + f_{YW} \circ g_{WU} & f_{YZ} \circ g_{ZV} + f_{YW} \circ g_{WV} \end{pmatrix}$$

where $(f_{AB}: A \rightarrow B)_{A,B}$ and $(g_{BC}: B \rightarrow C)_{B,C}$ are the matrix decompositions of f and g .

Definition 2.9 In a pre-additive category, a (co)kernel of $f: X \rightarrow Y$ is, if it exists, a (co)equalizer of f and $0_{X,Y}$. An *abelian category* is a pre-additive category with finite limits and finite colimits such that every monomorphism is a kernel of some arrow, and every epimorphism is a cokernel of some arrow.

By the definition of abelian categories, every additive category is an abelian category. Categories **Ab**, **Mod** $_R$, **Vect** $_K$ and **fdVect** $_K$ are abelian categories. Let \mathcal{C} be an abelian category. For an arrow $f: X \rightarrow Y$ in \mathcal{C} , we write $Y \twoheadrightarrow \text{coker}(f)$ for the cokernel of $f: X \rightarrow Y$, and we write $\ker(f) \rightarrowtail X$ for the kernel of $f: X \rightarrow Y$. Because any epimorphism in \mathcal{C} is a cokernel, epimorphisms in \mathcal{C} are regular epimorphisms. It is known that any arrow $f: X \rightarrow Y$ in \mathcal{C} has a regular-epi/mono factorization

$$X \xrightarrow{e} \text{im}(f) \rightarrowtail Y$$

where m is the kernel of the cokernel $Y \twoheadrightarrow \text{coker}(f)$ and e is the cokernel of the kernel $\ker(f) \rightarrowtail X$. Furthermore, the pullback of an epimorphism along any arrow is an epimorphism. (See [3, Theorem 1.5.5 and Proposition 1.7.6].) Hence, any abelian category is a regular category. Because the definition of abelian category is self-dual, if \mathcal{C} is an abelian category, then \mathcal{C}^{op} is an abelian category. In particular, \mathcal{C}^{op} is a regular category.

An object X in a category \mathcal{C} is *projective* when for every epimorphism $f: Y \twoheadrightarrow Z$ and for every arrow $g: X \rightarrow Z$ in \mathcal{C} , there is $h: X \rightarrow Y$ such that $h \circ f = g$. An object $X \in \mathcal{C}$ is *injective* when X is projective in \mathcal{C}^{op} . The following definition is from [10].

Definition 2.10 An abelian category is *semisimple* when one of the following equivalent conditions is true:

- Every object is projective.
- Every object is injective.

The category \mathbf{Vect}_K is semisimple for any field K . More generally, when a ring R is semisimple, i.e., R is a direct sum of its irreducible left R -submodules, then the category \mathbf{Mod}_R of left R -modules is semisimple. For example, finite direct products $K_1 \times \cdots \times K_n$ of fields are semisimple. It follows from Maschke's theorem [11] that given a field K and a finite group G of order n such that the characteristic of K does not divide n , the category $\mathbf{Mod}_{K[G]}$ of left $K[G]$ -modules is semisimple where $K[G]$ is the group ring of G over K .

Remark 2.11 In the sequel, we always regard an additive category \mathcal{C} as a symmetric monoidal category $(\mathcal{C}, 0, \oplus)$. As usually, we omit canonical isomorphisms such as $0 \oplus X \cong X \cong X \oplus 0$ and $(X \oplus Y) \oplus Z \cong X \oplus (Y \oplus Z)$ when we can infer them from the context, and we simply write $X \oplus Y \oplus \cdots \oplus Z$ for $((X \oplus Y) \oplus \cdots) \oplus Z$.

Concrete description of \mathbf{fdVect}

As an illustration, we explain how the abelian category \mathbf{fdVect}_K looks like. We first note that we can regard each arrow $f: K^n \rightarrow K^m$ in \mathbf{fdVect}_K as an $n \times m$ -matrix. Below, we identify $n \times m$ -matrices with arrows from K^n to K^m , and we also identify the monoidal product $K^n \oplus K^m$ with K^{n+m} in the obvious way. Under this identification, the identity on K^n is the $n \times n$ identity matrix I_n , and the composition of arrows in \mathbf{fdVect}_K is the multiplication of matrices. The zero arrow $0_{K^n, K^m}: K^n \rightarrow K^m$ is the zero matrix, and the addition $f + g: K^n \rightarrow K^m$ is given by the pointwise manner. The projections $\pi_{K^n, K^m}: K^{n+m} \rightarrow K^n$ and $\pi'_{K^n, K^m}: K^{n+m} \rightarrow K^m$ are given by

$$\pi_{K^n, K^m} = \begin{pmatrix} I_n \\ 0_{K^m, K^n} \end{pmatrix}, \quad \pi'_{K^n, K^m} = \begin{pmatrix} 0_{K^n, K^m} \\ I_m \end{pmatrix},$$

and the injections $\iota_{K^n, K^m}: K^n \rightarrow K^{n+m}$ and $\iota'_{K^n, K^m}: K^m \rightarrow K^{n+m}$ are given by

$$\iota_{K^n, K^m} = \begin{pmatrix} I_n & 0_{K^n, K^m} \end{pmatrix}, \quad \iota'_{K^n, K^m} = \begin{pmatrix} 0_{K^m, K^n} & I_m \end{pmatrix}.$$

It is easy to check the four equations (2) are true for these projections and injections. For example, we have $\iota_{K^n, K^m} \circ \pi_{K^n, K^m} = \text{id}_{K^n}$ because

$$\begin{pmatrix} I_n & 0_{K^n, K^m} \end{pmatrix} \begin{pmatrix} I_n \\ 0_{K^m, K^n} \end{pmatrix} = I_n \circ I_n + 0_{K^n, K^m} \circ 0_{K^m, K^n} = I_n.$$

The matrix decomposition of an arrow $f: K^{n+m} \rightarrow K^{n'+m'}$ is the partitioning of f into the following 4 blocks:

$$\begin{pmatrix} x_{1,1} & \cdots & x_{1,n'} & z_{1,1} & \cdots & z_{1,m'} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ x_{n,1} & \cdots & x_{n,n'} & z_{n,1} & \cdots & z_{n,m'} \\ \hline y_{1,1} & \cdots & y_{1,n'} & w_{1,1} & \cdots & w_{1,m'} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ y_{m,1} & \cdots & y_{m,n'} & w_{m,1} & \cdots & w_{m,m'} \end{pmatrix}.$$

Given an arrow $f: K^n \rightarrow K^{n'}$ in \mathbf{fdVect}_K , let $u: K^n \rightarrow K^n$ and $v: K^{n'} \rightarrow K^{n'}$ be isomorphisms in \mathbf{fdVect}_K such that

$$f = u \circ \begin{pmatrix} I_l & 0_{K^l, K^{n'-l}} \\ 0_{K^{n-l}, K^l} & 0_{K^{n-l}, K^{n'-l}} \end{pmatrix} \circ v. \quad (3)$$

The natural number l is the rank of f . The regular-epi/mono factorization of f is of the following form:

$$K^n \xrightarrow{e} K^l \xrightarrow{m} K^{n'}$$

where e and m are given by

$$e = u \circ \begin{pmatrix} I_l \\ 0_{K^{n-l}, K^l} \end{pmatrix}, \quad m = \begin{pmatrix} I_l & 0_{K^l, K^{n'-l}} \end{pmatrix} \circ v.$$

Because u and v in (3) are not unique, the above decomposition of f is not unique. Still, the decomposition is unique up to isomorphism.

3 Partial traces on additive categories

Because an additive category with a total trace is equivalent to the trivial one-object additive category $\{0\}$ (see [8]), interesting partial traces on additive categories must be strictly partial. We give some examples of partial traces. The first two examples are from [5,13].

Example 3.1 We define a partial trace $\mathbf{ex}_{\mathcal{C}}$ on an additive category \mathcal{C} by


$$(\mathbf{ex}_{\mathcal{C}})_{X,Y}^Z(f) = \begin{cases} f_{XY} + f_{XZ} \circ (\mathrm{id}_Z - f_{ZZ})^{-1} \circ f_{ZY}, & \text{if } \mathrm{id}_Z - f_{ZZ} \text{ is invertible,} \\ \text{undefined,} & \text{otherwise} \end{cases}$$

where $(f_{AB}: A \rightarrow B)_{A,B}$ is the matrix decomposition of $f: X \oplus Z \rightarrow Y \oplus Z$. When

f_{ZZ} is nilpotent, $\text{id}_Z - f_{ZZ}$ is an isomorphism, and we have

$$(\mathbf{ex}_C)_{X,Y}^Z(f) = f_{XY} + \sum_{n \in \mathbb{N}} f_{XZ} \circ f_{ZZ}^n \circ f_{ZY}. \quad (4)$$

The right hand side is called the *execution formula* [6]. Informally, the result of the execution formula is the accepting language of the following automaton:



where ϵ is the ϵ -transition, and Y is the final state. We note that (4) does not make sense in general as the execution formula subsumes infinite summation.

Example 3.2 We give another partial trace \mathbf{ki}_C called the *kernel-image trace* on an additive category \mathcal{C} . For $f: X \oplus Z \rightarrow Y \oplus Z$, if there are $g: X \rightarrow Z$ and $h: Z \rightarrow Y$ such that

$$\begin{array}{ccccc} X & \xrightarrow{f_{XZ}} & Z & & \\ & \searrow g & \uparrow \text{id}_Z - f_{ZZ} & \searrow h & \\ & & Z & \xrightarrow{f_{ZY}} & Y \end{array},$$

we define $(\mathbf{ki}_C)_{X,Y}^Z(f)$ to be $f_{XY} + f_{XZ} \circ h$; otherwise, $(\mathbf{ki}_C)_{X,Y}^Z(f)$ is undefined. We note that $(\mathbf{ki}_C)_{X,Y}^Z(f)$ is independent of the choice of g and h . It is easy to see that when $\text{id}_Z - f_{ZZ}$ is invertible, $(\mathbf{ki}_C)_{X,Y}^Z(f)$ is defined and is equal to $(\mathbf{ex}_C)_{X,Y}^Z(f)$. Hence, we have $\mathbf{ex}_C \leq \mathbf{ki}_C$. For the definition of the partial order \leq between partial traces, see Example 2.5.

We say that a partial trace \mathbf{tr} on an additive category \mathcal{C} satisfies *strong naturality* when \mathbf{tr} satisfies

$$g \circ \mathbf{tr}_{X,Y}^Z(f) \circ h \simeq \mathbf{tr}_{U,V}^Z((g \oplus Z) \circ f \circ (h \oplus Z))$$

for all $f: X \oplus Z \rightarrow Y \oplus Z$, $g: U \rightarrow X$ and $h: Y \rightarrow V$ in \mathcal{C} . We give a characterization of \mathbf{ex}_C .

Theorem 3.3 Let \mathcal{C} be an additive category. We have:

- The partial trace \mathbf{ex}_C satisfies strong naturality.
- If a partial trace \mathbf{tr} on \mathcal{C} satisfies strong naturality, then $\mathbf{tr} \leq \mathbf{ex}_C$.

Hence, \mathbf{ex}_C is the largest partial trace that satisfies strong naturality.

Proof. It is easy to check the first claim. For the second claim, the point of the proof is that for a partial trace \mathbf{tr} on \mathcal{C} and for an arrow $f: X \oplus Z \rightarrow Y \oplus Z$ in \mathcal{C} ,

if

$$\mathbf{tr}_{Z,Z}^Z((Z \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus f_{ZZ}) \circ (Z \oplus \gamma_Z)): Z \rightarrow Z \quad (5)$$

is defined, then it is the inverse of $\mathrm{id}_Z - f_{ZZ}$ where $f_{ZZ}: Z \rightarrow Z$ is the (Z, Z) th entry of the matrix decomposition of f . We use strong naturality to show that if $\mathbf{tr}_{X,Y}^Z(f)$ is defined, then (5) is defined \square

We give an example of a partial trace that is smaller than $\mathbf{ex}_{\mathbf{Vect}_K}$. Let $G \subseteq K \setminus \{0\}$ be a subgroup. We define a partial trace \mathbf{tr} on \mathbf{Vect}_K by

$$\mathbf{tr}_{X,Y}^Z(f) = \begin{cases} f_{XY} + f_{XZ} \circ (\mathrm{id}_Z - f_{ZZ})^{-1} \circ f_{ZY}, & \text{if } \det(\mathrm{id}_Z - f_{ZZ}) \in G, \\ \text{undefined}, & \text{otherwise} \end{cases}$$

where $\{f_{AB}: A \rightarrow B\}_{A,B}$ is the matrix decomposition of f . We can check that \mathbf{tr} is a partial trace as in the proof of $\mathbf{ex}_{\mathbf{Vect}_K}$ being a partial trace given in [5, Section 2.2]. When G is not equal to $K \setminus \{0\}$, we have $\mathbf{tr} < \mathbf{ex}_{\mathbf{Vect}_K}$.

4 Partial traces on abelian categories

In this section, we study partial traces on abelian categories. When \mathcal{C} is an abelian category, \mathcal{C} and $\mathcal{C}^{\mathrm{op}}$ are regular categories. Therefore, as we observed in Example 2.7, there are two different constructions of partial traces on \mathcal{C} : one is $\mathbf{rel}_{\mathcal{C}}$ that is induced by the strong symmetric monoidal functor $H_{\mathcal{C}}: \mathcal{C} \rightarrow \mathbf{Rel}(\mathcal{C})$, and the other is $\mathbf{rel}_{\mathcal{C}^{\mathrm{op}}}$ that is induced by the strong symmetric monoidal functor $(H_{\mathcal{C}^{\mathrm{op}}})^{\mathrm{op}}: \mathcal{C} \rightarrow \mathbf{Rel}(\mathcal{C}^{\mathrm{op}})^{\mathrm{op}}$. The goal of this section is to clarify how these partial traces are related and to give their characterizations. Below, we fix an abelian category \mathcal{C} .

Proposition 4.1 $\mathbf{rel}_{\mathcal{C}} = \mathbf{rel}_{\mathcal{C}^{\mathrm{op}}}$.

Proof. (Sketch) We define a functor $\Theta: \mathbf{Rel}(\mathcal{C}) \rightarrow \mathbf{Rel}(\mathcal{C}^{\mathrm{op}})^{\mathrm{op}}$ by $\Theta X = X$ and

$$\Theta(\langle r, s \rangle: R \rightarrowtail X \oplus Y) = [q, p]: Y \oplus X \rightarrowtail S \quad \text{where} \quad \begin{array}{ccc} R & \xrightarrow{s} & Y \\ r \downarrow & \text{p.o} & \downarrow q \\ X & \xrightarrow{p} & S \end{array}.$$

It is easy to check that $H_{\mathcal{C}} \circ \Theta$ is equal to $(H_{\mathcal{C}^{\mathrm{op}}})^{\mathrm{op}}$. The functor Θ is a symmetric monoidal isomorphism and preserves total traces. Hence, by the definition of $\mathbf{rel}_{\mathcal{C}}$ and $\mathbf{rel}_{\mathcal{C}^{\mathrm{op}}}$, they are the same. \square

We next show that $\mathbf{rel}_{\mathcal{C}}$ is the largest partial trace on \mathcal{C} . We prepare two lemmas.

Lemma 4.2 Let $f: X \oplus Z \rightarrow Y \oplus Z$ be an arrow in \mathcal{C} , let $(f_{AB}: A \rightarrow B)_{A=X,Z; B=Y,Z}$ be the matrix decomposition of f , and let

$$Z \xrightarrow{e} \twoheadrightarrow W \xrightarrow{m} Z$$

be the regular-epi/mono factorization of $\text{id}_Z - f_{ZZ}$. If there are arrows $g: X \rightarrow W$ and $h: W \rightarrow Y$ such that the following triangles commute:

$$\begin{array}{ccc} X & \xrightarrow{f_{XZ}} & Z \\ & \searrow g & \uparrow m \\ & & W \\ & & \uparrow e \\ Z & \xrightarrow{f_{ZY}} & Y \end{array} \quad , \quad \begin{array}{ccc} & & h \\ & & \searrow \\ & & Y \end{array}$$

then $(\mathbf{rel}_C)_{X,Y}^Z(f)$ is defined and is equal to $f_{XY} + g \circ h$.

Proof. We define $p: P \rightarrow X$ to be the pullback of $e: Z \rightarrow W$ along $g: X \rightarrow W$:

$$\begin{array}{ccc} P & \xrightarrow{q} & Z \\ p \downarrow & \text{p.b} & \downarrow e \\ X & \xrightarrow{g} & W \end{array} .$$

By the definition of arrows p and q , the tuple $\langle p, q \rangle: P \rightarrow X \oplus Z$ is the kernel of $\pi_{X,Z} \circ g - \pi'_{X,Z} \circ e$. Since m is a monomorphism, $\langle p, q \rangle: P \rightarrow X \oplus Z$ coincides with the kernel of $(\pi_{X,Z} \circ g - \pi'_{X,Z} \circ e) \circ m$, which is equal to $f \circ \pi'_{Y,Z} - \pi'_{X,Z}$. Hence,

$$\begin{array}{ccc} P & \xrightarrow{q} & Z \\ \langle p, q \rangle \downarrow & & \downarrow \delta_Z \\ X \oplus Z & \xrightarrow{\langle \pi'_{X,Z}, f \circ \pi'_{Y,Z} \rangle} & Z \oplus Z \end{array}$$

is a pullback square. Because p is an epimorphism, the right hand side of

$$\langle p, q \rangle \circ \langle \pi_{X,Z}, f \circ \pi_{Y,Z} \rangle = p \circ \langle \text{id}_X, f_{XY} + g \circ h \rangle$$

is the regular-epi/mono factorization of the left hand side of this equation. Hence, $(\mathbf{rel}_C)_{X,Y}^Z(f)$ is defined and is equal to $f_{XY} + g \circ h$. \square

Lemma 4.3 Let \mathcal{C} be an additive category, and let \mathbf{tr} be a partial trace on \mathcal{C} . For all arrows $f: X \rightarrow Z$, $g: Z \rightarrow Z$ and $h: Z \rightarrow Y$ in \mathcal{C} , if

$$\mathbf{tr}_{X,Y}^Z((f \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus g) \circ (h \oplus \gamma_Z))$$

is defined, then for all $k: Z \rightarrow W$,

$$\mathbf{tr}_{X,W}^Z((f \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus g) \circ ((\text{id}_Z - g) \circ k) \oplus \gamma_Z))$$

is defined and is equal to $f \circ k$.

Lemma 4.4 Let $f: X \oplus Z \rightarrow Y \oplus Z$ be an arrow in \mathcal{C} , let $(f_{AB}: A \rightarrow B)_{A=X,Z; B=Y,Z}$ be the matrix decomposition of f , and let

$$Z \xrightarrow{e} \twoheadrightarrow W \xrightarrow{m} Z$$

be the regular-epi/mono factorization of $\text{id}_Z - f_{ZZ}$. Given a partial trace \mathbf{tr} on \mathcal{C} , if $\mathbf{tr}_{X,Y}^Z(f)$ is defined, then there are arrows $g: X \rightarrow W$ and $h: W \rightarrow Y$ such that the following triangles commute:

$$\begin{array}{ccc} X & \xrightarrow{f_{XZ}} & Z \\ & \searrow g & \uparrow m \\ & & W \\ & \uparrow e & \searrow h \\ Z & \xrightarrow{f_{ZY}} & Y \end{array} \quad ,$$

and $\mathbf{tr}_{X,Y}^Z(f)$ is equal to $f_{XY} + g \circ h$.

Proof. We first note that $m: W \rightarrowtail Z$ is the kernel of the cokernel of $\text{id}_Z - f_{ZZ}$, and $e: Z \twoheadrightarrow W$ is the cokernel of the kernel of $\text{id}_Z - f_{ZZ}$. Because $\mathbf{tr}_{X,Y}^Z(f)$ is defined, the right hand side of

$$\begin{aligned} -f_{XY} + \mathbf{tr}_{X,Y}^Z(f) &\simeq \gamma_X \circ ((-f_{XY}) \oplus \mathbf{tr}_{X,Y}^Z(f)) \circ \delta_Y \\ &\preceq \mathbf{tr}_{X,Y}^Z((f_{XZ} \oplus \gamma_Z) \circ (\sigma_{Z,Z} \oplus f_{ZZ}) \circ (f_{ZY} \oplus \delta_Z)) \end{aligned}$$

is defined. Therefore, by Lemma 4.3,

$$\mathbf{tr}_{X,Z}^Z((f_{XZ} \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus f_{ZZ}) \circ ((\text{id}_Z - f_{ZZ}) \oplus \gamma_Z))$$

is defined and is equal to f_{XZ} . Let $c: Z \rightarrow C$ be the cokernel of $\text{id}_Z - f_{ZZ}$. Then by naturality, we obtain

$$f_{XZ} \circ c \preceq \mathbf{tr}_{X,C}^Z((f_{XZ} \oplus f_{ZZ}) \circ \gamma_Z \circ \iota'_{C,Z}) \simeq \mathbf{tr}_{X,C}^Z(f \circ (0_{Y,C} \oplus Z)) \succeq \mathbf{tr}_{X,Y}^Z(f) \circ 0_{Y,C} \preceq 0_{X,C}.$$

Hence, there exists a unique $g: X \rightarrow W$ such that

$$f_{XZ} = X \xrightarrow{g} W \rightarrowtail \xrightarrow{m} Z.$$

In the same way, we can show that there exists a unique $h: W \rightarrow Y$ such that

$$f_{ZY} = Z \xrightarrow{e} \twoheadrightarrow W \xrightarrow{h} Y.$$

It remains to check that $\mathbf{tr}_{X,Y}^Z(f)$ is equal to $f_{XY} + g \circ h$. We define $p: P \twoheadrightarrow X$ to be the pullback of $e: Z \twoheadrightarrow W$ along $g: X \rightarrow W$:

$$\begin{array}{ccc} P & \xrightarrow{q} & Z \\ p \downarrow & \text{p.b} & \downarrow e \\ X & \xrightarrow{g} & W \end{array} \quad .$$

Because

$$\begin{aligned} p \circ (-f_{XY} + \mathbf{tr}_{X,Y}^Z(f)) &\simeq p \circ \gamma_X \circ ((-f_{XY}) \oplus \mathbf{tr}_{X,Y}^Z(f)) \circ \delta_Y \\ &\preceq \mathbf{tr}_{P,Y}^Z(((p \circ f_{XZ}) \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus f_{ZZ}) \circ (f_{ZY} \oplus \gamma_Z)) \\ &\simeq \mathbf{tr}_{P,Y}^Z(((q \circ (\mathrm{id}_Z - f_{ZZ})) \oplus \delta_Z) \circ (\sigma_{Z,Z} \oplus f_{ZZ}) \circ (f_{ZY} \oplus \gamma_Z)), \end{aligned}$$

it follows from Lemma 4.3 that $p \circ (-f_{XY} + \mathbf{tr}_{X,Y}^Z(f))$ is equal to $q \circ f_{ZY} = p \circ g \circ h$. Hence, we see that $\mathbf{tr}_{X,Y}^Z(f)$ is equal to $f_{XY} + g \circ h$. \square

The following theorems are straightforward consequences of Lemma 4.2 and Lemma 4.4. The first one means that $\mathbf{rel}_{\mathcal{C}}$ is the largest partial trace on \mathcal{C} , and the second one means that we can define $\mathbf{rel}_{\mathcal{C}}$ without referring to $\mathbf{Rel}(\mathcal{C})$.

Theorem 4.5 *If \mathbf{tr} is a partial trace on \mathcal{C} , then $\mathbf{tr} \leq \mathbf{rel}_{\mathcal{C}}$.*

Theorem 4.6 *Let $f: X \oplus Z \rightarrow Y \oplus Z$ be an arrow in \mathcal{C} , let $(f_{AB}: A \rightarrow B)_{A=X,Z; B=Y,Z}$ be the matrix decomposition of f , and let*

$$Z \xrightarrow{e} W \xrightarrow{m} Z$$

be the regular-epi/mono factorization of $\mathrm{id}_Z - f_{ZZ}$. Then we have

$$(\mathbf{rel}_{\mathcal{C}})_{X,Y}^Z(f) = \begin{cases} f_{XY} + h \circ k, & \text{if } f_{XZ} = h \circ m \text{ and } f_{ZY} = e \circ k \text{ for some} \\ & h: X \rightarrow W \text{ and } k: W \rightarrow Y, \\ \text{undefined,} & \text{otherwise.} \end{cases}$$

Corollary 4.7 *Any partial trace \mathbf{tr} on an abelian category \mathcal{C} is uniform in the following sense: for all arrows $f: X \oplus Z \rightarrow Y \oplus Z$ and $g: X \oplus W \rightarrow Y \oplus W$ such that $\mathbf{tr}_{X,Y}^Z(f)$ and $\mathbf{tr}_{X,Y}^W(g)$ are defined, if there is an arrow $h: Z \rightarrow W$ satisfying*

$$\begin{array}{ccc} X \oplus Z & \xrightarrow{f} & Y \oplus Z \\ \downarrow X \oplus h & & \downarrow Y \oplus h \\ X \oplus W & \xrightarrow{g} & Y \oplus W \end{array},$$

then $\mathbf{tr}_{X,Y}^Z(f) = \mathbf{tr}_{X,Y}^W(g)$.

Proof. We can easily check that $\mathbf{rel}_{\mathcal{C}}$ is uniform in the above sense. Uniformity of arbitrary traces \mathbf{tr} on \mathcal{C} follows from $\mathbf{tr} \leq \mathbf{rel}_{\mathcal{C}}$. \square

When \mathcal{C} is a semisimple abelian category, the kernel-image trace $\mathbf{ki}_{\mathcal{C}}$ is equal to the largest partial trace $\mathbf{rel}_{\mathcal{C}}$.

Corollary 4.8 *Let \mathcal{C} be a semisimple abelian category. If \mathbf{tr} is a partial trace on \mathcal{C} , then $\mathbf{tr} \leq \mathbf{ki}_{\mathcal{C}}$.*

Proof. We show that $\mathbf{ki}_{\mathcal{C}}$ coincides with $\mathbf{rel}_{\mathcal{C}}$. Let $f: X \oplus Z \rightarrow Y \oplus Z$ be an arrow in \mathcal{C} such that $(\mathbf{rel}_{\mathcal{C}})_{X,Y}^Z(f)$ is defined. Let $(f_{AB}: A \rightarrow B)_{A=X,Z; B=Y,Z}$ be

the matrix decomposition of f , and let

$$Z \xrightarrow{e} W \xrightarrow{m} Z$$

be the regular-epi/mono factorization of $\text{id}_Z - f_{ZZ}$. By Theorem 4.6, there are $g: X \rightarrow W$ and $h: W \rightarrow Y$ such that $f_{XZ} = g \circ m$ and $f_{ZY} = e \circ h$. Because every object is projective and injective, we obtain arrows $g': X \rightarrow Z$ and $h': Z \rightarrow Y$ such that every triangle in

$$\begin{array}{ccccc} X & \xrightarrow{f_{XZ}} & Z & & \\ & \searrow g & \uparrow m & \searrow h' & \\ & & W & & \\ & \searrow g' & \uparrow e & \searrow h & \\ & & Z & \xrightarrow{f_{ZY}} & Y \end{array}$$

commutes. Hence, $(\mathbf{ki}_C)_{X,Y}^Z(f)$ is defined and is equal to $f_{XY} + g' \circ f_{ZY} = (\mathbf{rel}_C)_{X,Y}^Z(f)$. \square

We note that Corollary 4.8 does not hold for \mathbf{Ab} , which is not a semisimple abelian category. To see this we show that $\mathbf{ki}_{\mathbf{Ab}} < \mathbf{rel}_{\mathbf{Ab}}$. Let $f: \mathbb{Z}_2 \oplus (\mathbb{Z} \oplus \mathbb{Z}_2) \rightarrow \mathbb{Z}_2 \oplus (\mathbb{Z} \oplus \mathbb{Z}_2)$ be an arrow in \mathbf{Ab} given by

$$f([x], y, [z]) = ([x], y, [x + y + z])$$

where we write \mathbb{Z}_2 for $\mathbb{Z}/2\mathbb{Z}$, and $[n] \in \mathbb{Z}_2$ is the equivalence class of an integer $n \in \mathbb{Z}$. We shall show that $(\mathbf{ki}_{\mathbf{Ab}})_{\mathbb{Z}_2, \mathbb{Z}_2}^{\mathbb{Z} \oplus \mathbb{Z}_2}(f)$ is not defined and that $(\mathbf{rel}_{\mathbf{Ab}})_{\mathbb{Z}_2, \mathbb{Z}_2}^{\mathbb{Z} \oplus \mathbb{Z}_2}(f)$ is defined and is equal to $\text{id}_{\mathbb{Z}_2}$.

- $(\mathbf{ki}_{\mathbf{Ab}})_{\mathbb{Z}_2, \mathbb{Z}_2}^{\mathbb{Z} \oplus \mathbb{Z}_2}(f)$ is not defined because there is no $g: \mathbb{Z}_2 \rightarrow \mathbb{Z} \oplus \mathbb{Z}_2$ such that

$$\iota_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2} \circ f \circ \pi'_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2} = g \circ (\text{id}_{\mathbb{Z} \oplus \mathbb{Z}_2} - \iota'_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2} \circ f \circ \pi'_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2}).$$

We note that the left hand side is equal to $\iota'_{\mathbb{Z}, \mathbb{Z}_2}$ and $\text{id}_{\mathbb{Z} \oplus \mathbb{Z}_2} - \iota'_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2} \circ f \circ \pi'_{\mathbb{Z}_2, \mathbb{Z} \oplus \mathbb{Z}_2}$ maps $(y, [z])$ to $(0, [y])$.

- $(\mathbf{rel}_{\mathbf{Ab}})_{\mathbb{Z}_2, \mathbb{Z}_2}^{\mathbb{Z} \oplus \mathbb{Z}_2}(f)$ is defined and is equal to $\text{id}_{\mathbb{Z}_2}$ because the triangles

$$\begin{array}{ccccc} \mathbb{Z}_2 & \xrightarrow{\iota'_{\mathbb{Z}, \mathbb{Z}_2}} & \mathbb{Z} \oplus \mathbb{Z}_2 & & \\ & \searrow \text{id}_{\mathbb{Z}_2} & \uparrow \iota'_{\mathbb{Z}, \mathbb{Z}_2} & \searrow 0_{\mathbb{Z}_2, \mathbb{Z}_2} & \\ & & \mathbb{Z}_2 & & \\ & & \uparrow e & & \\ & & \mathbb{Z} \oplus \mathbb{Z}_2 & \xrightarrow{0_{\mathbb{Z} \oplus \mathbb{Z}_2, \mathbb{Z}_2}} & \mathbb{Z}_2 \end{array}$$

commute where the vertical arrow $e \circ \iota'_{\mathbb{Z}, \mathbb{Z}_2}$ is the regular-epi/mono factorization of h . The epimorphism $e: \mathbb{Z} \oplus \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is given by $e(x, [y]) = [x]$.

References

- [1] Abramsky, S., E. Haghverdi and P. J. Scott, *Geometry of interaction and linear combinatory algebras*, Mathematical Structures in Computer Science **12** (2002), pp. 625–665.
- [2] Blute, R. and P. J. Scott, “Category Theory for Linear Logicians,” London Mathematical Society Lecture Note Series, Cambridge University Press, 2004 pp. 3–64.
- [3] Borceux, F., “Handbook of Categorical Algebra,” Encyclopedia of Mathematics and its Applications **2**, Cambridge University Press, 1994.
- [4] Girard, J.-Y., *Geometry of interaction 1: Interpretation of system F*, in: R. Ferro, C. Bonotto, S. Valentini and A. Zanardo, editors, *Logic Colloquium '88*, Studies in Logic and the Foundations of Mathematics **127**, Elsevier, 1989 pp. 221–260.
- [5] Haghverdi, E. and P. J. Scott, *Towards a typed geometry of interaction*, Mathematical Structures in Computer Science **20** (2010), pp. 473–521.
- [6] Haghverdi, E. and P. J. Scott, “Geometry of Interaction and the Dynamics of Proof Reduction: A Tutorial,” Springer-Verlag Berlin Heidelberg, 2011 pp. 357–417.
- [7] Hasegawa, M., “Models of Sharing Graphs: A Categorical Semantics of let and letrec,” Ph.D. thesis, University of Edinburgh (1997), ECS-LFCS-97-360.
- [8] Hoshino, N., *Partial traces on additive categories*, http://www.kurims.kyoto-u.ac.jp/~naophiko/mfps_proof.tex
- [9] Joyal, A., R. Street and D. Verity, *Traced monoidal categories*, Mathematical Proceedings of the Cambridge Philosophical Society **119** (1996), pp. 447–468.
- [10] Kashiwara, M. and P. Schapira, “Categories and Sheaves,” Grundlehren der mathematischen Wissenschaften **332**, Springer-Verlag Berlin Heidelberg, 2006.
- [11] Lang, S., “Algebra,” Graduate Texts in Mathematics **211**, Springer-Verlag New York, 2002.
- [12] Mac Lane, S., “Categories for Working Mathematician,” Springer-Verlag New York, 1978.
- [13] Malherbe, O., P. J. Scott and P. Selinger, *Partially traced categories*, Journal of Pure and Applied Algebra **216** (2012), pp. 2563–2585.
- [14] Selinger, P., *A survey of graphical languages for monoidal categories*, in: B. Coecke, editor, *New Structures for Physics*, Lecture Notes in Physics **813**, Springer-Verlag Berlin Heidelberg, 2011 pp. 289–355.