



Inside Every Model of Abstract Stone Duality Lies an Arithmetic Universe

Paul Taylor¹

Department of Computer Science, University of Manchester

Abstract

The first paper published on *Abstract Stone Duality* showed that the *overt discrete* objects (those admitting \exists and $=$ internally) form a *pretopos*, *i.e.* a category with finite limits, stable disjoint co-products and stable effective quotients of equivalence relations. Using an \mathbb{N} -indexed least fixed point axiom, here we show that this full subcategory is an *arithmetic universe*, having a *free semilattice* (“collection of *Kuratowski-finite* subsets”) and a *free monoid* (“collection of *lists*”) on any overt discrete object. Each finite subset is represented by its pair (\Box, \Diamond) of *modal operators*, although a tight correspondence with these depends on a stronger Scott-continuity axiom. Topologically, such subsets are both *compact and open* and also involve *proper open maps*. In applications of ASD this can eliminate lists in favour of a *continuation-passing interpretation*.

Key words: Abstract Stone Duality, pretopos, arithmetic universe, free semilattice, Kuratowski-finite, lists, modal logic, powerdomains, proper open maps.

Contents		
1.	Introduction	247
2.	Proofs and natural numbers	252
3.	Finite subsets as modalities	259
4.	Fixed point properties	263
5.	Stages in the construction	267
6.	Admissible implies modal	271
7.	K as a functor	273
8.	Lists, heads and tails	278
9.	Recursion over lists	282
10.	The free semilattice	287
11.	Overt compact subspaces	290

1 Introduction

In Abstract Stone Duality the topology on a space X is treated as an exponential Σ^X with a λ -calculus rather than as a lattice with arbitrary joins. This has given accounts of the category of locally compact spaces, both over an

¹ Correspondence by email only, please, to [<pt@cs.man.ac.uk>](mailto:pt@cs.man.ac.uk)

elementary topos [H], and for a logic in which the maps $\mathbb{N} \rightarrow \mathbb{N}$ are precisely the provably total general recursive functions [G].

Remark 1.1 It is an important feature of ASD that its spaces do not have “underlying sets” of points — or even of open subspaces, as in locale theory. ASD is a *direct* axiomatisation of the category \mathcal{S} of “spaces”, amongst which the “discrete” ones serve in the role of “sets”. However, as we take the word discrete to mean that there is an internal notion of equality, $(=_X) : X \times X \rightarrow \Sigma$ (*i.e.* the diagonal $X \subset X \times X$ is open), we actually say ***overt discrete***, meaning that there is also an “existential quantifier”, $\exists_X : \Sigma^X \rightarrow \Sigma$.

Remark 1.2 Having postulated a notion of the category of “sets” in this roundabout fashion, *i.e.* as the full subcategory \mathcal{E} of overt discrete types in a certain λ -calculus, we are faced with the challenge of showing that it has enough of the usual features of set theory or categorical logic to warrant the name, as none of these went in as ingredients. In fact, \mathcal{E} is a topos if we assume the existence of “underlying sets”, *i.e.* a right adjoint to the inclusion $\mathcal{E} \hookrightarrow \mathcal{S}$ [H], so it is important *not* to make such an assumption if we want to develop a computational axiomatisation of topology.

Giraud’s theorem, which characterised Grothendieck toposes in terms of the limits and (infinitary) colimits that they admit, suggested the first categorical approximation to the “finitary” aspects of the category of sets: a ***pre-topos*** is a category with finite limits, stable disjoint coproducts and stable effective quotients of equivalence relations. Then André Joyal introduced ***arithmetic universes*** to prove Gödel’s incompleteness theorem in a categorical style: they have just enough structure to form the *free internal* thing of the same kind. Specifically, an arithmetic universe is a pretopos with free internal monoids ($\text{List}X$), from which the free internal arithmetic universe can be obtained by means of generators and relations.

Actually, these structures are just what is taught in a first year “discrete math” course intended for computer scientists, *i.e.* one in which only the collection of *listable* subsets is considered, instead of a full set-theoretic powerset. We shall argue in future work (on the construction of \mathcal{S} from \mathcal{E}) that the other substitutes for the powerset, *i.e.* the collections of *recursively enumerable* and of *decidable* subsets of X , are the non-discrete spaces Σ^X and $\mathbf{2}^X$. If Σ^X had an underlying set, this would be the usual powerset of X .

Remark 1.3 Unfortunately, despite their 30-year history, knowledge of arithmetic universes circulates literally by word of mouth. The only only refereed (or even obtainable) papers on them are those by Maria Emilia Maietti, who provided a Martin-Löf-style type theory for them [6]. She claims that her

notion of arithmetic universe [7] is stronger than that used here. Indeed, some disagreement over detail is known by those who have worked with arithmetic universes — but they hardly help matters by failing to write any papers. However, the best way to settle doubts over correct definitions in mathematics is to prove their equivalence with other structures that come from other intuitions. The construction in this paper of an arithmetic universe from a model of ASD and the proposed converse will, I believe, not only serve this purpose, but also provide a more expressive calculus than Maietti's, allowing domain-theoretic methods to be used to define structures within arithmetic universes more fluently.

Remark 1.4 In ASD the full subcategory of overt discrete objects is already known to be a pretopos [C], so the present work is concerned with the free monoid and the free semilattice (KX).

Note, however, that, as an existential question, this is redundant in the two cases on which we usually focus: in the classical models (the source of our intuitions) \mathcal{E} is a topos, whilst in the free one (the target of our computations) every object of \mathcal{E} is a subquotient of \mathbb{N} by an open partial equivalence relation; in both cases lists can be encoded in well known ways.

This paper is also arguably unnecessary from a topological point of view, in that the existence of the free monoid could reasonably be taken as another axiom, so our conclusion is merely that this axiom is redundant.

Nevertheless, ASD sets itself apart from other approaches to topology by having a *freestanding* (technically) elementary axiomatisation that does not rely on a pre-existing category of sets or spaces. The intention is to use this as a route to computation, so our *representation* of KX is of interest even when we already know abstractly that it exists.

Remark 1.5 The two semilattice structures that we use most often are \wedge and \vee in powers of Σ , and it turns out that these are jointly faithful, indeed that KX is a subspace of $\Omega \equiv \Sigma^{\Sigma^X} \times \Sigma^{\Sigma^X}$, where each finite subset ℓ is represented by the two *modal operators*,

$$[\ell] \equiv \lambda\phi. \forall x \in \ell. \phi x \quad \text{and} \quad \langle \ell \rangle \equiv \lambda\phi. \exists x \in \ell. \phi x.$$

It is well known that modal logic is related to the three *powerdomains*, in one of which the inclusion order agrees with the intrinsic one, in another they are contravariant, whilst in the third inclusion involves both the intrinsic order and its opposite [4,9,10]. However, we only consider the convex powerdomain of an (overt) discrete space. The reason for this lack of ambition is that this paper forms part of the “bootstrapping” of the theory

of ASD: much more can be done using the whole thing, in particular [G]; the analogous structure in an Hausdorff space such as \mathbb{R} is considered in [I]. Also, we represent *elements* of the powerdomain by modal operators, whereas the works cited use them to generate its *topology*, relying on the prior existence of the free semilattice.

Remark 1.6 Our translation of finite subsets into higher order λ -terms may be of computational value, like other examples of the *continuation passing style*. The theory of locally compact spaces is developed in [G] using bases of open and compact subspaces, and that paper concludes with a sketch of how continuous functions (for example $\mathbb{R} \rightarrow \mathbb{R}$) might be manipulated computationally using relations. Unfortunately, at least as the theory is expressed there, such bases have to be indexed by (finitary) *lattices*, and heavy use is made of the free distributive lattice generated by an overt discrete object. This may be constructed *via* the free semilattice, whose elements might in turn be represented as *lists*, but possibly at quite a heavy computational cost.

On the other hand, as functional programmers know very well, there are plenty of situations in which (nested) lists provide excellent data structures. For this, it would be absurd to take a diversion *via* logic, especially by means of the construction that we use in Sections 8–9. What this paper provides, therefore, is a *choice* between lists (as traditionally implemented) and a mathematically isomorphic structure that encodes finite subsets using modal logic, λ -calculus and continuation-passing. Empirical study will be needed to make this choice in particular applications.

Remark 1.7 Returning to type theory, the disagreement with Maietti regarding the definition of an arithmetic universe appears to concern *parametric* list types, which are not covered in this paper. Certainly these are needed, but there are ways of obtaining them without first developing dependent spaces in general.

A family of overt discrete objects indexed by another such object is given, as in a topos, by *any* map $\delta : X \rightarrow N$, where $X[n] \equiv \delta^{-1}(n)$. (We call δ the “display” map [8, Chapter VIII].) Then $\text{List}X[n]$ is given by the display $N + P \rightarrow N$, where

$$\begin{array}{ccc}
 P & \xrightarrow{\quad} & N \\
 \downarrow \lrcorner & & \downarrow \{-\} \\
 \text{List}X & \xrightarrow{\text{List}\delta} \text{List}N \longrightarrow & KN
 \end{array}$$

provides the non-empty lists and the extra term N the empty ones.

Remark 1.8 Another connection between our construction and domain theory is that it uses the methods of that subject. Specifically, the ASD calculus provides for the introduction of subtypes of a given ambient type Ω , defined by an endomorphism \mathcal{E} of Σ^Ω . In this paper we obtain that endomorphism as the least fixed point of an operator $\$$ of yet higher type.

To do this, a fixed point axiom must be added to what [C] used to show that \mathcal{E} is a pretopos. To construct KX and $ListX$ it is enough to assume

Axiom 1.9 The “linear countable” *fixed point axiom* is

$$\frac{\Gamma, n : \mathbb{N} \vdash \phi n : \Sigma^U \quad \Gamma, n : \mathbb{N} \vdash \phi n \leq \phi(n+1) : \Sigma^U}{\Gamma, F : \Sigma^U \rightarrow \Sigma^V \vdash F(\exists n. \phi n) = \exists n. F(\phi n) : \Sigma^V}$$

However, in order to obtain all of the topological results that we expect [G], and in particular to show that every pair (\Box, \Diamond) of operators satisfying the modal laws actually corresponds to a listable subset (Section 11), the following stronger² assumption is needed. Despite the conceptual simplification that it would bring throughout this paper, we are unable even to state it before proving our main result, as it involves either $ListX$ or KX .

Axiom 1.10 The *Scott-continuity axiom* is

$$\frac{\begin{array}{l} \Gamma, \ell : KX \vdash \phi^\ell : \Sigma^U \quad \Gamma, \ell_1, \ell_2 : KX \vdash \phi^{\ell_1} \vee \phi^{\ell_2} \leq \phi^{\ell_1 + \ell_2} \\ \Gamma, \ell : KX \vdash \alpha_\ell : \Sigma \quad \Gamma \vdash \alpha_{nil} = \top \quad \Gamma, \ell_1, \ell_2 : KX \vdash \alpha_{\ell_1 + \ell_2} = \alpha_{\ell_1} \wedge \alpha_{\ell_2} \end{array}}{\Gamma, F : \Sigma^U \rightarrow \Sigma^V \vdash F(\exists \ell. \alpha_\ell \wedge \phi^\ell) = \exists \ell. \alpha_\ell \wedge F\phi^\ell.}$$

A family (α_ℓ, ϕ^ℓ) satisfying the five premises is called a *directed diagram*. In the free model every overt discrete object such as KX is a subquotient of \mathbb{N} ; in this case general Scott-continuity may be derived from the fixed point axiom above. In other models, such as the classical ones, a Scott-continuity axiom is needed for each overt discrete object X .

Remark 1.11 This brings us back to the study of *general* models of ASD (for which the existential question about KX and $ListX$ is not trivial). Besides

² The fixed point axiom says that F preserves *countable* directed joins, but in the classical models *uncountable* directed joins are also needed. Recall, for example, that in universal algebra the free functor for a theory of infinite arity κ preserves κ -filtered colimits of diagrams of any size λ . The two versions of our axiom correspond very roughly to the roles of these two cardinal parameters in the classical case.

the free and classical ones, we may obtain other models by strengthening the axioms with more ways of forming types, terms or equations. For example, the “underlying set” is a new *type* [H]. An oracle for termination would say that the (already definable) space \bar{T} of codes for non-terminating programs is overt, providing a new *term* $\exists_{\bar{T}} : \Sigma^{\bar{T}} \rightarrow \Sigma$.

The very meaning of the assertions that we made in the opening paragraph depends on what we can prove about *equality* of terms. A morphism of the category is a class of terms that are provably equivalent according to a certain logic, whilst a partial morphism $f : \mathbb{N} \multimap \mathbb{N}$ is total by definition iff there is a proof that Σ^f satisfies a certain equation [D]. Since the definitions of *prime* and *nucleus* [A,B] depend on equations, if the logic proves more of them then it also defines more terms and (sub)types.

What counts as a legitimate proof (in particular of the equality of two terms) is an issue that we have to consider more carefully in this paper than has been done hitherto in the ASD literature, because this is the first *serious* use that we have made of recursion and induction over \mathbb{N} . That is the subject of the next section.

Section 3 gives the intuition and notation for the construction of KX and its modal logic, which are developed in Sections 4–6. Section 7 considers the properties of K as a functor. The universal property of KX as the free semilattice is, however, only proved in Section 10, being derived from that for $ListX$, which we construct in Sections 8–9. Section 11 reconsiders the sense in which KX is a finite powerset.

2 Proofs and natural numbers in ASD

Before we begin the construction of the free monoid and semilattice, we have to consider the form of proofs in the λ -calculus for ASD more carefully than has been done in previous papers. Nevertheless, our purpose is a domain-theoretic construction, not a proof-theoretic analysis of ASD, so we shall not give the complete set of rules. See in particular [B, §8] for a summary of the λ -calculus for $\{X \mid E\}$ that handles the monadic property.

The main issue to be considered is induction over \mathbb{N} , the point being that the definition of the natural numbers object is not adequate as it is usually given. It is well known that an object Γ of *parameters* has to be added explicitly to the definition when we work in a category that is not *cartesian closed*. Similarly, *equational hypotheses* have to be considered explicitly when the category does not have all *equalisers*.

Axiom 2.1 The λ -calculus for ASD consists of types, terms and equations.

Its judgements assert that

- types such as $\mathbf{0}$, $\mathbf{1}$, \mathbb{N} , Σ , $X \times Y$, Σ^X and $\{X \mid E\}$ are well formed,
- terms are well formed and of particular types, or
- equations hold between terms.

Such judgements about terms and equations are made in certain *contexts*, *i.e.* on the assumption that their free variables have certain types.

However, even though the definition of the subtype $\{X \mid E\}$ involves the term E , we insist (as the theory is currently formulated) that it be formed in the empty (global) context, *i.e.* without free variables. Even the pure syntax of dependent type theories is very complicated, and becomes more so in a semantic situation, where we have to choose a class of display maps [8, Chapter VIII]. Of course, we must also perform the semantic calculations, which form the main task of this paper. Families of overt and compact objects are encoded as open and proper maps respectively [C, §7].

Judgements are therefore of the four forms

$$\vdash X \text{ type}, \quad \Gamma \vdash a : X, \quad \Gamma \vdash a = b : X \quad \text{and} \quad \Gamma \vdash \alpha \leq \beta : X.$$

The last of these arises from the lattice structure on types of the form Σ^U , where $(\alpha \leq \beta)$ means $\alpha = (\alpha \wedge \beta)$ or $\beta = (\alpha \vee \beta)$. The order can be extended to other objects, but we shall not need that.

In this paper we find that *equations* (and inequalities) *are also needed as assumptions*. In other words, the context Γ may include equations and inequalities between terms, as well as a list of typed variables. Any provable judgement attests to the validity of a certain *fragment* of reasoning from its hypotheses to its conclusions, so, as we want to form bigger arguments by concatenating such fragments (*i.e.* by means of a cut rule), all forms of assertion that are allowed as conclusions should also be allowed as hypotheses. In particular this is needed for induction over \mathbb{N} , which internalises the process of concatenation. (Adding such hypotheses to Martin-Löf type theory leads to undecidability [2, §3.2], but the ASD calculus is Turing-complete by design anyway.)

Axiom 2.2 Terms of type Σ may be seen as predicates in coherent logic:

- (a) \top and \perp ;
- (b) $\alpha \wedge \beta$ and $\alpha \vee \beta$, where $\alpha, \beta : \Sigma$;
- (c) ϕa , where $\phi : \Sigma^X$ and $a : X$;
- (d) $(a =_X b)$, where $a, b : X$, and X is discrete (such as \mathbb{N} but not \mathbb{R} or $\mathbf{2}^{\mathbb{N}}$);
- (e) $(a \neq_X b)$, where X is a Hausdorff type (such as \mathbb{N} , \mathbb{R} or $\mathbf{2}^{\mathbb{N}}$);

- (f) $\exists_X \phi \equiv \exists x. \phi x$, where $\phi : \Sigma^X$ and X is an overt type (\mathbb{N} , \mathbb{R} , $2^{\mathbb{N}}$);
 (g) $\forall_X \phi \equiv \forall x. \phi x$, where $\phi : \Sigma^X$ and X is a compact type (such as $[0, 1] \subset \mathbb{R}$ or $2^{\mathbb{N}}$ but not \mathbb{N}).

We shall find that a type X has all four properties (d–g) iff it is finite, whilst it is *Kuratowski*-finite iff it has properties (d,f,g), that notion being the main subject of this paper. Terms of type $\phi : \Sigma^X$ are formed with a λ -calculus. As usual, the introduction of λx , $\exists x$ or $\forall x$ discharges the variable from the context, which must therefore contain no equational assumption in which x is free.

Remark 2.3 Notice that this logic does not include implication. Indeed, every term is monotone considered as a function of its free variables of type Σ^U . But a strictly limited form of implication is allowed in that assumptions and conclusions of judgements may be of the form $\alpha \leq \beta : \Sigma^U$, so judgements may be of the form

$$\dots, \alpha \leq \beta : \Sigma^U, \dots \vdash \gamma \leq \delta : \Sigma^V,$$

which means $(\forall u. \alpha u \Rightarrow \beta u) \Rightarrow (\forall v. \gamma v \Rightarrow \delta v)$ in traditional notation.

When we need to make an assertion α that is a term of type Σ , we shall often follow mathematical idiom by saying simply “ α ” alone; by this we mean the judgement $\Gamma \vdash \alpha = \top : \Sigma$, where the context Γ has been established implicitly in the argument.

Of course, any equation or inequality that is an assumption may be used directly as the conclusion of an “identity” or “axiom” judgement, or may be discharged by a “cut” rule. They are also discharged by logical rules that we now give.

Axiom 2.4 Even though we cannot define a term $\alpha \Rightarrow \beta$ of type Σ , the usual rule for it does introduce an inequality, in both an intuitionistic way and an apparently classical one:

$$\frac{\Gamma, \alpha = \top : \Sigma \vdash \beta = \top : \Sigma}{\Gamma \vdash \alpha \leq \beta : \Sigma} \qquad \frac{\Gamma, \alpha = \perp : \Sigma \vdash \beta = \perp : \Sigma}{\Gamma \vdash \alpha \geq \beta : \Sigma}$$

The intuitionistic rule is equivalent to the *Euclidean principle*,

$$F : \Sigma^{\Sigma}, \alpha : \Sigma \vdash \alpha \wedge F\alpha = \alpha \wedge F\top,$$

and the other rule to its lattice dual. Writing α and $\neg\beta$ for $\alpha = \top$ and $\beta = \perp$, together they yield rules similar to those for negation in Gerhard Gentzen’s classical sequent calculus.

They and monotonicity are together equivalent to the **Phoa principle**,

$$F : \Sigma^\Sigma, \alpha : \Sigma \vdash F\alpha = F\perp \vee \alpha \wedge F\top.$$

Note that these things are valid in intuitionistic locale theory [C, Section 5]. The Euclidean principle is equivalent (in the context of the monadicity assumption) to saying that $\top : \Sigma$ classifies a certain class of monos in \mathcal{S} , which we call **open**, whilst its dual says that $\perp : \Sigma$ classifies **closed inclusions**.

An example of the first principle that's obvious in the abstract but may look strange when it's used in practice is the rule (where X and Y are discrete)

$$\frac{\Gamma, a = b : X \vdash c = d : Y}{\Gamma \vdash (a =_X b) \leq (c =_Y d) : \Sigma} \quad \square$$

Turning to recursion and induction, we have to reconsider the definition of \mathbb{N} in a category without equalisers, such as that of locally compact spaces. See also [1] and [5, §II.4].

Axiom 2.5 The **recursion scheme** introduces *terms* of any type X , dependent on $n : \mathbb{N}$:

$$\frac{\Gamma \vdash z : X \quad \Gamma, n : \mathbb{N}, x : X \vdash s(n, x) : X}{\Gamma, n : \mathbb{N} \vdash \text{rec}(n, z, s) : X}$$

Its meaning is given by the β -rules

$$\begin{aligned} \Gamma &\vdash \text{rec}(0, z, s) = z : X \\ \Gamma, n : \mathbb{N} &\vdash \text{rec}(n+1, z, s) = s(n, \text{rec}(n, z, s)). \end{aligned}$$

This is called **recursion at type** X , the point being that as the class of types at which the recursion scheme is asserted grows, so considerably does the power of the logic.

The corresponding diagrammatic property is

$$\begin{array}{ccccc} \Gamma & \xrightarrow{0} & \Gamma \times \mathbb{N} & \xleftarrow{\text{id} \times \text{succ}} & \Gamma \times \mathbb{N} \\ & \searrow z & \downarrow \text{rec} & & \downarrow (\text{id}, \text{rec}) \\ & & X & \xleftarrow{s} & \Gamma \times \mathbb{N} \times X. \end{array}$$

If the category is cartesian closed then this diagram may be rewritten with $\mathbb{N} \times X^\Gamma$ in place of X , the object Γ itself being removed from the top line.

Symbolically, this means that the parameters can be embedded in both the data and the new term by means of λ -abstraction. If, as in ASD, the category (has finite products but) is not cartesian closed then the object Γ of parameters is needed.

Axiom 2.6 The *equational induction scheme* for $\Gamma, n : \mathbb{N} \vdash a_n, b_n : X$:

$$\frac{\Gamma \vdash a_0 = b_0 : X \quad \Gamma, n : \mathbb{N}, a_n = b_n : X \vdash a_{n+1} = b_{n+1} : X}{\Gamma, n : \mathbb{N} \vdash a_n = b_n : X}$$

Notice that this says nothing about where a_n and b_n themselves *came* from. If they had been defined by the same base and step data on X , an η -rule or uniqueness hypothesis in the recursion scheme above would have made them equal. However, the scheme is more general than that: the term a_n need not arise *directly* from recursion, but may perhaps be of the form $f(\text{rec}(n, z, s))$ where f is some function, whilst b_n is some unrelated $f'(\text{rec}(n, z', s'))$.

Remark 2.7 Just as the generalisation with parameters was redundant in a cartesian closed category, so the equational induction scheme can be derived from the usual universal property of \mathbb{N} when the category has equalisers:

$$E \equiv \{(\gamma, n) \mid a_n(\gamma) = b_n(\gamma) : X\} \longrightarrow \Gamma \times \mathbb{N} \begin{array}{c} \xrightarrow{a_n} \\ \xrightarrow{b_n} \end{array} X.$$

The base case says that $(\text{id}, 0) : \Gamma \rightarrow \Gamma \times \mathbb{N}$ factors through E , whilst the induction step says that $\text{id} \times \text{succ}$ restricts to a map $E \rightarrow E$. Then Axiom 2.5 provides

$$\text{rec}_E : \Gamma \times \mathbb{N} \rightarrow E.$$

This is inverse to the inclusion, one equation being given by uniqueness of

$$\text{rec}_{\mathbb{N}} : \Gamma \times \mathbb{N} \rightarrow \Gamma \times \mathbb{N}$$

and the other by the fact that $E \hookrightarrow \Gamma \times \mathbb{N}$ is mono. □

The equational induction scheme (and its equivalent version for inequalities $\alpha_n \leq \beta_n$) will be used in Propositions 4.4, 4.12, 5.9 and 7.10, and we shall deduce its analogue for lists and finite subsets in Propositions 9.7 and 10.11. It ought to have been stated as [A, Remark 2.5], since it was actually used in [A, Lemmas 8.9 and 9.6] and [B, Lemma 8.14].

Since the proof given for [A, 9.6] was sketchy and contained other errors, we give the correct version here by way of an example of equational induction.

It turns recursion at type \mathbb{N} into recursion at type $\Sigma^{\mathbb{N}}$, as part of the process of bringing descriptions (**the** m .) to the outside of a term.

Lemma 2.8 $\Gamma \vdash rn \equiv \text{rec}(n, z, \lambda n' u. s(n', u)) = \text{the } m. \rho(n, m)$, where

$$\begin{aligned}\zeta &\equiv (\lambda m. m = z), \\ \sigma(n, \phi) &\equiv (\lambda m. \exists m'. \phi m' \wedge m = s(n, m')) \\ \rho(n, m) &\equiv \text{rec}(n, \zeta, \sigma)m.\end{aligned}$$

Proof The base case (in context Γ) is

$$\begin{aligned}\lambda m. \rho(0, m) &\equiv \lambda m. \text{rec}(0, \zeta, \sigma)m = \lambda m. \zeta m && \beta\text{-rule} \\ &\equiv \lambda m. (m = z) && \text{def } \zeta \\ &= \lambda m. (m = r0) && \beta\text{-rule}\end{aligned}$$

The induction step, with the hypothesis $\lambda m'. \rho(n, m') = \lambda m'. (m = rn)$, is

$$\begin{aligned}\lambda m. \rho(n+1, m) &\equiv \lambda m. \text{rec}(n+1, \zeta, \sigma)m \\ &= \lambda m. \sigma(n, \text{rec}(n, \zeta, \sigma))m && \beta\text{-rule} \\ &= \lambda m. \exists m'. \text{rec}(n, \zeta, \sigma)m' \wedge (m = s(n, m')) && \text{def } \sigma \\ &\equiv \lambda m. \exists m'. \rho(n, m') \wedge (m = s(n, m')) \\ &= \lambda m. \exists m'. (m' = rn) \wedge (m = s(n, m')) && \text{hypothesis} \\ &= \lambda m. (m = s(n, rn)) && \text{equality rules} \\ &= \lambda m. (m = r(n+1)) && \beta\text{-rule}\end{aligned}$$

So $\Gamma, n : \mathbb{N} \vdash \lambda m. \rho(n, m) = \lambda m. (m = rn) : \Sigma$ by Axiom 2.6. \square

Remark 2.9 Recall from [B, Section 8] that the rules for the introduction of **focus** and **admit** terms have equational premises. This now means that there are such terms that are only defined in certain contexts that contain particular equational assumptions.

Recall, however, from *op. cit.* that, for any term $\Gamma \vdash \alpha : \Sigma$, there is another term $\Gamma \vdash \bar{\alpha} : \Sigma$ not involving **focus** or **admit**, with the property that $\Gamma \vdash \alpha = \bar{\alpha} : \Sigma$. (In fact, $\bar{\alpha}$ is obtained, essentially, by *erasing* **admit**.) Now, as $\bar{\alpha}$ does not contain **focus** or **admit**, no equational hypotheses are ever used in its formation, and so it is defined in the context $\bar{\Gamma}$ without them. \square

Corollary 2.10 Σ is injective in the category of contexts (possibly involving equations) and substitutions. \square

Remark 2.11 How might such equational hypotheses be interpreted in a category \mathcal{S} that does not necessarily have all equalisers? The obvious way is that, for $f_1, f_2 : X \rightarrow A$ and $g_1, g_2 : X \rightarrow B$ in \mathcal{S} , the statement

$$x : X, f_1x = f_2 : A \vdash g_1x = g_2x : B$$

means that

$$\text{for every } a : \Gamma \rightarrow X \text{ in } \mathcal{S}, \quad \text{if } (f_1 \cdot h) = (f_2 \cdot h) \quad \text{then } (g_1 \cdot h) = (g_2 \cdot h).$$

The restricted λ -calculus was formally extended with (abstract) sobriety [A] and monadicity [B]. This interpretation of equational hypotheses could be used to make another similar formal extension.

However, the result of such an extension would be an account of *spatial* locales (equivalently, sober spaces — in the traditional sense rather than that of [A]), not general ones. This is because the generality of the test object Γ is spurious: locally compact locales have enough points (classically, at least), and so this formula only uses the global points of Γ to test the equations. Equalisers (and, in fact, products) of locales and sober spaces are not the same [3, §II 2.13].

Remark 2.12 The category whose objects are contexts with equations has finite products, but it no longer has the exponentials $\Sigma^{(-)}$ that originally motivated ASD — these are only defined on the subcategory of contexts without equations. The problem is inescapable since, as we have just noted, such hypotheses have already been used, albeit inadvertently. This means that the theory really captures, not the category of locally compact spaces on its own, but that category embedded in either the category of locales or or spatial locales.

Nevertheless, the way forward is not to rewrite what has been done in this hybrid fashion, but to study the (*substantially*) larger structure that includes both equalisers and exponentials. Preliminary investigations of this structure, relating not only to the whole of the category of locales but also to cartesian closed extensions, may be found in [H]. However, this structure not only contains spaces of a much more general kind, but also captures a much stronger logic of (implications amongst) equations. This means that, in order to nail down the precise equivalence with arithmetic universes (Remark 1.3), the proposed converse construction will probably be of a category analogous to spatial locales.

But for the purposes of the present paper, the equational hypotheses are simply a device for managing proofs.

3 Finite subsets as modal operators

Throughout, let X be an overt discrete space. Its typical values will be called x, y , etc. and its predicates ϕ, ψ , etc. We shall construct an object $\mathbf{K}X$ and then show that it is the free semilattice on X . Although $\mathbf{K}X$ is intended to be the space of Kuratowski-finite *subsets* of X , we shall use *list* notation for it, writing $\ell : \mathbf{K}X$ for a typical value. So $\text{nil}, \{\!\{x\}\!\} : \mathbf{K}X$ denote the empty and singleton subsets, $+$ is union and $x :: \ell = \{\!\{x\}\!\} + \ell$. Note that we also use $\{x\}$ for $\lambda y. (x =_X y) : \Sigma^X$. We shall actually need to switch back and forth between $\mathbf{K}X$ and $\text{List}X$, and between the “mathematical” notion of monoid based on the $+$ operation and the “computer science” notion based on $::$.

Remark 3.1 $\mathbf{K}X$ will be constructed as a Σ -split subspace of $\Omega \equiv \Sigma^{\Sigma^X} \times \Sigma^{\Sigma^X}$.

The typical values of Ω will be called $L = (N, P)$, where the letters stand for *necessity* and *possibility*, as the central idea is to represent a subset $\ell : \mathbf{K}X$ by means of its *modal operators* $[\ell], \langle \ell \rangle : \Sigma^{\Sigma^X}$, which also vary *negatively* and *positively* with respect to the inclusion of lists. Along with $P : \Sigma^{\Sigma^X}$ we shall also make frequent use of $\pi \equiv \lambda x. P\{x\} : \Sigma^X$.

Remark 3.2 If we already had $\mathbf{K}X$ and $\text{List}X$ at our disposal, we would define

$$[\ell] \equiv \lambda \phi. \forall x \in \ell. \phi x \quad \text{and} \quad \langle \ell \rangle \equiv \lambda \phi. \exists x \in \ell. \phi x,$$

and the *membership predicate* $(x \in \ell) \equiv \pi_\ell x = P\{x\}$, where $P \equiv \langle \ell \rangle$. Then

$$\begin{array}{lll} [\text{nil}] = \lambda \phi. \top & \langle \text{nil} \rangle = \lambda \phi. \perp & \pi_{\text{nil}} = \lambda y. \perp \\ [x :: \ell] = \phi x \wedge [\ell] \phi & \langle x :: \ell \rangle = \phi x \vee \langle \ell \rangle \phi & \pi_{x::\ell} = \lambda y. (x = y) \vee \pi_\ell y \\ [\ell_1 + \ell_2] = [\ell_1] \wedge [\ell_2] & \langle \ell_1 + \ell_2 \rangle = \langle \ell_1 \rangle \vee \langle \ell_2 \rangle & \pi_{\ell_1 + \ell_2} = \pi_{\ell_1} \vee \pi_{\ell_2}, \end{array}$$

so that $[\{\!\{x\}\!\}] = \langle \{\!\{x\}\!\} \rangle = \eta x = \lambda \phi. \phi x$ and $\pi_{\{\!\{x\}\!\}} = \{x\}$. Also

$$(\ell_1 \subset \ell_2) \equiv \forall x \in \ell_1. \exists y \in \ell_2. (x =_X y) \equiv [\ell_1](\lambda x. \langle \ell_2 \rangle \{x\}) \equiv [\ell_1]\pi_2$$

and $(\ell_1 = \ell_2) \equiv (\ell_1 \subset \ell_2) \wedge (\ell_2 \subset \ell_1)$, which are values of type Σ . Finally, the “Curried” operations $\Box \phi \equiv \lambda \ell. [\ell] \phi$ and $\Diamond \phi \equiv \lambda \ell. \langle \ell \rangle \phi$ generate the topology on the powerdomains (Remark 1.5), but are not very useful here.

Notation 3.3 As we want to construct $\mathbf{K}X$ as a subspace of Ω , we need to extend this notation to $L \equiv (N, P) : \Omega \equiv \Sigma^{\Sigma^X} \times \Sigma^{\Sigma^X}$. For the modal notation itself, we simply use $[\]$ and $\langle \ \rangle$ for the product projections, so $\Box \phi L \equiv [L] \phi \equiv$

$N\phi$ and $\diamond \phi L \equiv \langle L \rangle \phi \equiv P\phi$. Then we define

$$\begin{array}{ll}
 \text{nil} & \equiv (\top, \perp) : \Omega \\
 x :: (N, P) & \equiv (x :: N, x :: P) \equiv (\lambda\phi. N\phi \wedge \phi x, \lambda\phi. P\phi \vee \phi x) \\
 (N_1, P_1) + (N_2, P_2) & \equiv (N_1 \wedge N_2, P_1 \vee P_2) \\
 & \equiv (\lambda\phi. N_1\phi \wedge N_2\phi, \lambda\phi. P_1\phi \vee P_2\phi) \\
 x \in L & \equiv \langle L \rangle \{x\} \equiv P\{x\} \equiv \pi x \\
 L_1 \subset L_2 & \equiv [L_1](\lambda x. \langle L_2 \rangle \{x\}) \equiv N_1(\lambda x. P_2\{x\}) \equiv N_1\pi_2 \\
 L_1 \sim L_2 & \equiv (L_1 \subset L_2) \wedge (L_2 \subset L_1) \\
 L \sim \text{nil} & = [L]\perp \equiv N\perp \\
 L \not\sim \text{nil} & = \langle L \rangle \top \equiv P\top.
 \end{array}$$

Clearly $+$ is an associative, commutative, idempotent binary operation on Ω , and nil is a unit for it, with $\text{nil} \subset L$; we shall derive their other algebraic properties of $+$ and \subset shortly. Notice, however, that for P they behave like \vee and \leq , but for N they are like \wedge and \geq . This means that they are not *intrinsic* structure on Ω (as \leq , \vee and \wedge are in Σ), but *imposed* on it by specifying certain maps.

The \wedge/\vee -ambiguity in the $::$ notation will always be resolved by the context, but we shall not risk confusion by further overloading of the important sign for equality.

Plainly not every pair (N, P) will arise as $([\ell], \langle \ell \rangle)$ from a finite subset.

Definition 3.4 We say that the pair $\Gamma \vdash (N, P) : \Omega$ is *modal* if N and P satisfy

$$\begin{array}{ll}
 N\top = \top & P\perp = \perp \\
 N(\phi \wedge \psi) = N\phi \wedge N\psi & P(\phi \vee \psi) = P\phi \vee P\psi \\
 N(\phi \vee \psi) \leq N\phi \vee P\psi & P(\phi \wedge \psi) \geq P\phi \wedge N\psi \\
 N(\lambda x. P\{x\}) = \top & P\phi = \exists x. \phi x \wedge P\{x\}.
 \end{array}$$

The last law, which recovers P from $\pi \equiv \lambda x. P\{x\}$, implies that P preserves \perp , \vee and \exists . Recalling the classical connection between finiteness and Scott continuity, it is perhaps not surprising that Axiom 1.10 provides converses. If P is Scott continuous and preserves \perp and \vee then it also preserves \exists and satisfies the last law. From Scott-continuity of N , Theorem 11.7 shows that the modal laws exactly characterise Kuratowski-finite subsets.

In view of the number of things to be checked, we shall omit most of the proofs that pairs (N, P) are modal, recommending them as exercises. Note that the sixth and fifth laws usually require the Euclidean principle and its

dual respectively.

When the ambient space is Hausdorff instead of discrete (e.g. \mathbb{R}), the seventh law is replaced by $P(\lambda x. N\overline{\{x\}})$, where $\overline{\{x\}} \equiv (\lambda y. x \neq y)$ [1].

Lemma 3.5 Modal operators satisfy the Frobenius law and its dual:

$$\phi : \Sigma^X, \sigma : \Sigma \vdash P(\sigma \wedge \phi) = \sigma \wedge P\phi \quad \text{and} \quad N(\sigma \vee \phi) = \sigma \vee N\phi.$$

Proof By the Phoa principle (Axiom 2.4), since $P\perp = \perp$ and $N\top = \top$. \square

Lemma 3.6 If (N, P) is modal then $P\{x\} \wedge N\phi \leq \phi x$.

Proof

$$\begin{aligned} P\{x\} \wedge N\phi &\leq P(\{x\} \wedge \phi) \\ &\equiv P(\lambda y. (x = y) \wedge \phi y) \\ &= \exists y. P\{y\} \wedge (x = y) \wedge \phi y \\ &= P\{x\} \wedge \phi x \leq \phi x \end{aligned} \quad \square$$

Using these laws we can already recover the algebraic structure of \mathbf{KX} .

Proposition 3.7 Definable finite subsets give rise to modal pairs because

- (a) $\text{nil} \equiv (\top, \perp)$ is modal;
- (b) $x : X \vdash \{x\} \equiv (\lambda \phi. \phi x, \lambda \phi. \phi x)$ is modal;
- (c) if $\Gamma \vdash (N_1, P_1), (N_2, P_2) : \Omega$ are modal then so is $(N_1, P_1) + (N_2, P_2)$. \square

A similar study of the intersection operation raises topological questions.

Proposition 3.8 There is a greatest modal (N, P) iff X is **compact**, in which case $N = \forall_X$, $P = \exists_X$ and $\pi = \top$.

Proof Suppose that there is a bound for all singletons:

$$\lambda x. (\{x\} \subset (N, P)) = \lambda x. [\{x\}](\lambda y. P\{y\}) = \lambda x. P\{x\} = \pi.$$

Hence if (N, P) is the greatest modal pair then $\pi = \top$. In this case,

$$P = \lambda \phi. \exists x. \pi x \wedge \phi x = \lambda \phi. \exists x. \phi x = \exists_X.$$

Also, $x : X, \phi : \Sigma^X \vdash N\phi = N\phi \wedge P\{x\} = \phi x$,

so $\Sigma^{!x}\sigma \equiv \lambda x. \sigma \leq \phi$ iff $\sigma \leq N(\lambda x. \sigma) \leq N\phi \leq \phi x$, which means that $\Sigma^{!x} \dashv N$, i.e. $N = \forall_X$ and X is compact.

Now $N(\lambda x. \exists_X \{x\}) = N(\lambda x. \exists y. x = y) = N\top = \top$ so if X is compact then $(N, P) \subset (\forall_X, \exists_X)$. This pair is modal because, in particular,

$$\exists x. (\phi x \wedge \psi x) \geq \exists x. \phi x \wedge (\forall y. \psi y) \geq (\exists x. \phi x) \wedge (\forall y. \psi y)$$

$$\forall x. (\phi x \vee \psi x) \leq \forall x. \phi x \vee (\exists y. \psi y) \leq (\forall x. \phi x) \vee (\exists y. \psi y)$$

by the Frobenius law and its dual. \square

Proposition 3.9 If $(=_X)$ is decidable (in which case we say that X is **Hausdorff** as well as discrete) and (N_1, P_1) and (N_2, P_2) are modal then so is (N, P) , where

$$\begin{aligned} N\phi &= N_1(\lambda x. \phi x \vee N_2(\lambda y. x \neq y)) \\ P\phi &= P_1(\lambda x. \phi x \wedge P_2(\lambda y. x = y)) \\ &= \exists x. P_1\{x\} \wedge P_2\{x\} \wedge \phi x, \end{aligned}$$

and this is the meet with respect to \subset . Decidability is necessary because, if $\{x\}$ and $\{y\}$ have a meet (N, P) that is modal, then $N\perp = (x \neq y)$, so $(x =_X y)$ is decidable. Intersection of decidable lists can also be defined by a “filtering” program — after we have proved that recursion is valid. \square

The modal laws are also enough to ensure that \sim provides the internal equality for KX .

Proposition 3.10 Let $\Gamma \vdash L_1, L_2 : \Omega$ be modal. Then these are equivalent:

$$(L_1 \subset L_2) = \top \quad \langle L_1 \rangle \leq \langle L_2 \rangle \quad \pi_1 \leq \pi_2 \quad [L_2] \leq [L_1].$$

Proof By the 8th modal law, $P_1 \leq P_2$ iff $\pi_1 \leq \pi_2$. Recall that

$$(L_1 \subset L_2) \equiv [L_1](\lambda x. \langle L_2 \rangle \{x\}) \equiv N_1(\lambda x. P_2\{x\}) \equiv N_1\pi_2.$$

By the 7th modal law this is implied by $\pi_1 \leq \pi_2$ since $\top = N_1\pi_1 \leq N_1\pi_2 \equiv (L_1 \subset L_2)$, or by $N_2 \leq N_1$ since $\top = N_2\pi_2 \leq N_1\pi_2 \equiv (L_1 \subset L_2)$.

Conversely, from $L_1 \subset L_2$, we deduce successively that

$$\begin{array}{lll} \Gamma, \phi : \Sigma^X, N_2\phi = \top & \vdash \pi_2 \leq \phi & \text{Lemma 3.6} \\ \Gamma, N_1\pi_2 = \top & \vdash \pi_1 \leq \pi_2 & \text{similarly} \\ \Gamma, N_1\pi_2 = \top & \vdash P_1 \leq P_2 & \text{8th modal law} \\ \Gamma, N_1\pi_2 = \top, \phi : \Sigma^X, N_2\phi = \top & \vdash \pi_1 \leq \pi_2 \leq \phi & \\ \Gamma, N_1\pi_2 = \top, \phi : \Sigma^X, N_2\phi = \top & \vdash N_1\pi_1 \leq N_1\phi & \text{monotonicity} \\ \Gamma, N_1\pi_2 = \top, \phi : \Sigma^X, N_2\phi = \top & \vdash \top \leq N_1\phi & \text{7th modal law} \\ \Gamma, N_1\pi_2 = \top, \phi : \Sigma^X & \vdash N_2\phi \leq N_1\phi & \text{Axiom 2.4} \\ \Gamma, N_1\pi_2 = \top & \vdash N_2 \leq N_1 & \square \end{array}$$

Corollary 3.11 Amongst modal pairs, \sim provides the internal equality relation, whilst \subset is the *imposed* partial order for which the imposed semilattice structure $+$ is the join.

Proof The relation \subset is reflexive and transitive because the equivalent conditions on $[L]$ or $\langle L \rangle$ are. We have $\Gamma \vdash (L_1 \sim L_2) \equiv (L_1 \subset L_2) \wedge (L_2 \subset L_1) = \top$ iff $\Gamma \vdash [L_1] \geq [L_2]$, $[L_1] \leq [L_2]$, $\langle L_1 \rangle \geq \langle L_2 \rangle$ and $\langle L_1 \rangle \leq \langle L_2 \rangle$, which happens iff $\Gamma \vdash L_1 = L_2 : \Omega$. As for the relationship with $+$,

$$\begin{aligned} (L_1 \subset L_2) = \top &\dashv\vdash \langle L_1 \rangle \leq \langle L_2 \rangle, [L_1] \geq [L_2] \\ &\dashv\vdash \langle L_2 \rangle = \langle L_1 \rangle \vee \langle L_2 \rangle, [L_2] = [L_1] \wedge [L_2] \\ &\dashv\vdash L_2 = L_1 + L_2 \end{aligned} \quad \square$$

Proposition 3.12 If (N, P) are modal then $((N, P) \sim \text{nil})$ is decidable:

$$((N, P) \sim \text{nil}) = N\perp \quad \text{and} \quad ((N, P) \not\sim \text{nil}) = P\top = \exists x. x \in L.$$

Proof From the modal laws, $N\perp \vee P\top \geq N(\perp \vee \top) = N\top = \top$ and $N\perp \wedge P\top \leq P(\perp \wedge \top) = P\perp = \perp$. On the other hand, $((N, P) \sim \text{nil}) = N\perp$ by definition, whilst $P\top = P(\exists x. \{x\}x) = \exists x. P\{x\} = \exists x. x \in L$. \square

Corollary 3.13 $\mathbf{K0} = \mathbf{1}$ and $\mathbf{K1} = \mathbf{2}$.

Proof $\Omega_0 = \Sigma^{\Sigma^0} \times \Sigma^{\Sigma^0} \cong \Sigma \times \Sigma$ and $\Omega_1 = \Sigma^{\Sigma^1} \times \Sigma^{\Sigma^1} \cong \Sigma^\Sigma \times \Sigma^\Sigma$. By the Phoa principle and the constraints $N\top = \top$, $P\perp = \perp$, $N\perp \wedge P\top = \perp$ and $N\perp \vee P\top = \top$, we have $((N, P) \sim (\top, \perp)) \vee ((N, P) \sim (\text{id}, \text{id}))$. \square

Although we have seen that the modal laws characterise the *algebraic* structure of \mathbf{KX} , we still have to show that it is

- (a) a Σ -split subspace of Ω , as modal laws just define an equaliser;
- (b) the free semilattice on X , with induction and recursion, and
- (c) overt, making the quantifier $\exists \ell : \mathbf{KX}$ in Axiom 1.10 legitimate.

We shall do this by developing another characterisation of \mathbf{KX} .

4 Fixed point properties

Since it is the purpose of the paper to *define* the space of Kuratowski-finite subsets, we have to eliminate them from the notation in the previous section. We illustrate this first with the extension of the *existential quantifier* on $\exists : \Sigma^{\mathbf{KX}} \rightarrow \Sigma$ to an operator $\mathbf{E} : \Sigma^\Omega \rightarrow \Sigma$.

Remark 4.1 Transforming $\exists \ell. \theta \ell$ into $\theta \text{nil} \vee \exists x. \exists \ell. \theta(x :: \ell)$, we have

$$\begin{aligned} \exists_{\text{KX}} \theta &\equiv \exists \ell. \theta \ell = \theta \text{nil} \vee \exists x. \exists \ell. \theta(x :: \ell) \\ &= \theta \text{nil} \vee \exists x. \exists_{\text{KX}} (\lambda \ell. \theta(x :: \ell)) \\ &\equiv \theta \text{nil} \vee \exists x. \exists_{\text{KX}} (\mathbb{S}x\theta). \end{aligned}$$

We can therefore define \mathbf{E} as the least fixed point of $\mathbf{E} = \lambda \Theta. \Theta \text{nil} \vee \exists x. \mathbf{E}(\mathbb{S}x\Theta)$, where we write Θ for a typical value of type Σ^Ω or predicate on Ω . Notice that unwinding this fixed point equation reveals the list representation of subsets that we had managed to conceal behind the semilattice structure in the previous section.

Notation 4.2 The *shift operator* $\mathbb{S} : X \times \Sigma^\Omega \rightarrow \Sigma^\Omega$ is defined by

$$\mathbb{S}x\Theta(N, P) \equiv \Theta(x :: N, x :: P) \equiv \Theta(\lambda \phi. N\phi \wedge \phi x, \lambda \phi. P\phi \vee \phi x),$$

and the *exception operator* $\mathcal{S} : X \times \Sigma^{\Sigma^X} \rightarrow \Sigma^{\Sigma^X}$ by

$$\mathcal{S}xN\phi \equiv N(\lambda y. x = y \vee \phi y).$$

Remark 4.3 Similarly, to prove that the embedding $i : \text{KX} \rightarrow \Omega$ is Σ -split, we must also show how to extend *any* predicate θ from KX to Ω , by means of a morphism $I : \Sigma^{\text{KX}} \rightarrow \Sigma^\Omega$. The composite $\mathcal{E} \equiv I \cdot \Sigma^i$ is called a *nucleus* (Definition 5.1), *starting* from which, [B] shows how to *define* KX formally as a subspace of Ω . Like \mathbf{E} , \mathcal{E} will be defined by a fixed point equation, the idea being that

$$I\theta L \equiv \exists \ell. (L \sim \ell) \wedge \theta \ell, \quad \text{so} \quad \mathcal{E}\Theta L \equiv \exists \ell. (L \sim \ell) \wedge \Theta([\ell], \langle \ell \rangle).$$

We can expand this as before, since $((N, P) \sim \ell) = N\pi_\ell \wedge [\ell]\pi_P$:

$$\begin{aligned} \mathcal{E}\Theta(N, P) &= \exists \ell. N\pi_\ell \wedge [\ell]\pi_P \wedge \Theta([\ell], \langle \ell \rangle) \\ &= N\pi_{\text{nil}} \wedge [\text{nil}]\pi_P \wedge \Theta \text{nil} \\ &\quad \vee \exists x. \exists \ell. N\pi_{x::\ell} \wedge [x :: \ell]\pi_P \wedge \Theta([x :: \ell], \langle x :: \ell \rangle) \\ &= (N\perp \wedge \top \wedge \Theta \text{nil}) \\ &\quad \vee \exists x. \exists \ell. \mathcal{S}xN\pi_\ell \wedge P\{x\} \wedge [\ell]\pi_P \wedge \mathbb{S}x\Theta([\ell], \langle \ell \rangle) \\ &= (N\perp \wedge \Theta \text{nil}) \vee \exists x. P\{x\} \wedge \mathcal{E}(\mathbb{S}x\Theta)(\mathcal{S}xN, P). \quad \square \end{aligned}$$

The reasoning that has led up to this fixed point equation depends on the prior existence of lists, so we have to start again from this formula as our “guess” for the definition of the nucleus \mathcal{E} . First let us recall the fixed

point property itself, which follows from Axiom 1.9, but also uses equational induction (Axiom 2.6).

Proposition 4.4 Let $A = \Sigma^U$ and $\Gamma \vdash F : A^A$. Then $\Gamma \vdash \alpha \equiv \exists n. F^n \perp : A$ satisfies $\Gamma \vdash F\alpha = \alpha$. Moreover, if $\Gamma \vdash \beta : A$ is a **pre-fixed point**, that is, $\Gamma \vdash F\beta \leq \beta$, then $\Gamma \vdash \alpha \leq \beta$. \square

Definition 4.5 Let \mathcal{E}_∞ and \mathbf{E}_∞ be the least fixed points of their respective equations above. In the next section we shall prove that \mathcal{E}_∞ is a nucleus on Ω , also satisfying

$$\vdash \mathcal{E}_\infty \leq \text{id} \quad \text{and} \quad x : X, \Theta : \Sigma^\Omega \vdash \mathcal{E}_\infty(\mathbb{S}x\Theta) \leq \mathbb{S}x(\mathcal{E}_\infty\Theta),$$

so we define $\mathsf{K}X \equiv \{\Theta \mid \mathcal{E}_\infty\}$. We call $\Gamma \vdash L : \Omega$ **admissible** if \mathcal{E}_∞ admits it:

$$\Gamma, \Theta : \Sigma^\Omega \vdash \mathcal{E}_\infty\Theta L = \Theta L.$$

In Section 6 we shall show that all *admissible* L are *modal*, so we have the benefit of the algebraic structure that we described in the previous section.

Lemma 4.6 If $\Theta \leq \lambda L. \sigma$ then $\mathbf{E}_\infty\Theta \leq \sigma$.

Proof First, $\mathbf{E}_0\Theta = \Theta \text{nil} \leq \sigma$. Now, if $\mathbf{E} \leq \lambda\Theta. \sigma$ then $\mathbf{E}(\mathbb{S}x\Theta) \leq \sigma$, so

$$\Phi\mathbf{E} \equiv \lambda\Theta. \Theta \text{nil} \vee \exists x. \mathbf{E}(\mathbb{S}x\Theta) \leq \lambda\Theta. \sigma.$$

This amounts to saying that $\Phi(\lambda\Theta. \sigma) \leq (\lambda\Theta. \sigma)$, *i.e.* that $\lambda\Theta. \sigma$ is a pre-fixed point of Φ ; then by Proposition 4.4, the *least* fixed point \mathbf{E}_∞ satisfies $\mathbf{E}_\infty \leq \lambda\Theta. \sigma$. \square

Lemma 4.7 $\mathcal{E}_\infty\Theta L \leq \mathbf{E}_\infty\Theta$.

Proof First, $\mathcal{E}_0\Theta L = \Theta \text{nil} \wedge [L]\perp \leq \Theta \text{nil} = \mathbf{E}_0\Theta \leq \mathbf{E}_\infty\Theta$, so $\mathcal{E}_0 \leq \lambda\Theta L. \mathbf{E}_\infty\Theta$. Now suppose that $\mathcal{E} \leq \lambda\Theta L. \mathbf{E}_\infty\Theta$. Then

$$\begin{aligned} \mathcal{E}\Theta L &= \Theta \text{nil} \wedge [L]\perp \vee \exists x. \mathcal{E}(\mathbb{S}x\Theta)(\mathcal{S}xN, P) \wedge P\{x\} \\ &\leq \mathbf{E}_\infty\Theta \vee \exists x. \mathbf{E}_\infty(\mathbb{S}x\Theta) = \Phi\mathbf{E}_\infty\Theta = \mathbf{E}_\infty\Theta. \end{aligned}$$

Again we have shown that $\lambda\Theta L. \mathbf{E}_\infty\Theta$ is a pre-fixed point of \mathcal{E} , so it is greater than the least fixed point, \mathcal{E}_∞ . \square

Lemma 4.8 $\mathbf{E}_\infty = \mathbf{E}_\infty \cdot \mathcal{E}_\infty$.

Proof First, $\mathbf{E}_0\Theta = \Theta \text{nil} = \mathcal{E}_0\Theta \text{nil} = \mathbf{E}_0(\mathcal{E}_0\Theta) \leq \mathbf{E}_\infty(\mathcal{E}_\infty\Theta)$.

Now suppose that $\mathbf{E} \leq \mathbf{E}_\infty \cdot \mathcal{E}_\infty$. Then

$$\begin{aligned}
 \Phi \mathbf{E} \Theta &= \Theta \text{nil} \vee \exists x. \mathbf{E}(\mathbb{S}x\Theta) \\
 &\leq \mathbf{E}_\infty(\mathcal{E}_\infty \Theta) \vee \exists x. \mathbf{E}_\infty(\mathcal{E}_\infty(\mathbb{S}x\Theta)) && \text{hypothesis} \\
 &\leq \mathbf{E}_\infty(\mathcal{E}_\infty \Theta) \vee \exists x. \mathbf{E}_\infty(\mathbb{S}x(\mathcal{E}_\infty \Theta)) && \text{Definition 4.5} \\
 &= \Phi \mathbf{E}_\infty(\mathcal{E}_\infty \Theta) = \mathbf{E}_\infty(\mathcal{E}_\infty \Theta),
 \end{aligned}$$

i.e. $\mathbf{E}_\infty \cdot \mathcal{E}_\infty$ is a pre-fixed point of Φ , so $\mathbf{E}_\infty \leq \mathbf{E}_\infty \cdot \mathcal{E}_\infty$, but they are equal since $\mathcal{E}_\infty \leq \text{id}$. \square

Proposition 4.9 $KX \subset \Omega$ is overt (not open), with existential quantifier and \Diamond -modal operator

$$\exists_{KX} \equiv \mathbf{E} \cdot I \dashv \Sigma^{\text{!}KX} \quad \text{and} \quad \mathbf{E} = \exists_{KX} \cdot \Sigma^i.$$

Proof We require $\theta \leq \lambda \ell$. σ iff $\exists \ell. \theta \ell \leq \sigma$. Forwards, $\Theta \leq \lambda L$. $\sigma \vdash \mathbf{E}_\infty \Theta \leq \sigma$ (Lemma 4.6). Conversely, $\mathcal{E}_\infty \Theta L \leq \mathbf{E}_\infty \Theta = \mathbf{E}_\infty(\mathcal{E}_\infty \Theta)$ (Lemmas 4.7 and 4.8). From this we also have $\exists_{KX} \cdot \Sigma^i = \mathbf{E}_\infty \cdot I \cdot \Sigma^i = \mathbf{E}_\infty \cdot \mathcal{E}_\infty = \mathbf{E}_\infty$. \square

Besides equational and fixed point induction (Axiom 2.6 and Proposition 4.4), we shall need two other principles in the rest of the paper. The idea of the first one is that $\Theta_\top L$ means $\exists \ell. L \sim \ell$, and in fact $\Theta_\top = \mathcal{E}_\infty \top$.

Lemma 4.10 Let $\vdash \Theta_\top : \Sigma^\Omega$ be the least solution of

$$\Theta_\top(N, P) = N \perp \vee \exists x. \Theta_\top(\mathbb{S}xN, P) \wedge P\{x\}.$$

Then $L : KX \vdash \Theta_\top L = \top$, *i.e.* $\Theta_\top(N, P) = \top$ for all admissible (N, P) . \square

Proposition 4.11 We have the induction scheme

$$\frac{\Gamma \vdash \Theta : \Sigma^\Omega \quad \Gamma \vdash \Theta \text{nil} = \top \quad \frac{\Gamma, L : \Omega, \Theta L = \top \vdash \Theta(x :: L) = \top}{\Gamma \vdash \Theta \leq \mathbb{S}x\Theta}}{\Gamma, L : KX \vdash \Theta L = \top}$$

Proof

$$\begin{aligned}
 \Theta(N, P) &\geq \mathcal{E}_\infty \Theta(N, P) \\
 &= \mathcal{E}_0 \Theta(N, P) \vee \exists x. \mathcal{E}_\infty(\mathbb{S}x\Theta)(\mathbb{S}xN, P) \wedge P\{x\} \\
 &\geq N \perp \vee \exists x. \mathcal{E}_\infty \Theta(\mathbb{S}xN, P) \wedge P\{x\}
 \end{aligned}$$

but Θ_\top is the least (pre)fixed point of this, so $\Theta_\top \leq \Theta$. Hence if (N, P) is admissible then $\Theta(N, P) \geq \Theta_\top(N, P) = \top$. \square

Although the form of this result is very familiar, its usefulness is rather limited, as it only tells us about *open* subspaces. Ultimately, Corollary 10.12 will provide an induction scheme of the form that *any* subspace that includes *nil* and is closed under $::$ is the whole space. The principle that we shall invoke repeatedly in Section 6 is the following.

Proposition 4.12 Let $\Gamma \vdash \Theta, \Phi : \Sigma^\Omega$ (*i.e.* particular Θ and Φ are given). Then

$$\frac{\Gamma, L : \Omega \vdash \mathcal{E}_0 \Theta L \leq \Phi L \quad \Gamma, \mathcal{E} \leq \text{id nucleus}, L : \Omega, \mathcal{E} \Theta L \leq \Phi L \vdash \$\mathcal{E} \Theta L \leq \Phi L}{\Gamma, L : \mathsf{K}X \vdash \Theta L \leq \Phi L}$$

i.e. $\Theta L \leq \Phi L$ for all admissible L .

Proof By equational induction (Axiom 2.6), $\Gamma, n : \mathbb{N} \vdash \$^n \mathcal{E}_0 \Theta \leq \Phi$, so $\Gamma \vdash \mathcal{E}_\infty \Theta = \exists n. \$^n \mathcal{E}_0 \Theta \leq \Phi$. Hence if L is admissible then $\Theta L = \mathcal{E}_\infty \Theta L \leq \Phi L$. \square

5 Stages in the construction

Definition 5.1 Recall from [B] that $\Gamma \vdash \mathcal{E} : \Sigma^\Omega \rightarrow \Sigma^\Omega$ is a **nucleus** on Ω if

$$\Gamma, H : \Sigma^3 \Omega \vdash \mathcal{E}(\lambda L : \Omega. H(\lambda \Theta : \Sigma^\Omega. (\mathcal{E} \Theta L)))L = \mathcal{E}(\lambda L : \Omega. H(\lambda \Theta : \Sigma^\Omega. \Theta L))L.$$

Beware that we have shamelessly appropriated this word from locale theory, in which a nucleus j satisfies $\text{id} \leq j \leq j^2$. Nuclei in ASD need not in general be order-related to id , but those in this paper will satisfy $\mathcal{E} \leq \text{id}$.

A nucleus $\Gamma \vdash \mathcal{E} : \Sigma^\Omega \rightarrow \Sigma^\Omega$ on Ω **admits** a term $\Gamma \vdash L : \Omega$ if

$$\Gamma, \Theta : \Sigma^\Omega \vdash \mathcal{E} \Theta L = \Theta L \quad \text{or} \quad \Gamma \vdash \lambda \Theta. \mathcal{E} \Theta L = \lambda \Theta. \Theta L.$$

If the context Γ is empty, [B] allows us to introduce a subtype $i : \{\Omega \mid \mathcal{E}\} \rightarrow \Omega$ with Σ -splitting I for which $\mathcal{E} = I \cdot \Sigma^i$. Then L belongs to the subtype iff \mathcal{E} admits it. In this paper we shall need to define nuclei in non-trivial contexts, but we will not introduce the corresponding dependent types.

Definition 5.2 Following Remark 4.3, we construct the subspace $\mathsf{K}X \subset \Omega$ using the nucleus \mathcal{E}_∞ that is defined as the least fixed point of the operator $\$: (\Sigma^\Omega \rightarrow \Sigma^\Omega) \rightarrow (\Sigma^\Omega \rightarrow \Sigma^\Omega)$, where

$$\$ \mathcal{E} \Theta(N, P) = \mathcal{E}_0 \Theta(N, P) \vee \exists x. \mathcal{E}(\$x \Theta)(\$x N, P) \wedge P\{x\}$$

and $\mathcal{E}_0 \equiv \$\perp \equiv \lambda\Theta L. [L]\perp \wedge \Theta \text{nil} \equiv \lambda\Theta NP. N\perp \wedge \Theta(\top, \perp)$.

The objects $X^{(n)} \equiv \{\Omega \mid \$^n \mathcal{E}_0\}$ that are obtained as the successive unwindings of this equation intuitively represent the collections of subsets of X with at most n elements.

Lemma 5.3 $\mathcal{F}\top \wedge F\perp \leq \mathcal{F}F$.

Proof Apply Axiom 2.4 to $F\perp = \top \vdash \top = F \vdash \mathcal{F}\top = \mathcal{F}F$. \square

This result shows how the singleton $\{\text{nil}\}$ is embedded as a subspace of Ω .

Lemma 5.4 The singleton $i : \{\text{nil}\} \longrightarrow \Omega$ is a Σ -split subspace, where

$$\begin{aligned} \sigma : \Sigma^{\{\text{nil}\}} \vdash I\sigma &\equiv \lambda NP. \sigma \wedge N\perp \\ \Theta : \Sigma^\Omega \vdash \mathcal{E}_0\Theta &\equiv I(\Sigma^i\Theta) = \lambda NP. \Theta \text{nil} \wedge N\perp. \end{aligned}$$

Moreover $\mathcal{E}_0 \leq \text{id}$, so $I \dashv \Sigma^i$, and if \mathcal{E} is any nucleus on Ω that admits nil then $\mathcal{E}_0 \leq \mathcal{E}$. \square

Lemma 5.5 $\Theta(N, x :: P) \wedge P\{x\} = \Theta(N, P) \wedge P\{x\}$
 $(x :: \mathcal{S}xN) = (x :: N) \leq N \leq \mathcal{S}xN = \mathcal{S}x(x :: N)$
 $P \leq (x :: P) \quad (x :: P)\{x\} = \top \quad P\{x\} \wedge \phi x \leq P\phi$

Proof Apply Axiom 2.4 to

$$\phi x = \top \vdash N\phi = \mathcal{S}xN\phi, \{x\} \leq \phi \quad \text{and} \quad P\{x\} = \top \vdash x :: P = P. \quad \square$$

Lemma 5.6 $\mathcal{E}(\mathbb{S}x\Theta) \leq \mathbb{S}x(\$ \mathcal{E}\Theta)$. \square

Lemma 5.7 If $\mathcal{E} \leq \text{id}$ then $\$ \mathcal{E} \leq \text{id}$. \square

Lemma 5.8 If \mathcal{E} is a nucleus with $\mathcal{E} \leq \text{id}$ then so is $\$ \mathcal{E}$.

Proof Expanding the definition of the *outer* $\$ \mathcal{E}$, in which $\mathsf{H} : \Sigma^3\Omega$,

$$\begin{aligned} &\$ \mathcal{E}(\lambda L. \mathsf{H}(\lambda \Theta. (\underline{\$ \mathcal{E}}\Theta L)))(N, P) \\ &= \mathcal{E}_0(\lambda L. \mathsf{H}(\lambda \Theta. (\underline{\$ \mathcal{E}}\Theta L))) \\ &\quad \vee \exists x. \mathcal{E}(\mathbb{S}x(\lambda L. \underline{\$ \mathcal{E}}\mathsf{H}(\lambda \Theta. \Theta L)))(\mathcal{S}xN, P) \wedge P\{x\}, \end{aligned}$$

and we have to show that we may delete the *inner* ones. First note that $\Theta \text{nil} \leq \$ \mathcal{E}\Theta \text{nil} \leq \Theta \text{nil}$, by Definition 5.2 and Lemma 5.7. Then

$$\mathcal{E}_0(\lambda L'. \mathsf{H}(\lambda \Theta. \$ \mathcal{E}\Theta L'))$$

$$\begin{aligned}
&= \lambda L. (\lambda L'. \mathsf{H}(\lambda \Theta. \$\mathcal{E}\Theta L')) \mathsf{nil} \wedge [L] \perp \\
&= \lambda L. \mathsf{H}(\lambda \Theta. \$\mathcal{E}\Theta \mathsf{nil} \wedge [L] \perp) \\
&= \lambda L. \mathsf{H}(\lambda \Theta. \Theta \mathsf{nil}) \wedge [L] \perp && \text{above} \\
&= \lambda L. (\lambda L'. \mathsf{H}(\lambda \Theta. \Theta L')) \mathsf{nil} \wedge [L] \perp \\
&= \mathcal{E}_0(\lambda L'. \mathsf{H}(\lambda \Theta. \Theta L')) \\
&\mathcal{E}(\mathbb{S}x(\lambda L. \mathsf{H}(\lambda \Theta. \Theta L))) \\
&\geq \mathcal{E}(\mathbb{S}x(\lambda L. \mathsf{H}(\lambda \Theta. \$\mathcal{E}\Theta L))) && \$\mathcal{E} \leq \mathsf{id} \\
&= \mathcal{E}(\lambda L. \mathsf{H}(\lambda \Theta. \$\mathcal{E}\Theta(x :: L))) && \text{def } \mathbb{S}x \\
&\geq \mathcal{E}(\lambda L. \mathsf{H}(\lambda \Theta. \mathcal{E}(\mathbb{S}x\Theta)L)) && \text{Lemma 5.6} \\
&= \mathcal{E}(\lambda L. (\mathsf{H} \cdot \Sigma^{\mathbb{S}x})(\lambda \Theta. \mathcal{E}\Theta L)) && \text{def } \Sigma^{(-)} \\
&= \mathcal{E}(\lambda L. (\mathsf{H} \cdot \Sigma^{\mathbb{S}x})(\lambda \Theta. \Theta L)) && \mathcal{E} \text{ nucleus wrt } \mathsf{H} \cdot \Sigma^{\mathbb{S}x} \\
&= \mathcal{E}(\lambda L. \mathsf{H}(\lambda \Theta. (\mathbb{S}x\Theta)L)) && \text{def } \Sigma^{(-)} \\
&= \mathcal{E}(\lambda L. \mathsf{H}(\lambda \Theta. \Theta(x :: L))) && \text{def } \mathbb{S}x \\
&= \mathcal{E}(\mathbb{S}x(\lambda L. \mathsf{H}(\lambda \Theta. \Theta L))) && \text{def } \mathbb{S}x \quad \square
\end{aligned}$$

As we have said, the ASD calculus does not currently allow us to define $X^{(n)}$ as a *type* dependent on n , but we can at least introduce $\$^n\mathcal{E}_0$ as a *term* dependent on n .

Proposition 5.9 $n : \mathbb{N} \vdash \$^n\mathcal{E}_0 : \Sigma^\Omega \rightarrow \Sigma^\Omega$ are nuclei with $n : \mathbb{N} \vdash \$^n\mathcal{E}_0 \leq \$^{n+1}\mathcal{E}_0 \leq \mathsf{id}$.

Proof The term $n : \mathbb{N} \vdash \$^n\mathcal{E}_0$ is formed by recursion. The base case of the induction is that $\vdash \mathcal{E}_0 \leq \mathsf{id}$ is a nucleus, which is Lemma 5.4. The induction step,

$$\mathcal{E} : (\Sigma^\Omega)^{(\Sigma^\Omega)}, \mathcal{E} \leq \mathsf{id}, \mathcal{E} \text{ nucleus} \vdash \mathcal{E} \leq \$\mathcal{E} \leq \mathsf{id}, \$\mathcal{E} \text{ nucleus},$$

was proved in Lemmas 5.7 and 5.8; it has equations as hypotheses and conclusions. From these things we may deduce the result by equational induction on \mathbb{N} (Axiom 2.6). \square

Proposition 5.10 $\vdash \mathcal{E}_\infty \equiv \exists n. \$^n\mathcal{E}_0$ is a nucleus with $\mathcal{E}_\infty \leq \mathsf{id}$.

Proof By the previous result, $\$^n\mathcal{E}_0 = \$^{n+1}\perp$ is an ascending chain of nuclei with $\$^n\mathcal{E}_0 \leq \mathsf{id}$. Writing

$$\mathcal{E} : (\Sigma^\Omega)^{(\Sigma^\Omega)} \vdash F\mathcal{E}, G\mathcal{E} : \Sigma$$

for the two sides of Definition 5.1 for a nucleus, Axiom 1.9 gives

$$\begin{aligned} \vdash F\mathcal{E}_\infty &= F(\exists n. \$^n \mathcal{E}_0) = \exists n. F(\$^n \mathcal{E}_0) \\ &= \exists n. G(\$^n \mathcal{E}_0) = G(\exists n. \$^n \mathcal{E}_0) = G\mathcal{E}_\infty, \end{aligned}$$

so \mathcal{E}_∞ is a nucleus. Lemma 5.7 showed that $\$id \leq id$, so id is a pre-fixed point of $\$$, whence $\mathcal{E}_\infty \leq id$ by Proposition 4.4. \square

This justifies Definition 4.5 and so the other results of the last section, apart from showing that all definable finite subsets give rise to admissible pairs.

Lemma 5.11 $\$ \mathcal{E}$ admits nil , since $\Theta nil \leq \$ \mathcal{E} \Theta nil \leq \Theta nil$. \square

Lemma 5.12 If \mathcal{E} admits L then $\$ \mathcal{E}$ admits $x :: L$.

Proof

$$\begin{aligned} \Theta(x :: L) &= (\$x\Theta)L = \mathcal{E}(\$x\Theta)L \\ &\leq \$x(\$ \mathcal{E} \Theta)L = \$ \mathcal{E} \Theta(x :: L) \\ &\leq (\$x\Theta)L. \end{aligned}$$

So $\Theta : \Sigma^\Omega \vdash \Theta(x :: L) = \mathcal{E} \Theta(x :: L)$ as required. \square

Lemma 5.13 If $\vdash \mathcal{E} \leq id$ is a nucleus then the pair

$$\mathbf{1} \xrightarrow{nil} \{\Omega \mid \$ \mathcal{E}\} \xleftarrow{::} X \times \{\Omega \mid \mathcal{E}\}$$

is jointly Σ -split epi.

Proof We have just shown that the two maps are well defined. In the diagram

$$\begin{array}{ccc} \{\Omega \mid \$ \mathcal{E}\} & \xleftarrow{\quad nil, :: \quad} & \mathbf{1} + X \times \{\Omega \mid \mathcal{E}\} \\ \Sigma^{\{\Omega \mid \$ \mathcal{E}\}} & \xrightleftharpoons{\quad \Sigma nil, :: \quad} & \Sigma \times \Sigma^{X \times \{\Omega \mid \mathcal{E}\}} \\ \uparrow \$ \mathcal{E} \quad \downarrow & & \uparrow id \times \mathcal{E}^X \quad \downarrow \\ \Sigma^\Omega & \xrightleftharpoons[\lambda L. \sigma \wedge [L] \perp \vee \exists x. \Phi x(\mathcal{S}x[L], \langle L \rangle) \wedge \langle L \rangle \{x\} \leftarrow (\sigma, \Phi)]{\Theta \mapsto (\Theta nil, \lambda x L. \mathcal{E}(\$x\Theta)L)} & \Sigma \times \Sigma^{X \times \Omega} \end{array}$$

we have to show that the inverse image map is given by the formula shown, and is split mono.

Let $\Theta : \Sigma^{\{\Omega | \mathcal{E}\}}$, $x : X$ and $L : \{\Omega | \mathcal{E}\}$, i.e. \mathcal{E} admits L . This means that

$$\Theta(x :: L) = \mathbb{S}x\Theta L = \mathcal{E}(\mathbb{S}x\Theta)L = \Phi xL,$$

which justifies the inverse image map. Now the composite takes

$$\begin{aligned} \Theta &\mapsto (\sigma, \Phi) \mapsto \lambda L. \sigma \wedge [L] \perp \vee \exists x. \Phi x(\mathcal{S}x[L], \langle L \rangle) \wedge \langle L \rangle \{x\} \\ &= \lambda L. \Theta \text{ nil} \wedge [L] \perp \vee \exists x. \mathcal{E}(\mathbb{S}x\Theta)(\mathcal{S}x[L], \langle L \rangle) \wedge \langle L \rangle \{x\}, \end{aligned}$$

which is $\mathcal{E}\mathcal{E}\Theta$, but the hypothesis on Θ says that $\Theta = \mathcal{E}\mathcal{E}\Theta$. \square

Hence $\mathbf{1} + X \twoheadrightarrow X^{(1)}$, $\mathbf{1} + X + X^2 \twoheadrightarrow X^{(2)}$, $\mathbf{1} + X + X^2 + X^3 \twoheadrightarrow X^{(3)}$, ...

Proposition 5.14 The pair $\mathbf{1} \xrightarrow{\text{nil}} \mathbb{K}X \xleftarrow{\cdot} X \times \mathbb{K}X$ is jointly Σ -split epi. \square

6 Admissible implies modal

Now we use the induction scheme in Proposition 4.12 to prove that all admissible pairs $\Gamma \vdash (N, P) : \mathbb{K}X$ satisfy the modal laws, starting with the properties of the “possibility” operator P .

Lemma 6.1 $\Sigma^X \triangleleft \Sigma^{\Sigma^X}$ by $\pi \mapsto \lambda \phi. \exists x. \phi x \wedge \pi x$ and $P \mapsto \lambda x. P\{x\}$. \square

Indeed, if (N, P) is admissible then P is recovered from π :

Lemma 6.2

$$L : \mathbb{K}X, \phi : \Sigma^X \vdash \langle L \rangle \phi = \exists x. \langle L \rangle \{x\} \wedge \phi x \equiv \exists x. \pi x \wedge \phi x \equiv \exists x. x \in L \wedge \phi x.$$

Proof $(\exists x. P\{x\} \wedge \phi x) \leq P\phi$ by Lemma 5.5, so we have to prove \geq .

Consider $\Theta \equiv \lambda N P. P\phi$ and $\Phi \equiv \lambda N P. \exists y. P\{y\} \wedge \phi y$ in the context $\Gamma \equiv [\phi : \Sigma^X]$. Then $\vdash \mathcal{E}_0\Theta \leq \Phi$ and $\mathcal{E}, \mathcal{E}\Theta \leq \Phi \vdash \mathcal{E}\mathcal{E}\Theta \leq \Phi$. Hence $L : \mathbb{K}X, \phi : \Sigma^X \vdash \langle L \rangle \phi \equiv \Theta L \leq \Phi L \equiv \exists x. \langle L \rangle \{x\} \wedge \phi x$ by Proposition 4.12. \square

Corollary 6.3 $\langle L \rangle \perp = \perp$, $\langle L \rangle (\phi \vee \psi) = \langle L \rangle \phi \vee \langle L \rangle \psi$ $\langle L \rangle (\phi \wedge \pi) = \langle L \rangle \phi$. \square

Next we consider the “necessity” operator N .

Lemma 6.4 $(N, P) : \mathbb{K}X \vdash N\pi \equiv N(\lambda x. P\{x\}) = \top$, so $(N, P) \sim (N, P)$.

Proof Consider $\Theta(N, P) = N(\lambda x. P\{x\})$. Then $\Theta \text{nil} = \top$ and

$$\begin{aligned} (\mathbb{S}x\Theta)(N, P) &= (x :: N)(\lambda y. (x :: P)\{y\}) \\ &= N(\lambda y. P\{y\} \vee y = x) \wedge (P\{x\} \vee x = x) \\ &\geq N(\lambda y. P\{y\}) = \Theta(N, P), \end{aligned}$$

so by Proposition 4.11, $H(N, P) = \top$ for all admissible (N, P) . \square

Lemma 6.5 If $L : \mathsf{KX}$ then $[L]$ preserves finite meets.

Proof We can show that $N\top = \top$ using $\Theta(N, P) \equiv N\top$ in Proposition 4.11. For $N\phi \wedge N\psi \leq N(\phi \wedge \psi)$ consider $\Theta(N, P) \equiv N\phi \wedge N\psi$ and $\Phi(N, P) \equiv N(\phi \wedge \psi)$ in Proposition 4.12. \square

Corollary 6.6 $[L]\top = \top$, $[L](\phi \wedge \psi) = [L]\phi \wedge [L]\psi$ and $[L](\phi \wedge \pi) = [L]\phi$.

Proof The last uses Lemma 6.4. \square

Lemma 6.7 $L : \mathsf{KX}$, $x : X$, $\phi : \Sigma^X \vdash x \in L \wedge [L]\phi \leq \phi x$, where $(x \in L) \equiv \langle L \rangle\{x\}$.

Proof With $L = (N, P)$, this says that $N\phi \wedge P\{x\} \leq \phi x$.

Consider $\Theta(N, P) = N\phi \wedge P\{x\}$ and $\Phi(N, P) = \phi x$ in Proposition 4.12. \square

Corollary 6.8 $[L](\phi \vee \psi) \leq [L]\phi \vee \langle L \rangle\psi$ and $\langle L \rangle(\phi \wedge \psi) \geq \langle L \rangle\phi \wedge [L]\psi$. \square

Theorem 6.9 KX is overt discrete and has no proper open subalgebra for nil and $::$.

Proof We have just shown that all $L : \mathsf{KX}$ are modal. For modal L_1, L_2 , Proposition 3.10 said that $L_1 = L_2 : \Omega$ iff $(L_1 \sim L_2) = \top$, which is an open equivalence relation. Proposition 4.9 said that KX is overt. Lemmas 5.11–5.12 provided the algebra structure and Corollary 4.11 said that this is minimal. \square

Unfortunately $+$ is missing: we are not yet in a position to show that $L_1 + L_2$ is *admissible* when L_1 and L_2 are. This is because if $L_1 : X^{(n)}$ and $L_2 : X^{(m)}$ we would expect $L_1 + L_2 : X^{(n+m)}$, whereas our method of induction only takes us from n to $n+1$. We shall show that $+$ is well defined in Lemma 10.9, using the recursion scheme for nil and $::$, but that is still a long way ahead.

7 K as a functor

We must show how K acts on $f : X \rightarrow Y$ between overt discrete objects, but we shall also prove that it preserves monos and inverse images, and that it takes coproducts ($X = Y + Z$) to products ($KY \times KZ$). This means that any Kuratowski-finite subset $L : KX$ may be *partitioned* between Y and Z . (We already know from Corollary 3.13 that $K0 = 1$.)

Remark 7.1 Recall that $f : X \rightarrow Y$ acts contravariantly on predicates, turning $\psi : \Sigma^Y$ into $\Sigma^f \psi \equiv \lambda x. \psi(fx) : \Sigma^X$, but covariantly on modal operators, so $P : \Sigma^{\Sigma^X}$ becomes

$$\Sigma^2 f P \equiv \Sigma^{\Sigma^f} P \equiv P \cdot \Sigma^f \equiv \lambda \psi. P(\Sigma^f \psi) \equiv \lambda \psi. P(\lambda x. \psi(fx)),$$

i.e. the composite with the *inverse* image Σ^f along f . In the case of an inclusion $i : Y \hookrightarrow X$, we may similarly compose with the *direct* image $\exists_i \dashv \Sigma^i$, *i.e.*

$$(P \cdot \exists_i) \psi \equiv P(\lambda x. \exists y. x = iy \wedge \psi y).$$

In the case of the coproduct, $Y \subset X = Y + Z$ is also closed, with $\Sigma^i \dashv \forall_i$. For $\psi : \Sigma^Y$ and $y : Y$, $\forall_i \psi(iy) = \psi y$, whilst for $z : Z$, $\forall_i \psi(jz) = \top$, so

$$(N \cdot \forall_i) \psi = N(\lambda x. (\exists y. x = iy \wedge \psi y) \vee (\exists z. x = jz)).$$

Locale theorists will recognise this construction as a special case of the relationship with open proper maps that we shall discuss in Section 11.

Lemma 7.2 If $\Gamma \vdash (N, P) : \Omega_X$ is modal then so are $\Gamma \vdash Kf(N, P) \equiv (N \cdot \Sigma^f, P \cdot \Sigma^f)$ and $Ki^{-1}(N, P) \equiv (N \cdot \forall_i, P \cdot \exists_i) : \Omega_Y$. \square

But we have to show that they are *admissible* to KY whenever $L : KX$. Writing $\mathcal{R}(N, P)$ for either of them, we shall do the two cases in parallel, the analogy being that the second is for a *partial* map $f : X \rightarrow Y$ with decidable support (the “inverse” of $i : Y \hookrightarrow X$). In Lemma 7.9 we shall have to expand an $\exists x$ in the definition of $\$ \mathcal{E}_X$; the second case has two sub-cases, in which $x = iy$ (so $y = fx$) and $x = jz$ (so fx is undefined). The following lemmas therefore have *three* cases. We write $\mathcal{E}_X^{(n)} = \$^n \mathcal{E}_0$ and $\mathcal{E}_X^{(\infty)} = \exists n. \$^n \mathcal{E}_0$ for the nuclei on Ω_X and similarly for Y .

In all three cases, \mathcal{R} preserves the empty subset and unions:

Lemma 7.3 $\mathcal{R} \text{ nil}_X = \text{nil}_Y$, so $[L] \perp \leq [\mathcal{R}L] \perp$ and $\mathcal{E}_X^{(0)}(\Theta \cdot \mathcal{R}) \leq (\mathcal{E}_Y^{(0)} \Theta) \cdot \mathcal{R}$.

Proof $\Sigma^f \top = \forall_i \top = \top$ and $\Sigma^f \perp = \exists_i \perp = \perp$.

$[\mathcal{R}L] \perp = [L](H \perp) \geq [L] \perp$ where $H = \Sigma^f$ or \forall_i .

$$\mathcal{E}_X^{(0)}(\Theta \cdot \mathcal{R})L = \Theta(\mathcal{R} \text{ nil}) \wedge [L] \perp \geq \Theta \text{ nil} \wedge [\mathcal{R}L] \perp = (\mathcal{E}_Y^{(0)}\Theta)(\mathcal{R}L). \quad \square$$

Lemma 7.4 $L_1, L_2 : \Omega_X \vdash \mathcal{R}(L_1 + L_2) = \mathcal{R}L_1 + \mathcal{R}L_2$,

so $+$ on Ω is natural in $f : X \rightarrow Y$.

Proof $(N_1 \wedge N_2) \cdot H = N_1 \cdot H \wedge N_2 \cdot H$ and $(P_1 \vee P_2) \cdot H = P_1 \cdot H \vee P_2 \cdot H$, where $H = \Sigma^f$, \forall_i or \exists_i . \square

\mathcal{R} either applies the function f to individual elements x , or “filters” the defined values $y = fx = f(iy)$ from the undefined ones with $x = jz$.

Lemma 7.5 $\mathcal{R}\{\!\{x}\!\} = \{\!\{fx}\!\}$, $\mathcal{R}\{\!\{iy}\!\} = \{\!\{y}\!\}$ and $\mathcal{R}\{\!\{jz}\!\} = \text{nil}$.

Proof We do \square , the proof for \diamond being similar.

$$[\{\!\{x}\!\}](\Sigma^f\psi) = \eta x(\Sigma^f\psi) = (\Sigma^f\psi)x = \psi(fx) = \eta(fx)\psi = [\{\!\{fx}\!\}]\psi$$

$$[\{\!\{iy}\!\}](\forall_i\psi) = \eta(iy)(\forall_i\psi) = (\forall_i\psi)(iy) = \psi y = \eta y\psi = [\{\!\{y}\!\}]\psi$$

$$[\{\!\{jz}\!\}](\forall_i\psi) = \eta(jz)(\forall_i\psi) = (\forall_i\psi)(jz) = \top = [\text{nil}]\psi. \quad \square$$

Corollary 7.6 $\mathbb{S}x(\Theta \cdot \mathcal{R}) = (\mathbb{S}(fx)\Theta) \cdot \mathcal{R}$, $\mathbb{S}(iy)(\Theta \cdot \mathcal{R}) = (\mathbb{S}y\Theta) \cdot \mathcal{R}$ and $\mathbb{S}(jz)(\Theta \cdot \mathcal{R}) = \Theta \cdot \mathcal{R}$.

Proof $\Theta \cdot \mathcal{R}(\{\!\{x}\!\} + L) = \Theta(\mathcal{R}\{\!\{x}\!\} + \mathcal{R}L)$, which is respectively $\Theta(\{\!\{fx}\!\} + \mathcal{R}L)$, $\Theta(\{\!\{y}\!\} + \mathcal{R}L)$ and $\Theta(\text{nil} + \mathcal{R}L)$. \square

Lemma 7.7 $P\{x\} \leq (P \cdot \Sigma^f)\{fx\}$, $P\{iy\} = (P \cdot \exists_i)\{y\}$ and $P\{jz\} = (P \cdot \exists_j)\{z\}$.

Proof $\{x\}x' = (x = x') \leq (fx = fx') = \Sigma^f\{fx\}x'$

and $\{iy\}x' = (iy = x') = \exists y'. (y = y') \wedge (iy' = x') = \exists_i\{y\}x'$. \square

Lemma 7.8 “ $fL \setminus fx \subset f(L \setminus x)$ ” in the sense that

$(\mathcal{S}xN) \cdot \Sigma^f \leq \mathcal{S}(fx)(N \cdot \Sigma^f)$, so $\mathcal{R}(\mathcal{S}xN, P) \leq \mathcal{S}(fx)(\mathcal{R}(N, P))$,

$(\mathcal{S}(iy)N) \cdot \forall_i = \mathcal{S}y(N \cdot \forall_i)$, so $\mathcal{R}(\mathcal{S}(iy)N, P) = \mathcal{S}y(\mathcal{R}(N, P))$,

and $(\mathcal{S}(jz)N) \cdot \forall_i = (N \cdot \forall_i)$, so $\mathcal{R}(\mathcal{S}(jz)N, P) = \mathcal{R}(N, P)$.

Proof

$$((\mathcal{S}xN) \cdot \Sigma^f)\psi = N(\lambda x'. x' = x \vee \psi(fx'))$$

$$\leq N(\lambda x'. fx' = fx \vee \psi(fx'))$$

$$= N(\Sigma^f(\lambda y. y = fx \vee \psi y))$$

$$= (\mathcal{S}(fx)(N \cdot \Sigma^f))\psi$$

$$((\mathcal{S}(iy)N) \cdot \forall_i)\psi = N(\lambda x. x = iy \vee \forall_i\psi x)$$

$$= N(\lambda x. \exists y'. x = iy' \wedge (iy' = iy \vee \forall_i\psi(iy')))$$

$$\begin{aligned}
& \vee \exists z'. x = jz' \wedge (jz' = iy \vee \forall_i \psi(jz')) \\
&= N(\lambda x. \exists y'. x = iy' \wedge (y' = y \vee \forall_i \psi y')) \\
& \quad \vee \exists z'. x = jz' \wedge (\perp \vee \top) \\
&= (N \cdot \forall_i)(\lambda y'. y = y' \vee \psi y') \\
&= \mathcal{S}y(N \cdot \forall_i)\psi \\
&((\mathcal{S}(jz)N) \cdot \forall_i)\psi = N(\lambda x. x = jz \vee \forall_i \psi x) \\
&= N(\lambda x. \exists y'. x = iy' \wedge (iy' = jz \vee \forall_i \psi(iy'))) \\
& \quad \vee \exists z'. x = jz' \wedge (jz' = jz \vee \forall_i \psi(jz')) \\
&= N(\lambda x. \exists y'. x = iy' \wedge (\perp \vee \psi y')) \\
& \quad \vee \exists z'. x = jz' \wedge (z = z' \vee \top) \\
&= N(\lambda x. \exists y'. x = iy' \wedge \psi y') \vee \exists z'. x = jz' \\
&= (N \cdot \forall_i)\psi \quad \square
\end{aligned}$$

Lemma 7.9 $\lambda\Phi. \mathcal{E}_X(\Phi \cdot \mathcal{R}) \leq \lambda\Phi. (\mathcal{E}_Y\Phi) \cdot \mathcal{R} \vdash \lambda\Theta. \$\mathcal{E}_X(\Theta \cdot \mathcal{R}) \leq \lambda\Theta. (\$ \mathcal{E}_Y\Theta) \cdot \mathcal{R}.$

Proof We expand $\$ \mathcal{E}_X$ (Definition 5.2). By Lemma 7.3,

$$\begin{aligned}
\$ \mathcal{E}_X(\Theta \cdot \mathcal{R})(N, P) &= \mathcal{E}_X^{(0)}(\Theta \cdot \mathcal{R})(N, P) \vee \exists x. (\dots) \\
&\leq (\mathcal{E}_Y^{(0)}\Theta)(\mathcal{R}(N, P)) \vee \exists x. (\dots),
\end{aligned}$$

where the term (\dots) has three cases:

In the first, for f , use Corollary 7.6 and the premise with $\Phi \equiv \mathbb{S}(fx)\Theta$:

$$\mathcal{E}_X(\mathbb{S}x(\Theta \cdot \mathcal{R})) \leq \mathcal{E}_X((\mathbb{S}(fx)\Theta) \cdot \mathcal{R}) \leq (\mathcal{E}_Y(\mathbb{S}(fx)\Theta)) \cdot \mathcal{R}.$$

Hence, using Lemmas 7.7 and 7.8, and putting $y \equiv fx$,

$$\begin{aligned}
(\dots) &\equiv \mathcal{E}_X(\mathbb{S}x(\Theta \cdot \mathcal{R}))(\mathcal{S}xN, P) \wedge P\{x\} \\
&\leq (\mathcal{E}_Y(\mathbb{S}(fx)\Theta))(\mathcal{R}(\mathcal{S}xN, P)) \wedge (P \cdot \Sigma^f)\{fx\} \\
&\leq (\mathcal{E}_Y(\mathbb{S}y\Theta))(\mathcal{S}y(\mathcal{R}(N, P))) \wedge \langle \mathcal{R}(N, P) \rangle \{y\} \\
&\leq \$ \mathcal{E}_Y\Theta(\mathcal{R}(N, P)) \quad \text{def } \$ \mathcal{E}_Y
\end{aligned}$$

Similarly, $\mathcal{E}_X(\mathbb{S}(iy)(\Theta \cdot \mathcal{R})) \leq \mathcal{E}_X((\mathbb{S}y\Theta) \cdot \mathcal{R}) \leq (\mathcal{E}_Y(\mathbb{S}y\Theta)) \cdot \mathcal{R}$, so

$$\begin{aligned}
(\dots) &\equiv \mathcal{E}_X(\mathbb{S}(iy)(\Theta \cdot \mathcal{R}))(\mathcal{S}(iy)N, P) \wedge P\{iy\} \\
&\leq (\mathcal{E}_Y(\mathbb{S}y\Theta))(\mathcal{R}(\mathcal{S}(iy)N, P)) \wedge (P \cdot \exists_i)\{y\} \\
&\leq (\mathcal{E}_Y(\mathbb{S}y\Theta))(\mathcal{S}y(\mathcal{R}(N, P))) \wedge \langle \mathcal{R}(N, P) \rangle \{y\} \\
&\leq \$ \mathcal{E}_Y\Theta(\mathcal{R}(N, P))
\end{aligned}$$

Finally, $\mathcal{E}_X(\mathbb{S}(jz)(\Theta \cdot \mathcal{R})) \leq \mathcal{E}_X(\Theta \cdot \mathcal{R}) \leq (\mathcal{E}_Y \Theta) \cdot \mathcal{R} \leq (\mathbb{S} \mathcal{E}_Y \Theta) \cdot \mathcal{R}$, so

$$\begin{aligned} (\cdots) &\equiv \mathcal{E}_X(\mathbb{S}(jz)(\Theta \cdot \mathcal{R}))(\mathcal{S}(jz)N, P) \wedge P\{jz\} \\ &\leq \mathbb{S} \mathcal{E}_Y \Theta(\mathcal{R}(\mathcal{S}(jz)N, P)) \\ &\leq \mathbb{S} \mathcal{E}_Y \Theta(\mathcal{R}(N, P)) \end{aligned} \quad \square$$

Proposition 7.10 If $L : \mathsf{K}X$ then $\mathcal{R}L : \mathsf{K}Y$ in all three cases.

Proof By Lemma 7.3, $\vdash \lambda \Theta. \mathcal{E}_X^{(0)}(\Theta \cdot \Omega_f) \leq \lambda \Theta. (\mathcal{E}_Y^{(0)} \Theta) \cdot \Omega_f$, and

$$\frac{n : \mathbb{N}, \lambda \Phi. \mathcal{E}_X^{(n)}(\Phi \cdot \mathcal{R}) \leq \lambda \Phi. (\mathcal{E}_Y^{(n)} \Phi) \cdot \mathcal{R}}{\lambda \Theta. \mathcal{E}_X^{(n+1)}(\Theta \cdot \mathcal{R}) \leq \lambda \Theta. (\mathcal{E}_Y^{(n+1)} \Theta) \cdot \mathcal{R}},$$

by Lemma 7.9. So by equational induction (Axiom 2.6),

$$n : \mathbb{N} \vdash \lambda \Theta. \mathcal{E}_X^{(n)}(\Theta \cdot \Omega_f) \leq \lambda \Theta. (\mathcal{E}_Y^{(n)} \Theta) \cdot \Omega_f,$$

whence $\Theta : \Sigma^{\Sigma_Y} \vdash \mathcal{E}_X^{(\infty)}(\Theta \cdot \Omega_f) \leq (\mathcal{E}_Y^{(\infty)} \Theta) \cdot \Omega_f$.

As $\Gamma \vdash L : \mathsf{K}X = \{\Omega_X \mid \mathcal{E}_X^{(\infty)}\}$, we have $\Gamma, \Phi : \Sigma^{\Omega_X} \vdash \Phi L = \mathcal{E}_X^{(\infty)} \Phi L$. Putting $\Phi = \Theta \cdot \mathcal{R}$,

$$\Theta(\mathcal{R}L) = \Phi L = (\mathcal{E}_X^{(\infty)} \Phi) L = \mathcal{E}_X^{(\infty)}(\Theta \cdot \mathcal{R}) L \leq (\mathcal{E}_Y^{(\infty)} \Theta)(\mathcal{R}L) \leq \Theta(\mathcal{R}L),$$

so these are equal and $\Gamma \vdash \mathcal{R}L : \{\Omega_Y \mid \mathcal{E}_Y^{(\infty)}\} = \mathsf{K}Y$. \square

We exploit the first case to define K as a functor.

Theorem 7.11 Let $f : X \rightarrow Y$ between overt discrete spaces and $L : \mathsf{K}X$.

- (a) Then $\mathsf{K}fL \equiv \mathcal{R}L : \mathsf{K}Y$,
- (b) the inclusion $\mathsf{K}(-) \rightarrow \Omega_{(-)}$ is natural in f ,
- (c) $\mathsf{K}f$ is a homomorphism in the sense that $\mathsf{K}f \mathsf{nil}_X = \mathsf{nil}_Y$ and $\mathsf{K}f(x :: L) = fx :: \mathsf{K}fL$;
- (d) K is a covariant functor, *i.e.* $\mathsf{Kid}_X = \mathsf{id}_{\mathsf{K}X}$ and $\mathsf{K}(g \cdot f) = \mathsf{K}g \cdot \mathsf{K}f$.
- (e) In fact, $\mathsf{K}f(L + L') = \mathsf{K}fL + \mathsf{K}fL'$, but we don't yet know that this is admissible.

Proof [a] Proposition 7.10, [b] Lemma 7.3 and Corollary 7.6, [c] Remark 7.1 and [d] Lemma 10.9. \square

The other two cases almost show that $\mathsf{K}(Y + Z) \cong \mathsf{K}Y \times \mathsf{K}Z$.

Lemma 7.12 If $(N, P) : \Omega_X$, $(N_1, P_1) : \Omega_Y$ and $(N_2, P_2) : \Omega_Z$ are modal then the following are inverse:

$$\begin{aligned} ((N_1, P_1), (N_2, P_2)) &\longmapsto (N_1 \cdot \Sigma^i \wedge N_2 \cdot \Sigma^j, P_1 \cdot \Sigma^i \vee P_2 \cdot \Sigma^j) \\ ((N \cdot \forall_i, P \cdot \exists_i), (N \cdot \forall_j, P \cdot \exists_j)) &\longleftarrow (N, P) \end{aligned}$$

If (N, P) is admissible then so are (N_1, P_1) and (N_2, P_2) . Conversely, if $(N_1, P_1) = \llbracket y \rrbracket$ and (N_2, P_2) is admissible then so is $(N, P) = iy :: K_j(N_2, P_2)$.

Proof The isomorphism uses the equations

$$\begin{aligned} \forall_i \cdot \Sigma^i \wedge \forall_j \cdot \Sigma^j &= \text{id}_{\Sigma^2 X} = \forall_i \cdot \Sigma^i \wedge \forall_j \cdot \Sigma^j \\ \Sigma^i \cdot \forall_i &= \text{id}_{\Sigma^Y} = \Sigma^i \cdot \exists_i & \Sigma^j \cdot \forall_j &= \text{id}_{\Sigma^Z} = \Sigma^j \cdot \exists_j \\ \Sigma^i \cdot \forall_j &= \top & \Sigma^j \cdot \forall_i &= \top & \Sigma^i \cdot \exists_j &= \perp & \Sigma^j \cdot \exists_i &= \perp \end{aligned}$$

Finally, $Ki\llbracket y \rrbracket + KjL = (iy) :: KjL$. \square

This is enough for us to proceed to the investigation of **ListX**. With the aid of that, we shall later be able to prove that $+$ is admissible, making KX a semilattice — indeed, the free one on X . This will also remove the restriction in the previous result.

We conclude this section by considering a *non-complemented* inclusion $i : U \hookrightarrow X$. In this case we no longer have \forall_i , but we can instead apply \exists_i to both N and P .

Lemma 7.13 Let $i : U \hookrightarrow X$ be a mono between overt discrete spaces, classified by $\phi : X \rightarrow \Sigma$. Then $Ki : KU \rightarrow KX$ is a split mono, classified by $\square\phi : KX \rightarrow \Sigma$.

$$\begin{array}{ccc} U & \xrightarrow{\quad} & \mathbf{1} \\ \downarrow i & \lrcorner & \downarrow \top \\ X & \xrightarrow{\quad \phi \quad} & \Sigma \end{array} \qquad \begin{array}{ccc} KU & \xrightarrow{\quad} & \mathbf{1} \\ \downarrow Ki & \lrcorner & \downarrow \top \\ KX & \xrightarrow{\quad \square\phi \quad} & \Sigma \end{array}$$

Proof Ki is split mono since

$$(N, P) \xrightarrow{Ki} (N \cdot \Sigma^i, P \cdot \Sigma^i) \xrightarrow{K\widehat{\exists}_i} (N \cdot \Sigma^i \cdot \exists_i, P \cdot \Sigma^i \cdot \exists_i) = (N, P).$$

The square on the right commutes, since

$$L : KU \vdash \square\phi(KiL) = [KiL]\phi = [L](\Sigma^i\phi) = [L]\top = \top.$$

If $\Gamma \vdash L : \mathbf{K}X$ with $\Box \phi L \equiv [L]\phi = \top$ then $[L](\phi \wedge \psi) = [L]\phi \wedge [L]\psi = [L]\psi$ and $\langle L \rangle \psi = [L]\phi \wedge \langle L \rangle \psi \leq \langle L \rangle (\phi \wedge \psi) \leq \langle L \rangle (\psi)$. Hence

$$\begin{aligned} \mathbf{K}i(\mathbf{K}\widehat{\exists}_i L) &= ([L] \cdot \exists_i \cdot \Sigma^i, \langle L \rangle \cdot \exists_i \cdot \Sigma^i) \\ &= (\lambda\psi. [L](\phi \wedge \psi), \lambda\psi. \langle L \rangle (\phi \wedge \psi)) \\ &= (\lambda\psi. [L]\psi, \lambda\psi. \langle L \rangle \psi) = L, \end{aligned}$$

so L belongs to the retract, which is KU . \square

Theorem 7.14 \mathbf{K} preserves monos and their inverse images.

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V & \xrightarrow{\quad} & \mathbf{1} \\ \downarrow i & \lrcorner & \downarrow j & \lrcorner & \downarrow \top \\ X & \xrightarrow{f} & Y & \xrightarrow{\phi} & \Sigma \end{array} \qquad \begin{array}{ccc} KU & \xrightarrow{\quad} & KV & \xrightarrow{\quad} & \mathbf{1} \\ \downarrow \mathbf{K}i & \lrcorner & \downarrow \mathbf{K}j & \lrcorner & \downarrow \top \\ KX & \xrightarrow{\mathbf{K}f} & KY & \xrightarrow{\Box \phi} & \Sigma \end{array}$$

Proof Since V and U are classified by ϕ and $\phi \cdot f$, we have just shown that KV and KU are classified by $\Box \phi$ and $\Box(\phi \cdot f)$, so we just need to check that

$$\Box(\phi \cdot f) = \lambda L. [L](\phi \cdot f) = \lambda L. ([L] \cdot \Sigma^f)\phi = \Box \phi \cdot \mathbf{K}f. \quad \square$$

8 Lists, heads and tails

We shall derive the recursive properties of $\mathbf{K}X$ from those of $\mathbf{List}X$. The idea of the representation is that a list or sequence of length n is a partial function $\mathbb{N} \rightharpoonup X$ with support $\{0, 1, \dots, n-1\}$, encoded as a finite set of pairs.

Definition 8.1 $\mathbf{List}X \subset \mathbf{K}(\mathbb{N} \times X)$ is the open subspace classified by

$$\begin{aligned} &\lambda L. [L](\lambda n x. [L](\lambda m y. n \neq m \vee x = y)) \\ &\wedge [L](\lambda n x. n = 0 \vee \langle L \rangle (\lambda m y. n = m + 1)). \end{aligned}$$

In more suggestive notation (which will be justified in Lemma 9.10) $\mathbf{List}X$ consists of the finite sets L of pairs such that $\forall (n, x), (m, y) \in L. n \neq m \vee x = y$ and $\forall (n, x) \in L. n = 0 \vee \exists (m, y) \in L. n = m + 1$. The first condition says that L , considered as a binary relation, is functional, and the second that this function is defined on an initial segment.

Proposition 8.2 $\mathbf{List}X$ is overt discrete, being an open subspace of an overt discrete space. \square

Definition 8.3 For $z : X$, $L : \text{List}X$, let $(z :: L) \equiv \text{Ki}\llbracket z \rrbracket + \text{K}jL$ and $\text{tail } L = \text{K}j^{-1}L$, where

$$X \xrightarrow{i = (0, \text{id})} \mathbb{N} \times X \xleftarrow{j = \text{succ} \times \text{id}} \mathbb{N} \times X.$$

Expanding the definitions from the previous section,

$$\begin{aligned} [z :: L]\phi &\equiv \phi(0, z) \wedge [L](\lambda my. \phi(m+1, y)) \\ \langle z :: L \rangle \phi &\equiv \phi(0, z) \vee \langle L \rangle (\lambda my. \phi(m+1, y)) \\ [\text{tail } L]\phi &\equiv [L](\lambda nx. n = 0 \vee \exists m. n = m+1 \wedge \phi(m, x)) \\ \langle \text{tail } L \rangle \phi &\equiv \langle L \rangle (\lambda nx. \exists m. n = m+1 \wedge \phi(m, x)). \end{aligned}$$

Lemma 8.4 $\text{nil} \in \text{List}X$, whilst if $z : X$ and $L : \text{List}X$ then $(z :: L) : \text{List}X$.

Proof As $[\text{nil}]\phi = \top$, the two conditions in Definition 8.1 are easily satisfied.

Lemma 7.12 is enough to show that $z :: L$ is in $\text{K}(\mathbb{N} \times X)$. To show that it is in $\text{List}X$, we expand first the outer $[z :: L]$ and then the inner $[z :: L]$ or $\langle z :: L \rangle$, using Corollaries 6.3 and 6.6 and some basic arithmetic.

$$\begin{aligned} [z :: L](\lambda nx. [z :: L](\lambda my. n \neq m \vee x = y)) \\ &= [z :: L](\lambda my. 0 \neq m \vee z = y) \\ &\quad \wedge [L](\lambda nx. [z :: L](\lambda my. n+1 \neq m \vee x = y)) \\ &= (0 \neq 0 \vee z = z) \wedge [L](\lambda my. 0 \neq m+1 \vee z = y) \\ &\quad \wedge [L](\lambda nx. n+1 \neq 0 \vee x = z) \\ &\quad \wedge [L](\lambda nx. [L](\lambda my. n+1 \neq m+1 \vee x = y)) \\ &= \top \wedge \top \wedge \top \wedge [L](\lambda nx. [L](\lambda my. n+1 \neq m+1 \vee x = y)) \\ &= [L](\lambda nx. [L](\lambda my. n \neq m \vee x = y)) = \top \\ [z :: L](\lambda nx. n = 0 \vee \langle z :: L \rangle (\lambda my. n = m+1)) \\ &= (0 = 0 \vee \langle z :: L \rangle (\lambda my. 0 = m+1)) \\ &\quad \wedge [L](\lambda nx. n+1 = 0 \vee \langle z :: L \rangle (\lambda my. n+1 = m+1)) \\ &= \top \wedge [L](\lambda nx. \perp \vee n+1 = 0+1 \vee \langle L \rangle (\lambda my. n+1 = m+1+1)) \\ &= [L](\lambda nx. n = 0 \vee \langle L \rangle (\lambda my. n = m+1)) = \top \quad \square \end{aligned}$$

From this we can show that **List** is a functor to the category of internal (imposed) monoids.

Theorem 8.5 Let $f : X \rightarrow Y$ between overt discrete spaces and $L : \text{List}X$.

Then

$$\text{List}(f)(L) \equiv \mathbf{K}(\mathbb{N} \times f)(L) : \text{List}(Y),$$

$\text{List}(f)$ is a homomorphism for nil and $::$, *i.e.*

$$\text{List}(f)(\text{nil}_X) = \text{nil}_Y \quad \text{and} \quad \text{List}(f)(x :: L) = fx :: \text{List}(f)(L),$$

and List is a covariant functor, *i.e.*

$$\text{List}(\text{id}_X) = \text{id}_{\text{List}(X)} \quad \text{and} \quad \text{List}(g \cdot f) = \text{List}(g) \cdot \text{List}(f).$$

Functional programmers write $\text{map}fL$ for $\text{List}fL$.

Proof We have $\mathbf{K}(\mathbb{N} \times f)(L) : \mathbf{K}(\mathbb{N} \times Y)$ by Proposition 7.11, so we have to show that it satisfies Definition 8.1, and also show that $\text{List}f$ is a homomorphism for $::$.

$$\begin{aligned} & [\mathbf{K}(\mathbb{N} \times f)(L)](\lambda ny. [\mathbf{K}(\mathbb{N} \times f)(L)](\lambda my'. n \neq m \vee y = y')) \\ &= \Sigma^2(\mathbb{N} \times f)[L](\lambda ny. \Sigma^2(\mathbb{N} \times f)[L](\lambda my'. n \neq m \vee y = y')) \quad 7.11 \\ &= [L](\lambda nx. [L](\lambda mx'. n \neq m \vee fx = fx')) \quad \text{Remark 7.1} \\ &\geq [L](\lambda nx. [L](\lambda mx'. n \neq m \vee x = x')) = \top (x = x') \leq (fx = fx') \\ &[\mathbf{K}(\mathbb{N} \times f)(L)](\lambda ny. n = 0 \vee \langle \mathbf{K}(\mathbb{N} \times f)(L) \rangle(\lambda my'. n = m + 1)) \\ &= [L](\lambda nx. n = 0 \vee \langle L \rangle(\lambda mx'. n = m + 1)) = \top \quad \text{Proposition 7.11} \\ &\mathbf{K}(\mathbb{N} \times f)(x :: L) \\ &= \mathbf{K}(\mathbb{N} \times f)((0, x) :: \mathbf{K}(\text{succ} \times X)L) \quad \text{Definition 8.3} \\ &= (0, fx) :: \mathbf{K}(\mathbb{N} \times f) \cdot \mathbf{K}(\text{succ} \times X)L \quad \text{Proposition 7.11} \\ &= (0, fx) :: \mathbf{K}(\text{succ} \times Y) \cdot \mathbf{K}(\mathbb{N} \times f)L \quad \text{def } (\text{succ} \times f) \\ &= \mathbf{K}(\text{succ} \times Y)(fx :: \mathbf{K}(\mathbb{N} \times f)L) \quad \text{Proposition 7.11} \end{aligned}$$

The other parts follow directly from Proposition 7.11. \square

Next we show that every list is either empty or has a unique head and tail.

Lemma 8.6 For $L : \text{List}X$ and $n : \mathbb{N}$,

$$(\exists x. (n + 1, x) \in L) \leq (\exists y. (n, y) \in L) \leq (\exists z. (0, z) \in L).$$

Proof Write $\alpha_n \equiv \langle L \rangle(\lambda mx. n = m) \equiv (\exists x. (n, x) \in L)$. The second part of Definition 8.1 for $L : \text{List}X$ says

$$[L](\lambda nx. n = 0 \vee \langle L \rangle(\lambda my. n = m + 1)),$$

which, together with the hypothesis $\alpha_{n+1} = \top$, gives

$$(n + 1 = 0 \vee \langle L \rangle (\lambda my. n + 1 = m + 1)) = \top,$$

which is equivalent to $\alpha_n = \top$. Hence by Axiom 2.4 we have $\alpha_{n+1} \leq \alpha_n$ and

$$L : \text{List}X, n : \mathbb{N}, \alpha_n \leq \alpha_0 \vdash \alpha_{n+1} \leq \alpha_0,$$

which is the induction step for proving $L : \text{List}X, n : \mathbb{N} \vdash \alpha_n \leq \alpha_0$. \square

Lemma 8.7 If $L : \text{List}X$ then $L = \text{nil} \vee \exists!z. (0, z) \in L$.

Proof From Proposition 3.12, either $L \sim \text{nil}$ or $\exists nx. (n, x) \in L$. In the latter case the previous result applies, giving $\exists z. (0, z) \in L$, which is unique by the first part of Definition 8.1. \square

Lemma 8.8 Let $L : \text{List}X$ and $x : X$. Then $\text{tail } L : \mathbb{K}(\mathbb{N} \times X)$, $\text{tail } (x :: L) = L$, and if $(0, z) \in L : \text{List}X$ then $L = z :: \text{tail } L$.

Proof These are corollaries of Lemma 7.12.

$$\begin{aligned} \text{tail } (z :: L) &= \mathbb{K}j^{-1}(\mathbb{K}i\{z\} + \mathbb{K}jL) \\ &= (\mathbb{K}j^{-1} \cdot \mathbb{K}i)\{z\} + (\mathbb{K}j^{-1} \cdot \mathbb{K}jL) = \text{nil} + L \\ z :: \text{tail } L &= \mathbb{K}i\{z\} + \mathbb{K}j(\text{tail } L) \\ &= \mathbb{K}i(\mathbb{K}i^{-1}L) + \mathbb{K}j(\mathbb{K}j^{-1}L) = L \quad \{z\} = \mathbb{K}i^{-1}L \quad \square \end{aligned}$$

Lemma 8.9 If $L : \text{List}X$ then $\text{tail } L : \text{List}X$.

Proof

$$\begin{aligned} &[\text{tail } L](\lambda nx. [\text{tail } L](\lambda my. n \neq m \vee x = y)) \\ &= [L](\lambda nx. n = 0 \vee (\exists n'. n = n' + 1 \wedge [L](\lambda my. m = 0 \\ &\quad \vee \exists m'. m = m' + 1 \wedge (n' \neq m' \vee x = y)))) \\ &\geq [L](\lambda nx. n = 0 \vee (n \neq 0 \wedge [L](\lambda my. n \neq m \vee x = y))) \\ &\geq [L](\lambda nx. [L](\lambda my. n \neq m \vee x = y)) \geq \top \\ &[\text{tail } L](\lambda nx. n = 0 \vee \langle \text{tail } L \rangle (\lambda my. n = m + 1)) \\ &= [L](\lambda nx. n = 0 \vee \exists n'. n = n' + 1 \wedge (n' = 0 \\ &\quad \vee \langle L \rangle (\lambda my. \exists m'. m = m' + 1 \wedge n' = m' + 1))) \\ &\geq [L](\lambda nx. n = 0 \vee n \neq 0 \wedge (n = 1 \vee \langle L \rangle (\lambda my. n = m + 1 \neq 1))) \\ &\geq [L](\lambda nx. n = 0 \vee n \neq 0 \wedge \langle L \rangle (\lambda my. n = m + 1)) \\ &\geq [L](\lambda nx. \langle L \rangle (\lambda my. n = m + 1)) \geq \top \quad \square \end{aligned}$$

Proposition 8.10 $\text{List}X \cong \mathbf{1} + X \times \text{List}X$, cf. Proposition 5.14 for $\mathbf{K}X$. In particular,

(a) $L : \text{List}X \vdash (L =_{\text{List}X} \text{nil} \vee \exists x L'. L =_{\text{List}X} x :: L') = \top$, and

(b) $x, y : X, L, L' : \text{List}X \vdash$

$$(x :: L =_{\text{List}X} y :: L') = (x =_X y) \wedge (L =_{\text{List}X} L').$$

□

9 Recursion over lists

Lemma 9.1 Define $\downarrow : \mathbb{N} \rightarrow \text{List}(\mathbf{1})$ by recursion,

$$\downarrow 0 = \text{nil} \quad \text{and} \quad \downarrow(n+1) = \star :: (\downarrow n),$$

where $\star : \mathbf{1}$, so \downarrow is a homomorphism. Then $((k, \star) \in \downarrow n) = (k < n)$,

$$[\downarrow n] = \lambda\phi. \forall k < n. \phi(k, \star) \quad \text{and} \quad \langle \downarrow n \rangle = \lambda\phi. \exists k < n. \phi(k, \star).$$

Proof The recursive definition says that $[\downarrow 0]\phi = \top$ and

$$[\downarrow(n+1)]\phi = \phi(0, \star) \wedge [\downarrow n](\lambda ky. \phi(k+1, y)),$$

but $\lambda\phi. \forall k < n. \phi(k, \star)$ satisfies the same equations (Remark 3.2), so they are equal by the universal property of \mathbb{N} . Similarly for $\langle \downarrow n \rangle$ with \perp , \vee and \exists in place of \top , \wedge and \forall . □

Remark 9.2 This means that

$$\downarrow 1 = \{(0, \star)\}, \quad \downarrow 2 = \{(0, \star), (1, \star)\}, \quad \downarrow 3 = \{(0, \star), (1, \star), (2, \star)\}, \dots$$

but we shall ignore the \star from now on, and regard $\text{List}(\mathbf{1}) \subset \mathbf{K}(\mathbb{N})$.

Lemma 9.3 There is a unique map $|-| : \text{List}(\mathbf{1}) \rightarrow \mathbb{N}$ such that

$$(n < |L|) = (n \in L) \equiv \langle L \rangle(\lambda x. x = n)$$

and

$$(|L| \leq n) = (n \notin L) \equiv [L](\lambda x. x \neq n).$$

Proof We must define $|L|$ in terms of the presence or absence of numbers in the list, as it would be begging the question to use recursion on $\text{List}(\mathbf{1})$.

The two properties $n \in L$ and $n \notin L$ (as defined by \equiv above) are complementary by the modal laws, in particular Corollary 6.8.

$$\top = [L](\lambda x. x \neq n \vee x = n)$$

$$\begin{aligned}
&\leq [L](\lambda x. x \neq n) \vee \langle L \rangle(\lambda x. x = n) \equiv (n \notin L) \vee (n \in L) \\
\perp &= \langle L \rangle(\lambda x. x \neq n \wedge x = n) \\
&\geq [L](\lambda x. x \neq n) \wedge \langle L \rangle(\lambda x. x = n) \equiv (n \notin L) \wedge (n \in L).
\end{aligned}$$

Now $\phi \equiv \lambda n. (x < n)$ satisfies the premise of Axiom 1.9, so

$$\begin{aligned}
\top &= [L](\lambda x. \exists n. x < n) = \exists n. [L](\lambda x. x < n) \\
&\leq \exists n. [L](\lambda x. x \neq n) = \exists n. n \notin L.
\end{aligned}$$

Hence we may use general recursion or sobriety [A, Lemma 9.11] to define $|L| \equiv \mu n. n \notin L$, which therefore satisfies $(n < |L|) \leq (n \in L)$ and $(|L| \leq n) \geq (n \notin L)$. These are actually equalities, as Lemma 8.6 said that

$$(n + 1 \in L) \leq (n \in L), \quad \text{and so} \quad (n \notin L) \leq (n + 1 \notin L),$$

and indeed $(m \geq n \notin L) \leq (m \notin L)$ by induction. \square

Proposition 9.4 $|-|$ and \downarrow make $\mathbf{List}(\mathbf{1}) \cong \mathbb{N}$.

Proof We show that \downarrow and $|-|$ are inverse. Using Lemmas 9.1, 9.3 and 6.2,

$$\begin{aligned}
\langle \downarrow |L| \rangle \phi &= \exists x < |L|. \phi x = \exists x. x < |L| \wedge \phi x \\
&= \exists x. \langle L \rangle(\lambda y. x = y) \wedge \phi x = \langle L \rangle \phi,
\end{aligned}$$

so $\downarrow |L| = L$ by Proposition 3.10. On the other hand,

$$(|\downarrow m| > n) = \langle \downarrow m \rangle(\lambda x. x = n) = (\exists x < m. x = n) = (n < m)$$

by Lemmas 9.3 and 9.1, so $|\downarrow m| = m$. \square

Corollary 9.5 For any overt discrete object X there is a map $|-| : \mathbf{List}X \rightarrow \mathbb{N}$ such that

$$|L| = 0 \iff L = \mathbf{nil}, \quad \text{and} \quad |z :: L| = |L| + 1.$$

The number $|L|$ is called the **length** of the list.

Proof Define $|L| \equiv |\mathbf{List}(!_X)L|$, which is the composite of one homomorphism with the inverse of another. By Lemma 8.7, $(L = \mathbf{nil})$ is decidable. If $L \neq \mathbf{nil}$ then $L = x :: L'$ for some x and L' , so $|L| = |L'| + 1 \neq 0$. \square

Now we can start to define recursion over lists, in a way that will be very familiar to functional programmers. The first result depends on Proposition 8.10 and establishes existence, and the second uses Corollary 9.5 and gives uniqueness.

Lemma 9.6 Let $A = \Sigma^U$ for some object U , equipped with an **action** of X :

$$\Gamma \vdash \zeta : A \quad \text{and} \quad \Gamma, x : X, \alpha : A \vdash \sigma(x, \alpha) : A.$$

Then the term $\Gamma \vdash \epsilon : A^{\text{List}X}$ is a homomorphism in the sense that

$$\epsilon \text{ nil} = \zeta \quad \text{and} \quad \epsilon(x :: L) = \sigma(x, \epsilon L).$$

iff it satisfies the fixed point equation

$$\epsilon = \lambda L. (L = \text{nil} \wedge \zeta) \vee \exists x : X. \exists L' : \text{List}X. (L = x :: L') \wedge \sigma(x, \epsilon L').$$

Proof If ϵ is a homomorphism then by Proposition 8.10(a),

$$\begin{aligned} \epsilon L &= (L = \text{nil} \wedge \epsilon L) \vee \exists x L'. (L = x :: L' \wedge \epsilon L) \\ &= (L = \text{nil} \wedge \zeta) \vee \exists x L'. L = x :: L' \wedge \sigma(x, \epsilon L') \end{aligned}$$

Conversely, if it is a fixed point then by Proposition 8.10(b),

$$\begin{aligned} \epsilon \text{ nil} &= (\text{nil} = \text{nil} \wedge \zeta) \vee \exists x L'. \perp \wedge \sigma(x, \epsilon L') = \zeta \\ \epsilon(y :: L) &= (y :: L = \text{nil} \wedge \zeta) \vee \exists x L'. (y :: L = x :: L') \wedge \sigma(x, \epsilon L') \\ &= \perp \vee \exists x L'. (x = y) \wedge (L = L') \wedge \sigma(x, \epsilon L') \\ &= \sigma(y, \epsilon L). \end{aligned} \quad \square$$

The simplicity of the last step is one reason why we have done this part of the argument for lists: the corresponding argument for finite subsets would be much more difficult. The other reason is that lists have a well defined length, which is the key to proving uniqueness, and more generally **equational induction for lists**.

Proposition 9.7 Let $\Gamma, L : \text{List}X \vdash \alpha L, \beta L : A \equiv \Sigma^U$. Then, cf. Axiom 2.6,

$$\frac{\Gamma \vdash \alpha \text{ nil} = \beta \text{ nil} \quad \Gamma, L : \text{List}X, x : X, \alpha L = \beta L \vdash \alpha(x :: L) = \beta(x :: L)}{\Gamma, L : \text{List}X \vdash \alpha L = \beta L}$$

Proof Consider $\alpha_n \equiv \lambda L. |L| \leq n \wedge \alpha L$ and $\beta_n \equiv \lambda L. |L| \leq n \wedge \beta L$, both of type $\Sigma^{A \times \text{List}X}$. Then $\alpha_0 = \beta_0$ because $(L = \text{nil}) \iff (|L| = 0)$ by Corollary 9.5.

Suppose that $\alpha_n = \beta_n$. Then if $L' : \text{List}X$ with $|L'| \leq n$, we have $\alpha L' = \alpha_n L' = \beta_n L' = \beta L'$ and so $\alpha(x :: L') = \beta(x :: L')$ by hypothesis. Hence, using Proposition 8.10(a) and Corollary 9.5,

$$\alpha_{n+1} = \lambda L. (L = \text{nil} \vee \exists x. \exists L'. (L = x :: L')) \wedge (|L| \leq n + 1) \wedge \alpha L$$

$$\begin{aligned}
&= \lambda L. (L = \text{nil} \wedge \alpha \text{nil}) \\
&\quad \vee \exists x. \exists L'. (L = x :: L') \wedge (|L'| \leq n) \wedge \alpha(x :: L') \\
&= (\text{the same for } \beta) = \beta_{n+1}.
\end{aligned}$$

Thus $\alpha_n = \beta_n$ by equational induction for \mathbb{N} (Axiom 2.6), and

$$\alpha L = (|L| \leq |L|) \wedge \alpha L = \alpha_{|L|} L = \beta_{|L|} L = \beta L. \quad \square$$

Theorem 9.8 $\text{List}X$ is the free (imposed, $::$) monoid on X . It also obeys equational induction at all types.

Proof Let M be any type with an action $\Gamma \vdash z : M, r : X \times M \rightarrow M$. We have already shown that some homomorphism $\text{List}X \rightarrow M$ exists and is unique in the case where $M = \Sigma^U$, so we consider $A \equiv \Sigma^{\Sigma^M}$, $\zeta = \lambda\psi. \psi x : A$ and $\sigma : X \times A \rightarrow A$ by $\sigma(x, F) = \lambda\psi. F(\lambda m. (\psi \cdot r)(x, m))$.

Then there is a unique map $\epsilon : \text{List}X \rightarrow A$ such that

$$\begin{aligned}
\epsilon \text{nil} &= \zeta = \lambda\psi. \psi z \\
\epsilon(x :: L) &= \sigma(x, \epsilon L) = \lambda\psi. \epsilon L(\lambda m. (\psi \cdot r)(x, m)).
\end{aligned}$$

Now we use Proposition 9.7 to show that ϵL is prime [A, §4]. Clearly ϵnil is prime. Suppose that ϵL is prime, so (with $\mathcal{F} : \Sigma^3 M$)

$$\lambda\mathcal{F}. \mathcal{F}(\epsilon L) = \lambda\mathcal{F}. \epsilon L(\lambda m. \mathcal{F}(\lambda\psi. \phi m)).$$

We shall show that $\epsilon(x :: L)$ is also prime (with respect to $\mathcal{G} : \Sigma^3 M$), using

$$\mathcal{F} \equiv \lambda F. \mathcal{G}(\lambda\psi. F(\lambda m. (\psi \cdot r)(x, m)))$$

in the primality equation for ϵL , so

$$\begin{aligned}
\mathcal{G}(\epsilon(x :: L)) &= \mathcal{G}(\lambda\psi. \epsilon L(\lambda m. (\psi \cdot r)(x, m))) && \text{def } \epsilon(x :: L) \\
&\equiv \mathcal{F}(\epsilon L) && \text{def } \mathcal{F} \\
&= \epsilon L(\lambda m. \mathcal{F}(\lambda\psi. \psi m)) && \epsilon L \text{ prime (hypothesis)} \\
&= \epsilon L(\lambda m. \mathcal{G}(\lambda\psi. (\psi \cdot r)(x, m))) && \text{def } \mathcal{F} \\
&= (\lambda\psi'. \epsilon L(\lambda m. (\psi' \cdot r)(x, m))) (\lambda m'. \mathcal{G}(\lambda\psi. \psi m')) (\lambda\beta)^{-1} \\
&= \epsilon(x :: L) (\lambda m'. \mathcal{G}(\lambda\psi. \psi m')) && \text{def } \epsilon(x :: L)
\end{aligned}$$

Hence $\epsilon L = \lambda\psi. \psi(eL)$ for some unique map $e : \text{List}X \rightarrow M$, which also satisfies $e \text{nil} = z$ and $e(x :: L) = r(x, eL)$. \square

We can now define all sorts of operations on lists in the usual way. We shall need the following in particular.

Lemma 9.9 $n < |L|$ iff $\exists!x:X. (n, x) \in L$, so we may define $L@n$ by description [A, Section 9]. \square

Lemma 9.10 $[L]\phi = \forall n < |L|. \phi(n, L@n)$ and $\langle L \rangle \phi = \exists n < |L|. \phi(n, L@n)$.

Proof They satisfy the same recursion equations (cf. Remark 3.2). \square

Definition 9.11 We define **concatenation** $(++)$ on $\text{List}X$ in the usual way by recursion:

$$\text{nil} ++ L_2 = L_2 \quad \text{and} \quad (x :: L_1) ++ L_2 = x :: (L_1 ++ L_2).$$

Beware that $++$ on $\text{List}X$ is not the same as $+$ on $\mathbf{K}(\mathbb{N} \times X)$ or $\Omega_{\mathbb{N} \times X}$.

Proposition 9.12 Using list induction we prove in the usual way that

- (a) $++$ is associative with unit nil ;
- (b) for any $f : X \rightarrow Y$, $\text{List}f$ is a homomorphism for nil and $++$; and
- (c) $\text{List}X$ is the free monoid on X with respect to $++$. \square

Lemma 9.13 $x = L@m$ for some m iff $L = L' ++ \{x\} ++ L''$ for some L', L'' , where $L' = \text{take}(L, m)$ and $L'' = \text{drop}(L, m+1)$ in functional programming notation. \square

Lemma 9.14 List preserves equalisers.

Proof Let $X \xrightarrow{i} Y \xrightleftharpoons[g]{f} Z$ be an equaliser of overt discrete spaces. (This exists, given f and g , because Z is discrete.) We show, using equational list induction on $\Gamma \vdash L : \text{List}Y$, that if $\Gamma \vdash \text{List}fL = \text{List}gL : \text{List}Z$ then $\Gamma \vdash \exists!L':\text{List}X. L = \text{List}iL'$. For the base case, $\text{nil}_Y = \text{List}i \text{nil}_X$. For the induction step,

$$fy :: \text{List}fL \equiv \text{List}f(y :: L) = \text{List}g(y :: L) \equiv gy :: \text{List}gL$$

iff $fy = gy \wedge \text{List}fL = \text{List}gL$. Hence if $L = \text{List}iL'$ (by the induction hypothesis) and $y = ix$ then $y :: L = \text{List}i(x :: L')$. \square

Lemma 9.15 List preserves pullback over $\mathbf{1}$, *i.e.*

$$\begin{array}{ccc}
 X + Y & \xrightarrow{\quad} & Y \\
 \downarrow \lrcorner & & \downarrow \\
 X & \xrightarrow{\quad} & \mathbf{1}
 \end{array}
 \quad
 \begin{array}{ccc}
 \text{List}(X \times Y) & \xrightarrow{\quad} & Y \\
 \downarrow \lrcorner & & \downarrow | - | \\
 \text{List}X & \xrightarrow{| - |} & \mathbb{N}
 \end{array}$$

Proof Define $\text{zip} : \text{List}X \times \text{List}Y \rightarrow \text{List}(X \times Y)$ by

$$\begin{aligned}
 \text{zip}(x :: L_1, y :: L_2) &= (x, y) :: \text{zip}(L_1, L_2) \\
 \text{zip}(L_1, \text{nil}) &= \text{nil} = \text{zip}(\text{nil}, L_2).
 \end{aligned}$$

Now if $\Gamma \vdash L_1 : \text{List}X$ and $\Gamma \vdash L_2 : \text{List}Y$ make a commutative square, *i.e.* $|L_1| = |L_2|$, then $L_1 = \text{List}\pi_0(\text{zip}(L_1, L_2))$ and $L_2 = \text{List}\pi_1(\text{zip}(L_1, L_2))$, and $\Gamma \vdash \text{zip}(L_1, L_2) : \text{List}(X \times Y)$ is the unique thing that does this. \square

Theorem 9.16 List preserves all finite connected limits.

Proof They may be obtained from equalisers and pullbacks like this. \square

10 The free semilattice

Now that we have the free monoid, and [C, Section 11] showed how to construct (stable effective) quotients of overt discrete objects by open equivalence relations, the free semilattice exists. We want to show that this free semilattice is in fact \mathbf{KX} . One way to do this would be to follow the motivation in Section 3, which we have now made legitimate. Instead, we shall identify the quotient map $\text{List}X \rightarrow \mathbf{KX}$ for the semilattice laws directly.

Recall from Theorem 7.11 that $\mathbf{K}\pi_1$ is a homomorphism in the sense that $\mathbf{K}\pi_1 \text{nil} = \text{nil}$ and $\mathbf{K}\pi_1(x :: L) = x :: \mathbf{K}\pi_1 L$.

Lemma 10.1 $\mathbf{K}\pi_1$ is a homomorphism: $\mathbf{K}\pi_1(L_1 \# L_2) = \mathbf{K}\pi_1 L_1 + \mathbf{K}\pi_1 L_2$. \square

Proposition 10.2 $\mathbf{K}\pi_1 : \text{List}X \rightarrow \mathbf{KX}$ is an open surjection.

Proof It is an open map since both $\text{List}X$ and \mathbf{KX} are overt discrete [C, Lemma 10.2]. For it to be an open surjection [C, Definition 10.4] we have to show that $\Theta \equiv \exists_{\mathbf{K}\pi_1} \top = \top : \Sigma^{\mathbf{KX}}$, where

$$S : \mathbf{KX} \vdash \Theta S \equiv \exists_{\mathbf{K}\pi_1} \top S = \exists L : \text{List}X. (\mathbf{K}\pi_1 L = S).$$

Since $\mathsf{K}\pi_1 \text{ nil} = \text{nil}$, we have $\Theta \text{ nil} = \top$. Then

$$\begin{aligned} \Theta S &\equiv \exists L. (\mathsf{K}\pi_1 L = S) \\ &\leq \exists L. (\mathsf{K}\pi_1(x :: L) = x :: S) && \text{above} \\ &\leq \exists L'. (\mathsf{K}\pi_1 L' = x :: S) && \text{where } L' = x :: L \\ &= \Theta(x :: S) = \mathbb{S}x\Theta S \end{aligned}$$

so $\Theta \leq \mathbb{S}x\Theta$. Hence $\Theta = \lambda S. \top$ by Corollary 4.12. \square

Lemma 10.3 $\mathsf{K}\pi_1$ is a natural transformation from **List** to **K**.

Proof Both composites in the square are of the form $(N, P) \mapsto (N \cdot H, P \cdot H)$, where $H = \Sigma^{\pi_1} \cdot \Sigma^f = \Sigma^{\text{id} \times f} \cdot \Sigma^{\pi_1}$. \square

Notation 10.4 Write $L_1, L_2 : \text{List}X \vdash L_1 \approx L_2 : \Sigma$ for the open congruence generated by the semilattice laws. Explicitly, this is of the form $\exists \mathcal{L} : \text{List}(\text{List}X). (\dots)$, where (\dots) says that the list \mathcal{L} of lists begins L_1 , ends L_2 and its successive members are related by one of the semilattice laws. Being a congruence also means that if $L_1 \approx L_2$ and $L_3 \approx L_4$ then $L_1 \# L_3 \approx L_2 \# L_4$.

Lemma 10.5 $\mathsf{K}\pi_1 : \text{List}X \rightarrow \mathsf{K}X$ coequalises these laws, *i.e.* if $L_1 \approx L_2$ then $\mathsf{K}\pi_1 L_1 = \mathsf{K}\pi_1 L_2$.

Proof This is an equational induction over the list \mathcal{L} , in which the induction step considers a single instance of a semilattice law $L_1 \approx L_2$ in which two elements are either interchanged or coalesced. But $\mathsf{K}\pi_1$ is a homomorphism for $+$, and these laws hold in $\mathsf{K}X$. \square

Lemma 10.6 $\mathsf{K}\pi_1 L_1 \subset \mathsf{K}\pi_1 L_2$ iff $(\forall n < |L_1|. \exists m < |L_2|. L_1 @ n = L_2 @ m)$.

Proof

$$\begin{aligned} \mathsf{K}\pi_1 L_1 \subset \mathsf{K}\pi_1 L_2 &= [\mathsf{K}\pi_1 L] (\lambda x. \langle \mathsf{K}\pi_1 L_2 \rangle (\lambda y. x = y)) && \text{Notation 3.3} \\ &= \Sigma^2 \pi_1 [L] (\lambda x. \Sigma^2 \pi_1 \langle L_2 \rangle (\lambda y. x = y)) && \text{Remark 7.1} \\ &= [L] (\lambda n x. \langle L_2 \rangle (\lambda m y. x = y)) \\ &= (\forall n < |L_1|. \exists m < |L_2|. L_1 @ n = L_2 @ m) L. && \text{9.10 } \square \end{aligned}$$

Lemma 10.7 If $\mathsf{K}\pi_1 L_1 \subset \mathsf{K}\pi_1 L_2$ then $L_1 \# L_2 \approx L_2$.

Proof By equational list induction on L_1 . For the base case, $\text{nil} \# L_2 = L_2$. Suppose that $x \in \mathsf{K}\pi_1 L$. Then $x = L @ m$ for some $m < |L|$, so $L = L \# \{x\} \# L$ for some lists L and L by Lemma 9.13, whence

$$(x \in \mathsf{K}\pi_1 L) \leq (\{x\} \# L \approx L) : \Sigma.$$

Hence for the induction step,

$$\begin{aligned}
 (\mathsf{K}\pi_1(x :: L_1) \subset \mathsf{K}\pi_1 L_2) &= (x \in \mathsf{K}\pi_1 L_2) \wedge (\mathsf{K}\pi_1 L_1 \subset \mathsf{K}\pi_1 L_2) \\
 &\leq (x \in \mathsf{K}\pi_1(L_1 + L_2)) \wedge (L_1 \# L_2 \approx L_2) \\
 &\leq (x :: (L_1 + L_2) \approx (L_1 + L_2)) \wedge (L_1 \# L_2 \approx L_2) \\
 &\leq ((x :: L_1) \# L_2 \approx L_2) \quad \square
 \end{aligned}$$

Corollary 10.8 $\mathsf{K}\pi_1 L_1 = \mathsf{K}\pi_1 L_2$ iff $L_1 \approx L_1 + L_2 \approx L_2$, and so $\mathsf{K}X = \text{List}X/\approx$. This coequaliser is valid in \mathcal{S} as well as in \mathcal{E} [C, Lemma 10.8]. \square

Lemma 10.9 $+: \mathsf{K}X \times \mathsf{K}X \rightarrow \mathsf{K}X$ is well defined and is preserved by $\mathsf{K}f$.

$$\begin{array}{ccccc}
 \text{List}X \times \text{List}X & \xrightarrow{\mathsf{K}\pi_1 \times \mathsf{K}\pi_1} & \mathsf{K}X \times \mathsf{K}X & \xrightarrow{\quad} & \Omega_X \times \Omega_X \\
 \downarrow \# & & \vdots & & \downarrow + \\
 \text{List}X & \xrightarrow{\mathsf{K}\pi_1} & \mathsf{K}X & \xrightarrow{\quad} & \Omega_X
 \end{array}$$

Proof The rectangle commutes by Lemma 10.1, the top left map is a surjection by Proposition 10.2 and the bottom right is a Σ -split mono, so there is a unique fill-in. The map $f : X \rightarrow Y$ turns the diagram into a commutative cuboid, since $\mathsf{K} \mapsto \Omega$. $\mathsf{K}\pi_1$, $+$ and $\#$ are natural by Theorem 7.11(b), Lemma 10.3, Lemma 7.4 and Proposition 9.12(b). \square

Theorem 10.10 $\mathsf{K}X$ is the free semilattice (in the sense of $+$) on X in \mathcal{S} .

$$\begin{array}{ccccc}
 & & \text{List}X & \xrightarrow{\mathsf{K}\pi_1} & \mathsf{K}X & \xrightarrow{\quad} & \Omega \\
 & & \uparrow & & \vdots & & \\
 & & \mathbb{I}(0, -)\mathbb{I} & \searrow \epsilon & \downarrow \delta & & \\
 & & X & \xrightarrow{\quad} & M & &
 \end{array}$$

Proof If M is an (imposed) semilattice then it is in particular a monoid, so by Proposition 9.12(c) there is a unique homomorphism (for nil , $:::$ and $\#$) $\epsilon : \text{List}X \rightarrow M$. But as M also obeys the semilattice laws, ϵ factors through the coequaliser (Corollary 10.8), giving the required mediator $\delta : \mathsf{K}X \rightarrow M$. This is a homomorphism (for nil and $+$) because $\mathsf{K}\pi_1$ is, and is surjective. If $\delta' : \mathsf{K}X \rightarrow M$ is another homomorphism for nil and $+$ then $\delta' \cdot \mathsf{K}\pi_1 :$

$\text{List}X \rightarrow M$ is a homomorphism for nil and \dashv , so $\delta' \cdot K\pi_1 = \delta \cdot K\pi_1$, and $\delta' = \delta$ because $K\pi_1$ is surjective. \square

Proposition 10.11 Let $\Gamma, L : KX \vdash \alpha L, \beta L : A \equiv \Sigma^U$. Then, cf. Axiom 2.6 and Proposition 9.7,

$$\frac{\Gamma \vdash \alpha \text{nil} = \beta \text{nil} \quad \Gamma, L : KX, x : X, \alpha L = \beta L \vdash \alpha(x :: L) = \beta(x :: L)}{\Gamma, L : KX \vdash \alpha L = \beta L}$$

Proof Consider $\bar{\alpha} \equiv \alpha \cdot K\pi_1, \bar{\beta} \equiv \beta \cdot K\pi_1 : \text{List}X \rightarrow KX \Rightarrow A$. These satisfy Proposition 9.7, so $\bar{\alpha} = \bar{\beta}$, whence $\alpha = \beta$ since $K\pi_1$ is surjective. \square

Corollary 10.12 Any Σ -split subspace $U \subset KX$ that contains nil and is closed under $::$ is $U = KX$.

Proof U can be expressed as the equaliser of some $\alpha, \beta : KX \Rightarrow A$ [B, Proposition 4.14]. \square

11 Overt compact subspaces

After this lengthy manipulation in the logical calculus of ASD, you may be left wondering what “admissible” or “modal” terms $L : KX$ ever had to do with “finite subsets” or “compact open subspaces” of X . In fact, the topological results follow from the *modal* laws in Definition 3.4, and it is these that we assume of $L : \Omega$, rather than admissibility ($L : KX$), which we deduce on the basis of Axiom 1.10. In view of the lack of a theory of parametric types for ASD, we first consider a *global* (non-parametric) modal element $\vdash L : \Omega$.

Proposition 11.1 The open subspace $i : U \subset X$ classified by π is overt (and discrete), with existential quantifier $\exists_U = \langle L \rangle \cdot \exists_i$ and modal operator $\langle L \rangle = \exists_U \cdot \Sigma^i$. This justifies the notation $\langle L \rangle \phi \equiv \exists x \in L. \phi x$, cf. the last version of Lemma 6.2.

Proof The space X is overt by hypothesis, and U is an open subspace of it classified by π , so U is itself overt, with $\exists_U = \exists_X \cdot \exists_i$. What we have to show, therefore, is that

$$\langle L \rangle \cdot \exists_i = \exists_X \cdot \exists_i \quad \text{and} \quad \langle L \rangle \cdot \exists_i \cdot \Sigma^i = \langle L \rangle.$$

As usual, we regard an open predicate θ on the open subspace $U \subset X$ (which is classified by π) as a predicate on X itself, with $\theta \leq \pi$. This means that we

represent \exists_i by id and Σ^i by $\pi \wedge (-)$. Using the last of the eight modal laws, the first equation is then

$$\langle L \rangle (\exists_i \theta) = \langle L \rangle \theta = \exists x. \pi x \wedge \theta x = \exists x. \theta x = \exists_X (\exists_i \theta)$$

and the second, for $\phi : \Sigma^X$, is $\langle L \rangle (\exists_i \cdot \Sigma^i \phi) = \langle L \rangle (\pi \wedge \phi) = \langle L \rangle \phi$. \square

Proposition 11.2 The subspace $i : U \subset X$ is also compact, with universal quantifier $\forall_U = [L] \cdot \exists_i$ and modal operator $[L] = \forall_U \cdot \Sigma^i$. This justifies the notation $[L]\phi \equiv \forall x \in L. \phi x$.

Proof We have to show that $\Sigma^! \dashv [L] \cdot \exists_i$ and $[L] \cdot \exists_i \cdot \Sigma^i = [L]$. Again we represent \exists_i by id and Σ^i by $\pi \wedge (-)$. Let $\sigma : \Sigma$ and $u : U$, *i.e.* $u : X$ with $\pi u = \top$.

Suppose $\sigma \leq [L]\psi$. Then $\sigma \leq [L]\psi \wedge \pi u \leq \psi u$ by Remark 3.6, so $(\lambda u : U. \sigma) \leq \psi$.

Conversely, suppose $\lambda u : U. \sigma \leq \psi$, so $x : X \vdash \sigma \wedge \pi x \leq \psi x$. Then

$$\begin{aligned} \sigma &= \sigma \wedge [L]\pi && \text{7th modal law} \\ &\leq [L](\lambda x. \sigma \wedge \pi x) && \text{Euclid} \\ &\leq [L]\psi && \text{hypothesis} \end{aligned}$$

For the modal operator, with $\phi : \Sigma^X$, $[L](\exists_i \cdot \Sigma^i \phi) = [L]\phi$. \square

The subspace U need not be Hausdorff, *i.e.* have decidable equality, as we haven't assumed this of X itself.

Proposition 11.3 Conversely, let $U \subset X$ be compact open. Then $\vdash (N, P) : \Omega$ is modal, where $P \equiv \exists_U \cdot \Sigma^i$ and $N \equiv \forall_U \cdot \Sigma^i$.

Proof As for Proposition 11.10 below without γ . \square

Once again we have a result for *modal* $L : \Omega$ when we really want one for *admissible* $L : \mathbf{K}X$, but now at last we are able to state, and so invoke, the Scott continuity Axiom 1.10.

Theorem 11.4 $\vdash L : \Omega_X$ is admissible iff it is modal, and such terms correspond bijectively to compact open subspaces $U \subset X$. Moreover, a subspace $U \subset X$ is compact open iff it is listable.

Proof Without loss of generality $U = X$. Applying Axiom 1.10 to $F \equiv \forall_X$, $\alpha_\ell = \top$ and $\phi_\ell = \pi_\ell$, where X is a compact overt discrete space, we have

$$\vdash \top = \forall_X \top = \exists \ell : \mathbf{K}X. \forall x : X. x \in \ell = \exists \ell : \mathbf{K}X. (\forall x : X. x \in \ell).$$

This means that $(\forall_X, \exists_X) \subset \ell$, where ℓ is admissible and (\forall_X, \exists_X) is modal, whilst $\ell \subset (\forall_X, \exists_X)$ since (\forall_X, \exists_X) is the greatest modal pair (Proposition 3.8). Hence $(\forall_X, \exists_X) \sim \ell$, which means that (\forall_X, \exists_X) is admissible (and $\exists \ell$ above is unique). Notice that the two universal quantifiers in

$$\exists \ell: \mathbf{K}X. (\forall y \in \ell. \exists u: U. x = iu) \wedge (\forall u: U. \exists x \in \ell. x = iu) : \Sigma$$

are legitimate because both U and ℓ define compact spaces.

In the last part, \Leftarrow follows from well known topological results, namely that the image of a compact overt space is compact overt, and that any overt subspace of a discrete space is open [C]. The more significant \Rightarrow comes from the fact that $\mathbf{List}X \rightarrow \mathbf{K}X$. \square

Remark 11.5 We have to be careful with the notion of listability: we may legitimately use a listing to prove results about U only if the statements of those results does not depend on the *choice* of listing [8, §6.6].

Definition 11.6 An object X that is listable in the above sense is called **Kuratowski finite**. If it is also Hausdorff then there is a listing without repetitions, in which case we say that X is simply **finite**. Recall from Propositions 3.8–3.9 that $\mathbf{K}X$ has a greatest element iff X is Kuratowski finite, and is a lattice (in fact a Boolean algebra) iff X is finite.

Encouraged by this success, we use the same Axiom and proof for the

Theorem 11.7 Any $\Gamma \vdash L : \Omega$ is admissible iff it is modal.

Proof We have a directed union in the sense of Axiom 1.10,

$$\pi : \Sigma^X, x : X \vdash \pi x = \exists \ell: \mathbf{K}X. (\forall y \in \ell. \pi y) \wedge (x \in \ell).$$

This is preserved by any $\Gamma \vdash N : \Sigma^{\Sigma^X}$, so

$$\begin{aligned} \Gamma \vdash \top &= N(\lambda x. P\{x\}) && \text{7th modal law} \\ &= N(\exists \ell. (\forall y \in \ell. P\{y\}) \wedge (\lambda x. x \in \ell)) && \text{above} \\ &= \exists \ell. (\forall y \in \ell. P\{y\}) \wedge N(\lambda x. x \in \ell) && \text{Axiom 1.10} \\ &= \exists \ell. (\ell \subset (N, P)) \wedge (N, P) \subset \ell) && \text{Notation 3.3} \\ &= \exists \ell: \mathbf{K}X. (\ell = (N, P)), && \text{Corollary 3.11} \end{aligned}$$

where ℓ is unique and admissible, so (N, P) is itself admissible. \square

What are the results for parametric $\Gamma \vdash L : \mathbf{K}X$ corresponding to the compact open subspace $U \subset X$ above? If we had a theory of parametric

types, in place of a single object U , we would have a **display map** $U \multimap \Gamma$ [8, Chapter VIII]. The idea that each U_γ is overt compact is expressed by saying that $p : U \multimap \Gamma$ is an open proper map [C, §7]. Also, where $U \subset X$ was an open subspace, $U \subset \Gamma \times X$ is an open binary relation $\Gamma \multimap X$. We call this a **Kuratowski-finite subset of X dependent on Γ** .

Proposition 11.8 Let $\Gamma \vdash (N, P) : \mathbf{K}X$. Then there is a diagram as shown, in which the open subspace $i : U \multimap X \times \Gamma$ is classified by $\lambda x \gamma. P\gamma\{x\} : \Sigma^{X \times \Gamma}$, the three squares are pullbacks, and p is an open proper map.

$$\begin{array}{ccccc}
 & U & \xrightarrow{\quad} & (\in) & \xrightarrow{\quad} & 1 \\
 & \downarrow i & & \downarrow & & \downarrow \top \\
 p \swarrow & X \times \Gamma & \xrightarrow{\quad} & X \times \mathbf{K}X & \xrightarrow{\quad} & \Sigma \\
 & \downarrow \pi_1 & & \downarrow \pi_1 & & \downarrow \\
 & \Gamma & \xrightarrow{(N, P)} & \mathbf{K}X & &
 \end{array}
 \quad
 \begin{array}{c}
 x, (N', P') \mapsto P'\{x\}
 \end{array}$$

Proof The composite $p = \pi_1 \cdot i$ is open since X is overt. To show that p is proper, we must find the right adjoint of Σ^p and verify the dual Frobenius law [C, Definition 7.3] and Beck–Chevalley condition. In fact, we shall also give another formula for the left adjoint, and I claim that $E \cdot \exists_i = \exists_X^\Gamma \cdot \exists_i = \exists_p \dashv \Sigma^p \dashv A \cdot \exists_i$.

$$\begin{array}{ccccc}
 \Sigma^U & \xleftarrow{\Sigma^i} & \Sigma^{X \times \Gamma} & \xrightarrow{A : \psi \mapsto \lambda \gamma. N\gamma(\lambda x. \psi(x, \gamma))} & \Sigma^\Gamma \\
 & \xrightarrow{\exists_i} & & \xrightarrow{E : \psi \mapsto \lambda \gamma. P\gamma(\lambda x. \psi(x, \gamma))} & \\
 & & \Sigma^p & &
 \end{array}$$

First recall that $(\exists_i \cdot \Sigma^i \psi)(x, \gamma) = \psi x \wedge (x, \gamma) \in U = \psi x \wedge P\gamma\{x\}$, and let $x : X, \gamma : \Gamma, \theta : \Sigma^\Gamma$ and $\psi : \Sigma^{X \times \Gamma}$.

For the left adjoint we require $E \cdot (\exists_i \cdot \Sigma^i) = \exists_X^\Gamma \cdot (\exists_i \cdot \Sigma^i)$.

For the unit of the right adjoint, $\text{id}_{\Sigma^\Gamma} \leq A \cdot (\exists_i \cdot \Sigma^i) \cdot \Sigma^{\pi_1}$.

For the counit, $(\exists_i \cdot \Sigma^i) \cdot (\Sigma^{\pi_1} \cdot A) \cdot (\exists_i \cdot \Sigma^i) \leq (\exists_i \cdot \Sigma^i)$.

For the dual Frobenius law, $A \cdot \exists_i(\phi \vee \Sigma^p \theta) \gamma = A(\exists_i \phi) \gamma \vee \theta \gamma$.

Finally, we must show stability under pullback along $s : \Delta \rightarrow \Gamma$. This follows by application of the same results, but for $\Delta \vdash (s^*N, s^*P) : \mathbf{K}X$.

$$\begin{array}{ccccc}
 \Sigma^{s^*U} & \xleftarrow{\quad} & \Sigma^U & & s^*U \xrightarrow{\quad} U \xrightarrow{\quad} (\in) \xrightarrow{\quad} \mathbf{1} \\
 \downarrow \exists_j & & \downarrow \exists_i & & \downarrow j \quad \downarrow i \quad \downarrow \quad \downarrow \top \\
 \Sigma^{X \times \Delta} & \xleftarrow{\Sigma^{X \times s}} & \Sigma^{X \times \Gamma} & \xrightarrow{X \times s} & X \times \Delta \xrightarrow{X \times s} X \times \Gamma \longrightarrow X \times \mathbf{K}X \longrightarrow \Sigma \\
 \downarrow A' & & \downarrow A & & \downarrow \pi_1 \quad \downarrow \pi_1 \quad \downarrow \pi_1 \\
 \Sigma^\Delta & \xleftarrow{\Sigma^s} & \Sigma^\Gamma & \xrightarrow{s} & \Delta \xrightarrow{s} \Gamma \xrightarrow{(N, P)} \mathbf{K}X
 \end{array}$$

The Beck–Chevalley condition for the top left square is that for the inverse image of an open inclusion [C, Proposition 3.11]. In the the bottom left square, where

$$A'\theta \equiv \lambda\delta. N(s\delta)(\lambda x. \theta(x, \delta)),$$

the Beck–Chevalley condition is

$$\Sigma^s(A\psi)\delta = N(s\delta)(\lambda x. \psi(x, s\delta)) = A'(\Sigma^{X \times s}\psi)\delta. \quad \square$$

Corollary 11.9 In particular, $(\in) \hookrightarrow X \times \mathbf{K}X \rightarrow \mathbf{K}X$ is open and proper. \square

Proposition 11.10 Conversely, let $i : U \hookrightarrow X \times \Gamma$ be open such that $p \equiv (\pi_1 \cdot i) : R \rightarrow \Gamma$ is (open and) proper. Then $N\gamma\phi = \forall_p(\lambda u. \phi(qu))$ and

$$P\gamma\phi = \exists_p(\lambda u. \phi(qu)) = \exists x. \phi x \wedge (x, \gamma) \in U = \exists x. \phi x \wedge P\gamma\{x\}$$

are modal (where $q \equiv (\pi_0 \cdot i) : R \rightarrow X$). We recover U from P as $P\gamma\{x\} = ((x, \gamma) \in U)$. \square

Theorem 11.11 $\mathbf{K}X$ classifies Kuratowski finite subsets of X , in the sense that

$$\begin{array}{c}
 \text{open proper relations } \Gamma \hookrightarrow X \text{ as above} \\
 \hline \hline
 \Gamma \longrightarrow \mathbf{K}X
 \end{array} \quad \square$$

Theorem 11.12 KX classifies Kuratowski finite subsets of X .

$$\begin{array}{ccc}
 U & \xrightarrow{\quad} & (\in) \\
 \downarrow p & \lrcorner & \downarrow \\
 \Gamma & \xrightarrow{\quad} & KX
 \end{array}$$

Indeed $(\in) : X \rightarrow KX$ is the generic Kuratowski-finite subset of X (dependent on KX). By this we mean that any pullback of it as shown is a Kuratowski-finite subset of X dependent on Γ , and every Kuratowski-finite subset of X dependent on Γ arises uniquely in this way. \square

Acknowledgement

I would like to thank Peter Aczel, Andrej Bauer, Robin Cockett, Maria Emilia Maietti and Gavin Wraith for their helpful comments. This paper was accepted for *Category Theory and Computer Science* **10**, which was held in Copenhagen in August 2004, and an abridged version appeared in the printed proceedings. However, I was unable to attend, as I had broken my leg two weeks earlier. Research on Abstract Stone Duality has been funded since September 2003 by the EPSRC project GR/S58522 of the same title, but the basic results of Sections 3–6 had been found in July–August 2003.

References

- [1] Cockett, J. R., *List-arithmetic distributive categories: Loco!*, Journal of Pure and Applied Algebra **66** (1990), pp. 1–29.
- [2] Hofmann, M., “Extensional concepts in intensional type theory,” Ph.D. thesis, University of Edinburgh (1995).
URL <http://www.lfcs.inf.ed.ac.uk/reports/95/ECS-LFCS-95-327>
- [3] Johnstone, P., “Stone Spaces,” Number 3 in Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1982.
- [4] Johnstone, P. T., *Vietoris locales and localic semi-lattices*, in: R.-E. Hoffmann and K. H. Hoffmann, editors, *Continuous Lattices and their Applications*, Marcel Dekker, 1985 pp. 155–180.
- [5] Lambek, J. and P. J. Scott, “Introduction to Higher Order Categorical Logic,” Cambridge University Press, 1986.
- [6] Maietti, M. E., *Joyal’s arithmetic universes via type theory*, Electronic Notes in Theoretical Computer Science **69** (2003).
- [7] Maietti, M. E., *Reflection into models of finite decidable FP-sketches in an arithmetic universe*, Electronic Notes in Theoretical Computer Science **416** (2005).

- [8] Taylor, P., “Practical Foundations of Mathematics,” Cambridge University Press, 1999.
- [9] Vickers, S. J., “Topology Via Logic,” Cambridge Tracts in Theoretical Computer Science **5**, Cambridge University Press, 1988.
- [10] Winskel, G., *Powerdomains and modality*, in: M. Karpinski, editor, *Foundations of Computation Theory*, number 158 in Lecture Notes in Computer Science (1983), pp. 505–514.

My papers on abstract Stone duality may be obtained from
www.cs.man.ac.uk/~pt/ASD

- [A] [Sober spaces and continuations](#). *Theory and Applications of Categories*, 10(12):248–299, 2002.
- [B] [Subspaces in abstract Stone duality](#). *Theory and Applications of Categories*, 10(13):300–366, 2002.
- [C] [Geometric and higher order logic using abstract Stone duality](#). *Theory and Applications of Categories*, 7(15):284–338, 2000.
- [D] [Non-Artin gluing in recursion theory and lifting in abstract Stone duality](#). 2000.
- [E] [Inside every model of Abstract Stone Duality lies an Arithmetic Universe](#). *Category Theory and Computer Science 10*, Copenhagen, 2004.
- [F] [Scott domains in abstract Stone duality](#). March 2002.
- [G–] [Local compactness and the Baire category theorem in abstract Stone duality](#). *Electronic Notes in Theoretical Computer Science* **69**, Elsevier, 2003.
- [G] [Computably based locally compact spaces](#). May 2003.
- [H–] [An elementary theory of the category of locally compact locales](#). APPSEM Workshop, Nottingham, March 2003.
- [H] [An elementary theory of various categories of spaces and locales](#). November 2004.
- [I] [The Dedekind reals in abstract Stone duality](#). December 2004.
- [J] [A \$\lambda\$ -calculus for real analysis](#). December 2004.