

Problem 1 (HW 7 Problem 7)

Let $X : \Omega \mapsto [0, \infty)$ be a random variable with distribution μ , distribution function

$$F(x) = P(X \leq x) = \mu((-\infty, x]) = \mu([0, x]) = \int_{\mathbb{R}} I_{[0, x]}(\xi) d\mu(\xi)$$

and Laplace transform for $\lambda \in (0, \infty)$:

$$\hat{F}(\lambda) = \mathbb{E}[e^{-\lambda X}] = \int_0^\infty e^{-\lambda \xi} d\mu(\xi)$$

Show that the Laplace transform \hat{F} determines the distribution function F , using the strong law of large numbers and the following steps:

(a) Show that the derivatives of this Laplace transform \hat{F} are given by

$$\frac{\partial^k}{\partial \lambda^k} \hat{F}(\lambda) = \hat{F}^{(k)}(\lambda) = \int_0^\infty (-1)^k \xi^k e^{-\lambda \xi} d\mu(\xi), \quad \lambda \in (0, \infty), k \in \mathbb{N}$$

(b) Suppose that $Z_1(\theta), Z_2(\theta), \dots$ are independent random variables with the same Poisson distribution of parameter $\theta = \mathbb{E}[Z_k(\theta)]$. Use the weak law of large numbers, to argue that

$$P\left(\frac{1}{n} \sum_{k=1}^n Z_k(\theta) \leq x\right) \rightarrow \begin{cases} 1 & \text{if } x > \theta \\ 0 & \text{if } x < \theta \end{cases}$$

and thus

$$\sum_{j \leq nx} e^{-n\theta} \frac{(n\theta)^j}{j!} \rightarrow \begin{cases} 1 & \text{if } x > \theta \\ 0 & \text{if } x < \theta \end{cases}$$

What is the above limit, if $x = \theta$?

(c) Conclude that, for any $x \in [0, \infty)$ which is a continuity point of the distribution function F , we have

$$\sum_{j \leq nx} \frac{(-1)^j}{j!} n^j \hat{F}^{(j)}(n) \rightarrow F(x)$$

(d) Argue that the distribution function F (thus also the measure μ) is determined uniquely from the Laplace transform \hat{F} .

(a) Consider the difference quotient:

$$\frac{\hat{F}(\lambda + h) - \hat{F}(\lambda)}{h} = \int_0^\infty \frac{e^{-(\lambda+h)\xi} - e^{-\lambda\xi}}{h} d\mu(\xi), \quad \text{where } \frac{e^{-(\lambda+h)\xi} - e^{-\lambda\xi}}{h} \rightarrow -\xi e^{-\lambda\xi}$$

Notice that when h is small enough, we always have

$$\left| \frac{e^{-(\lambda+h)\xi} - e^{-\lambda\xi}}{h} \right| \leq 1 + \xi e^{-\lambda\xi}$$

We claim that $1 + \xi e^{-\lambda \xi}$ is integrable. Since μ is a probability measure, 1 is integrable. Since $\lambda > 0$, $\xi e^{-\lambda \xi}$ is nonnegative and vanishes very fast as $\xi \rightarrow \infty$, the integral of $\xi e^{-\lambda \xi}$ over $[0, \infty)$ is also finite. Thus, the claim is true and we can apply DCT to conclude

$$\lim_{h \rightarrow 0} \frac{\hat{F}(\lambda + h) - \hat{F}(\lambda)}{h} = \int_0^\infty -\xi e^{-\lambda \xi} d\mu(\xi)$$

Thus, the desired result is true for $k = 1$. Repeat the same argument inductively, based on the fact that $e^{-\lambda \xi}$ is smooth and $\xi^k e^{-\lambda \xi}$ is integrable for any k , DCT implies the desired result is also true for any k .

(b) By weak law of large number, $\bar{Z}_k \rightarrow \theta$ in probability, and hence in distribution, so

$$\lim_{n \rightarrow \infty} P(\bar{Z}_k \leq x) = P(\theta \leq x) = \begin{cases} 1 & \text{if } x > \theta \\ 0 & \text{if } x < \theta \end{cases}$$

The case $x = \theta$ is not valid because convergence in distribution is only valid for continuous point, but $x = \theta$ is not continuous. However, consider

$$P(\bar{Z}_k \leq \theta) = P(\sqrt{n}(\bar{Z}_k - \theta)/\theta \leq 0) \rightarrow P(Z \leq 0) = \frac{1}{2}$$

where Z follows standard normal distribution by CLT. Since the sum of n i.i.d. $\text{Poisson}(\theta)$ is $\text{Poisson}(n\theta)$, we have

$$P(\bar{Z}_k \leq x) = P\left(\sum_{k=1}^n Z_k(\theta) \leq nx\right) = \sum_{j \leq nx} e^{-n\theta} \frac{(n\theta)^j}{j!}$$

and the desired result follows.

(c) Denote the RHS of (b) as $f(\theta; x)$ and LHS as $f_n(\theta; x)$. Notice that $f_n(\theta; x) \rightarrow f(\theta; x)$ a.e. with respect to θ , and $|f_n(\theta; x)| \leq 1$. Since F is a probability distribution, 1 is integrable, so by DCT,

$$\int_0^\infty \sum_{j \leq nx} e^{-n\theta} \frac{(n\theta)^j}{j!} d\mu(\xi) = \int_0^\infty f_n(\theta; x) d\mu(\xi) \rightarrow \int_0^\infty f(\theta; x) d\mu(\xi) = \int_0^x 1 d\mu(\xi) = F(x)$$

Observe that

$$\int_0^\infty \sum_{j \leq nx} e^{-n\theta} \frac{(n\theta)^j}{j!} d\mu(\xi) = \sum_{j \leq nx} \frac{n^j}{j!} \int_0^\infty e^{-n\theta} \theta^j d\mu(\xi) = \sum_{j \leq nx} \frac{n^j}{j!} (-1)^j \hat{F}^{(j)}(n)$$

where the last equality follows from the result in (a). Thus, the desired statement follows.

(d) From (c), we know F is uniquely determined from \hat{F} for all continuous points. Since F is a distribution function, it is right continuous with left limit and the discontinuous point is only countably many, so by taking the limit of a sequence of continuous point, every x is uniquely determined for $F(x)$.

Problem 2 (HW 7 Problem 8)

For $s > 1$ fixed, let $X : \Omega \mapsto \mathbb{N}$ be a random variable with the “zeta distribution”:

$$P(X = n) = \frac{1}{\zeta(s)n^s}, \quad n \in \mathbb{N}, \quad \zeta(s) = \sum_{n \in \mathbb{N}} \frac{1}{n^s}$$

where $\zeta(s)$ is the Riemann’s zeta-function. For each $m \in \mathbb{N}$, consider the event

$$A_m = \{\omega \in \Omega : m \text{ is a factor of } X(\omega)\} = \{\omega \in \Omega : X(\omega) = mk(\omega), \text{ for some } k(\omega) \in \mathbb{N}\}$$

- (i) Show that $P(A_m) = 1/m^s$.
- (ii) Conclude from (i) that the events in the collection $\{A_p\}_{p \in \mathcal{P}}$ are independent, where \mathcal{P} is the set of prime numbers.
- (iii) Use (i) and (ii) to derive the Euler formula

$$\frac{1}{\zeta(s)} = \prod_{p \in \mathcal{P}} \left(1 - \frac{1}{p^s}\right)$$

- (i) Notice that by definition, we can compute $P(A_m)$ by

$$P(A_m) = \sum_{k \in \mathbb{N}} \frac{1}{\zeta(s)(mk)^s} = \frac{1}{m^s \zeta(s)} \sum_{k \in \mathbb{N}} \frac{1}{k^s} = \frac{1}{m^s \zeta(s)} \zeta(s) = \frac{1}{m^s}$$

and the desired result follows.

- (ii) By definition of independence, it suffices to show that for any finite subsequence p_1, \dots, p_K of the collection $\{A_p\}_{p \in \mathcal{P}}$, we have $P(A_{p_1}, \dots, A_{p_K}) = \prod_{i=1}^K P(A_{p_i})$. To compute $P(A_{p_1}, \dots, A_{p_K})$, consider

$$P(A_{p_1}, \dots, A_{p_K}) = P(X = mp_1 \cdots p_K) = P(A_{p_1 \cdots p_K}) = \frac{1}{(p_1 \cdots p_K)^s} = \prod_{i=1}^K P(A_{p_i})$$

Thus, $\{A_p\}_{p \in \mathcal{P}}$ is an independent collection.

- (iii) Note that $P(X = 1) = \frac{1}{\zeta(s)}$ is the LHS of the equation. Observe that

$$P(X = 1) = P(\cap_{p \in \mathcal{P}} A_p^c) = \prod_{p \in \mathcal{P}} P(A_p^c) = \prod_{p \in \mathcal{P}} \left(1 - \frac{1}{p^s}\right)$$

because of the independence proved in (ii) and the formula in (i).