

# Part II Topics in Quantum Theory

William Royce

October 4, 2024

Part II Physics, The University of Cambridge



# Preface

---



# Contents

---

<b>1</b>	<b>Quantum Dynamics</b>	<b>1</b>
1.1	The Quantum Harmonic Oscillator . . . . .	1
1.1.1	Time Independent Case . . . . .	1
1.1.2	Time Dependent Force . . . . .	4
1.2	A Spin in a Field . . . . .	6
1.2.1	Rabi Oscillations . . . . .	9
1.3	The Adiabatic Approximation . . . . .	9
1.3.1	Landau–Zener Tunnelling . . . . .	11
<b>2</b>	<b>Introduction to Path Integrals</b>	<b>15</b>
<b>3</b>	<b>Scattering Theory</b>	<b>17</b>
<b>4</b>	<b>Identical Particles in Quantum Mechanics</b>	<b>19</b>
<b>5</b>	<b>Density Matrices</b>	<b>21</b>
<b>6</b>	<b>Lie Groups [<i>non-examinable</i>]</b>	<b>23</b>
<b>7</b>	<b>Relativistic Quantum Physics [<i>non-examinable</i>]</b>	<b>25</b>
<b>A</b>	<b>Appendix: Operator Algebra</b>	<b>A.1</b>
A.1	General Operator Algebra . . . . .	A.1

## List of Tables

## List of Figures

---

1.1	The action of a periodic trajectory is equal to the area enclosed in the phase plane. For a simple harmonic oscillator the curve is an ellipse and the action is the product of the energy and the period. If the period of the oscillator is altered slowly (by changing the length of a pendulum, say) the ellipse will distort but the area will remain fixed. . . . .	12
1.2	Instantaneous eigenvalues of the Landau–Zener problem. The dotted line schematically illustrates what happens when we pass over the branch point.	12
1.3	: (Top) Branch cut and contour of time evolution in the complex $t$ plane. (Bottom) Riemann surface of $\sqrt{(\beta t)^2 + \Delta^2}$ (real part). . . . .	12
1.4	Caption . . . . .	13



## CHAPTER 1

# Quantum Dynamics

---

In this first chapter we're going to introduce some general ideas of quantum dynamics, using the two simplest quantum systems: the harmonic oscillator and a single spin-1/2.

## 1.1 The Quantum Harmonic Oscillator

There's an old crack from the late quantum field theorist Sidney Coleman to the effect that

*The career of a young theoretical physicist consists of treating the harmonic oscillator in ever-increasing levels of abstraction.*

There's a large kernel of truth in this, for the simple reason that many systems in physics vibrate, from bridges to quantum fields, and within a certain approximation that vibration can be treated as harmonic. In this section we are going to remind ourselves about some features of quantum dynamics using this model as our basic example, as it allows most results to be expressed analytically. Along the way I'll try and point out which features generalise to more complicated systems (and which don't!).

### 1.1.1 Time Independent Case

The Hamiltonian is

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2, \quad (1.1)$$

where the position and momentum operators satisfy

$$[\hat{x}, \hat{p}] = i\hbar. \quad (1.2)$$

The state of the oscillator  $\bar{\psi}$  evolves in time according to the (time dependent) Schrödinger equation

$$i\hbar \frac{d}{dt}\bar{\psi} = \hat{H}\bar{\psi}. \quad (1.3)$$

This is a first order differential equation, and so the evolution is fixed once the initial state  $\psi(0)$  is specified. We can write the solution as

$$\psi(t) = \exp(-i\hat{H}t/\hbar)\psi(0) \equiv \hat{U}(t)\psi(0). \quad (1.4)$$

The operator  $\hat{U}(t) = e^{-i\hat{H}t/\hbar}$  is called the *evolution operator*, as it evolves the state  $\psi(0)$  forward in time.

Functions of operators can be thought of as defined by their power series expansions, in this case

$$\hat{U}(t) = \mathbb{I} - i\frac{\hat{H}t}{\hbar} - \frac{1}{2}\left(\frac{\hat{H}t}{\hbar}\right)^2 + \dots \quad (1.5)$$

Alternatively, if an operator has a complete orthonormal eigenbasis  $|n\rangle$  (as quantum observables do, being Hermitian operators), we can write any such function in terms of this basis and the corresponding function of the eigenvalues  $E_n$

$$\hat{U}(t) = \sum_n e^{-iE_n t/\hbar} |n\rangle \langle n|. \quad (1.6)$$

This latter point of view then focuses attention on the eigenstates  $|n\rangle$ . To find these there are at least two approaches.

1. **Brute force:** Take the position representation  $\hat{p} = -i\hbar d/dx$  and study the time independent Schrödinger equation in this representation

$$-\frac{\hbar^2}{2m} \frac{d^2\psi_n}{dx^2} + \frac{1}{2}m\omega^2 x^2 \psi_n = E_n \psi_n, \quad (1.7)$$

where  $\langle x|n\rangle = \psi_n(x)$ . The result is that the eigenfunctions have the form

$$\psi_n(x) = \frac{1}{\sqrt{2^n n!}} \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-m\omega x^2/2\hbar} H_n\left(\sqrt{\frac{m\omega}{\hbar}}x\right), \quad (1.8)$$

with eigenvalues  $E_n = \hbar\omega(n + 1/2)$ , where  $H_n(z)$  are the Hermite polynomials.

2. **More Sophisticated:** Define the hermitian conjugate pair

$$\begin{aligned} \hat{a} &= \sqrt{\frac{m\omega}{2\hbar}} \left( \hat{x} + i\frac{\hat{p}}{m\omega} \right) \\ \hat{a}^\dagger &= \sqrt{\frac{m\omega}{2\hbar}} \left( \hat{x} - i\frac{\hat{p}}{m\omega} \right), \end{aligned} \quad (1.9)$$

which satisfy

$$[\hat{a}, \hat{a}^\dagger] = 1. \quad (1.10)$$

The Hamiltonian is expressed as

$$\hat{H} = \frac{\hbar\omega}{2} [\hat{a}^\dagger \hat{a} + \hat{a} \hat{a}^\dagger] = \hbar\omega \left( N + \frac{1}{2} \right), \quad (1.11)$$

where  $N \equiv \hat{a}^\dagger \hat{a}$ . The commutation relation Eq. (1.10) implies

$$[N, \hat{a}] = -\hat{a}, \quad [N, \hat{a}^\dagger] = +\hat{a}^\dagger, \quad (1.12)$$

which in turn tells us that acting with  $\hat{a}^\dagger(\hat{a})$  on an eigenstate  $|n\rangle$  of  $N$  with eigenvalue  $n$  gives another eigenstate with eigenvalue increased (decreased) by 1.

Alternatively, we can try and find  $\hat{U}(t)$  indirectly, from the effect it has on operators. Recall that in the *Heisenberg picture* operators acquire a time dependence

$$\hat{O}(t) = \hat{U}^\dagger(t) \hat{O} \hat{U}(t), \quad (1.13)$$



equivalent to the Heisenberg equation of motion

$$\frac{d\hat{\mathcal{O}}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{\mathcal{O}}(t)]. \quad (1.14)$$

Let's see what this means for the Harmonic oscillator. Evidently  $\hat{H} = \hat{U}^\dagger(t) \hat{H} \hat{U}(t)$ , so

$$\hat{H} = \hat{U}^\dagger(t) \hat{H} \hat{U}(t) = \frac{\hat{p}(t)^2}{2m} + \frac{1}{2} m \omega^2 \hat{x}(t)^2. \quad (1.15)$$

We have

$$\frac{d\hat{x}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{x}(t)] = \frac{\hat{p}(t)}{m} \quad (1.16)$$

$$\frac{d\hat{p}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{p}(t)] = -m\omega^2 \hat{x}(t). \quad (1.17)$$

You may recognize these equations as identical to *Hamilton's equations* for the SHO

$$\frac{d\hat{x}}{dt} = \frac{\partial \hat{H}}{\partial \hat{p}} = \frac{\hat{p}}{m} \quad (1.18)$$

$$\frac{d\hat{p}}{dt} = -\frac{\partial \hat{H}}{\partial \hat{x}} = -m\omega^2 \hat{x}. \quad (1.19)$$

Considering the case for the more general Hamiltonian,

$$\hat{H} = \hat{T}(\hat{p}) + \hat{V}(\hat{x}) = \sum_n \frac{t_n \hat{p}^n}{n!} + \sum_n \frac{v_n \hat{x}^n}{n!}, \quad (1.20)$$

we find the Heisenberg equation of motion

$$\frac{d\hat{x}(t)}{dt} = \frac{i}{\hbar} \sum_n \left[ \frac{t_n \hat{p}^n}{n!}, \hat{x} \right] = \hat{T}'(\hat{p}), \quad (1.21)$$

which corresponds to Hamilton's equation  $\dot{\hat{x}} = \partial \hat{H} / \partial \hat{p}$ . The same proof works for  $d\hat{p}(t)/dt = -\hat{V}'(\hat{x})$ .

The general solution to Eqs. (1.18) and (1.19) is

$$\hat{x}(t) = \cos(\omega t) \hat{x}(0) + \sin(\omega t) \frac{\hat{p}(0)}{m\omega} \quad (1.22)$$

$$\hat{p}(t) = \cos(\omega t) \hat{p}(0) - m\omega \sin(\omega t) \hat{x}(0), \quad (1.23)$$

and corresponds to a point tracing out an elliptical trajectory centred at the origin in the  $x-p$  plane (*phase space*). From this point of view the operators  $\hat{a}$ ,  $\hat{a}^\dagger$  in Eq. (1.9) can be seen as complex amplitudes whose phase changes linearly in time

$$\hat{a}(t) = e^{-i\omega t} \hat{a}(0), \quad \hat{a}^\dagger(t) = e^{+i\omega t} \hat{a}^\dagger(0). \quad (1.24)$$

This follows by writing the Hamiltonian as  $\hat{H} = \omega(\hat{a}^\dagger \hat{a} + \frac{1}{2})$ , such that  $\dot{\hat{a}}(t) = -i\omega \hat{a}(t)$ , which can be integrated to yield  $\hat{a}(t) \exp(-i\omega t) \hat{a}(0)$ .

### 1.1.2 Time Dependent Force

Mostly we don't leave quantum systems to get on with their own time evolution, but disturb them in some way. For example, an atom may experience an external radiation field. The prototype for this situation is the SHO subject to a time dependent force<sup>1</sup>

$$\hat{H}(t) = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2 - F(t)\hat{x}. \quad (1.25)$$

How does such a system evolve? The important thing to realise is that the solution of the Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} |\psi\rangle = \hat{H}(t) |\psi\rangle \quad (1.26)$$

is *not*

$$\hat{U}(t) \neq \exp(-i\hat{H}(t)t/\hbar). \quad (1.27)$$

Let's consider the situation described by

$$F(t) = \begin{cases} F_1 & 0 \leq t < t_1 \\ F_2 & t_1 \leq t < t_2 \end{cases}. \quad (1.28)$$

The evolution operator is

$$\hat{U}(t) = \begin{cases} U_1(t) & 0 \leq t < t_1 \\ U_2(t - t_1)U_1(t_1) & t_1 \leq t < t_2 \end{cases}, \quad (1.29)$$

where  $U_i = e^{-i\hat{H}_i t/\hbar}$  and  $\hat{H}_i = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2 - F_i\hat{x}$ . It's important to realise that  $\hat{H}_1$  and  $\hat{H}_2$  don't commute with each other,

$$[\hat{H}_1, \hat{H}_2] = -i\hbar \frac{\hat{p}}{m} (F_1 - F_2), \quad (1.30)$$

thus  $U_1$  and  $U_2$  don't commute and the product of  $U_1$  and  $U_2$  in Eq. (1.29) is not easily written in terms of a single exponential.

The evolution operator corresponding to a general force  $F(t)$  can be understood by splitting the evolution up into many small stages

$$\begin{aligned} \hat{U}(t) &= \lim_{\Delta t \rightarrow 0} e^{-i\hat{H}(t-\Delta t)\Delta t/\hbar} e^{-i\hat{H}(t-2\Delta t)\Delta t/\hbar} \dots e^{-i\hat{H}(\Delta t)\Delta t/\hbar} e^{-i\hat{H}(0)\Delta t/\hbar} \\ &= \lim_{\Delta t \rightarrow 0} \left(1 - \frac{i\hat{H}(t-\Delta t)\Delta t}{\hbar}\right) \left(1 - \frac{i\hat{H}(t-2\Delta t)\Delta t}{\hbar}\right) \dots \left(1 - \frac{i\hat{H}(0)\Delta t}{\hbar}\right) \\ &= 1 - \frac{i}{\hbar} \int_0^t dt_1 \hat{H}(t_1) - \frac{1}{\hbar^2} \int_0^t dt_2 \int_0^{t_2} dt_1 \hat{H}(t_2)\hat{H}(t_1) + \dots \end{aligned} \quad (1.31)$$

Note that the time arguments of  $\hat{H}(t)$  are increasing from right to left. The final expression for  $\hat{U}(t)$  can be written in a dangerously compact fashion by using the notation

$$\mathcal{T}[\hat{H}(t_1)\hat{H}(t_2)] = \begin{cases} \hat{H}(t_1)\hat{H}(t_2) & t_1 \geq t_2 \\ \hat{H}(t_2)\hat{H}(t_1) & t_2 > t_1 \end{cases}, \quad (1.32)$$

---

<sup>1</sup>Recall that in the Heisenberg picture the Hamiltonian remained time independent. Now it has intrinsic time dependence.

and so on. The operation denoted by  $\mathcal{T}$  is usually called *time ordering*. We have

$$\hat{U}(t) = 1 - \frac{i}{\hbar} \int_0^t dt_1 \hat{H}(t_1) - \frac{1}{2\hbar^2} \int_0^t dt_2 \int_0^{t_2} dt_1 \mathcal{T}[\hat{H}(t_1)\hat{H}(t_2)] + \dots \quad (1.33)$$

Allowing the integrals to range over  $0 < t_i < t$  instead of ordering them necessitates the introduction of a factor  $\frac{1}{n!}$  at the  $n^{\text{th}}$  order. This allows us to write

$$\hat{U}(t) = \mathcal{T} \exp\left(-\frac{i}{\hbar} \int_0^t dt' \hat{H}(t')\right) \quad (1.34)$$

*This expression should be handled with extreme care!* It evidently reduces to  $e^{-i\hat{H}t/\hbar}$  in the case of a time-independent Hamiltonian. In the general case, it is only really useful in the form of the expansion Eq. (1.33).

To make progress in the case of the driven oscillator, it's useful to once again consider the Heisenberg equations of motion

$$\frac{d\hat{x}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{x}(t)] = \frac{\hat{p}(t)}{m} \quad (1.35)$$

$$\frac{d\hat{p}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{p}(t)] = -m\omega^2 \hat{x}(t) + F(t). \quad (1.36)$$

In terms of  $\hat{a}$  and  $\hat{a}^\dagger$

$$\hat{H}(t) = \frac{\hbar\omega}{2} (\hat{a}^\dagger \hat{a} + \hat{a} \hat{a}^\dagger) - F(t) \sqrt{\frac{\hbar}{2m\omega}} (\hat{a} + \hat{a}^\dagger), \quad (1.37)$$

and we have

$$\frac{d\hat{a}}{dt} = -i\omega \hat{a} + iF(t) \sqrt{\frac{1}{2m\hbar\omega}}. \quad (1.38)$$

If we define  $\tilde{\hat{a}}(t)e^{i\omega t}\hat{a}(t)$ , we get

$$\frac{d\tilde{\hat{a}}}{dt} = +i \frac{F(t)e^{i\omega t}}{\sqrt{2m\hbar\omega}}, \quad (1.39)$$

with general solution

$$\tilde{\hat{a}}(t) = \tilde{\hat{a}}(0) + \frac{i}{\sqrt{2m\hbar\omega}} \int_0^t F(t') e^{i\omega t'} dt', \quad (1.40)$$

and similarly

$$\tilde{\hat{a}}^\dagger(t) = \tilde{\hat{a}}^\dagger(0) - \frac{i}{\sqrt{2m\hbar\omega}} \int_0^t F(t') e^{-i\omega t'} dt'. \quad (1.41)$$

From this we have

$$\hat{x}(t) = \frac{1}{2} \sqrt{\frac{2\hbar}{m\omega}} (\hat{a}(t) + \hat{a}^\dagger(t)) = \hat{x}(0) - \frac{1}{m\omega} \int_0^t F(t') \sin(\omega t') dt', \quad (1.42)$$

this is the same as for a classical harmonic oscillator (in the rotating frame).

What can we do with this solution? Suppose we start from the ground state, which satisfies

$$\hat{a} |0\rangle = 0. \quad (1.43)$$

Since  $\hat{a}(t)\hat{U}^\dagger(t)\hat{a}\hat{U}(t)$  we have  $\hat{a}\hat{U}(t) = \hat{U}(t)\hat{a}(t)$  and thus

$$\hat{a}\hat{U}(t)|0\rangle = \frac{i}{\sqrt{2m\hbar\omega}} \int_0^t F(t')e^{i\omega(t'-t)} dt' \hat{U}(t)|0\rangle, \quad (1.44)$$

we have that  $\hat{U}(t)|0\rangle$  is an eigenstate of  $\tilde{\hat{a}}(0) = \hat{a}$  with eigenvalue

$$\frac{i}{\sqrt{2m\hbar\omega}} \int_0^t F(t')e^{i\omega(t'-t)} dt', \quad (1.45)$$

in other words, it is a *coherent state*. Recall from AQP that a coherent state  $|\alpha\rangle$  is defined as an eigenstate of  $\hat{a}$  with eigenvalue  $\alpha$  (generally complex, as  $\hat{a}$  is not Hermitian)

$$\hat{a}|\alpha\rangle = \alpha|\alpha\rangle. \quad (1.46)$$

The explicit form of a normalised coherent state is

$$|\alpha\rangle = e^{-|\alpha|^2/2} e^{\alpha\hat{a}^\dagger} |0\rangle, \quad (1.47)$$

where both the property Eq. (1.46) and the normalisation follow from the fundamental commutator  $[\hat{a}, \hat{a}^\dagger] = 1$ . Given Eq. (1.47),

$$\hat{a}|\alpha\rangle = e^{-|\alpha|^2/2} [\hat{a}, e^{\alpha\hat{a}^\dagger}] |0\rangle = \alpha|\alpha\rangle. \quad (1.48)$$

For the normalisation, we have

$$\langle\alpha|\alpha\rangle = e^{-|\alpha|^2} \sum_{n,m} \langle 0 | \frac{(\alpha^*\hat{a})^n}{n!} \frac{(\alpha\hat{a}^\dagger)^m}{m!} | 0 \rangle = e^{-|\alpha|^2} \sum_n \frac{|\alpha|^{2n}}{n!} = 1. \quad (1.49)$$

Note that the ground state  $|0\rangle$  is a coherent state with  $\alpha = 0$ .

Time-dependent perturbation theory (see AQP notes) gives

$$c_n^{(1)} = -\frac{i}{\hbar} \int_{t_0}^t dt' e^{i\omega_{n,i}t'} V_{n,i}(t'). \quad (1.50)$$

In our case the energy different between  $|0\rangle$  and  $|1\rangle$  is  $\hbar\omega$ , and the matrix element  $V_{1,0}(t') = -F(t')\hbar/(2m\omega)$ . This coincides with (1.45), because

$$\langle 1 | \hat{U}(t) | 0 \rangle = \langle 0 | \hat{U}(t) \hat{a}(t) | 0 \rangle = \langle 0 | \hat{U}(t) | 0 \rangle \frac{i}{\sqrt{2m\hbar\omega}} \int_0^t F(t')e^{i\omega(t'-t)} dt'. \quad (1.51)$$

Then  $\langle 0 | \hat{U}(t) | 0 \rangle = 1 + \mathcal{O}(F^2)$ .

## 1.2 A Spin in a Field

Two state systems abound in physics. Or rather, many physical situations can be approximated by considering only two states. Some important examples are the spin states of the electron, a pair of atomic states coupled by external radiation, and the two equivalent positions of the Nitrogen atom in the trigonal pyramid structure of Ammonia ( $\text{NH}_3$ ).

Quantum two state systems are central to the field of *quantum computing*, where they replace the classical bit of information and are often known as *qubits*.

The simplest quantum system we can write down consists of just two states. The Hilbert space is then two dimensional, and any operator can be thought of as a  $2 \times 2$  matrix. In this section, we'll see that there is a lot to be learnt from this seemingly elementary problem.

It's convenient to describe such a system using the language of spin-1/2, even though the two states may have nothing to do with real spin. The most general time dependent Hamiltonian can then be written using the spin-1/2 operators  $S_i = \frac{1}{2}\sigma_i$  as

$$\hat{H}(t) = \hat{\mathbf{H}}(t) \cdot \hat{\mathbf{S}}, \quad (1.52)$$

in terms of a time dependent 'magnetic field'  $\hat{\mathbf{H}}(t)$  (that again may have nothing to do with a real magnetic field). Using the Pauli matrices, we have the explicit form

$$\hat{H}(t) = \frac{1}{2} \begin{pmatrix} H_z(t) & H_x(t) - iH_y(t) \\ H_x(t) + iH_y(t) & -H_z(t) \end{pmatrix}. \quad (1.53)$$

The Schrödinger equation corresponding to Eq. (1.52) is

$$i\hbar \frac{d|\Psi\rangle}{dt} = \hat{H}(t) |\Psi\rangle, \quad (1.54)$$

where  $|\Psi\rangle = (\psi_\uparrow, \psi_\downarrow)$ .

As before, the formal solution to Eq. (1.54) can be written as

$$|\Psi(t)\rangle = \hat{U}(t, t') |\Psi(t')\rangle, \quad (1.55)$$

In the present case,  $\hat{U}(t, t')$  is a  $2 \times 2$  unitary matrix. It's perhaps a bit surprising that, for this most basic of all possible problems of quantum dynamics, there is no simple relationship between  $\hat{H}(t)$  and  $\hat{U}(t, t')$ . If we think of  $\hat{U}(t, t')$  as representing a kind of rotation in Hilbert space,  $\hat{H}(t)$  corresponds to an instantaneous "angular velocity" describing an infinitesimal rotation. Because these rotations do not commute at different times, the relationship between the infinitesimal rotations and the finite rotation that results is complicated.

The same picture emerges if we look at the Heisenberg equation of motion for  $\hat{\mathbf{S}}(t) = \hat{U}^\dagger(t, t') \hat{\mathbf{S}}(t') \hat{U}(t, t')$ , which take the form

$$\begin{aligned} \frac{d\hat{\mathbf{S}}}{dt} &= \frac{i}{\hbar} [\hat{H}(t), \hat{\mathbf{S}}] \\ &= \frac{1}{\hbar} \hat{\mathbf{H}}(t) \times \hat{\mathbf{S}} \end{aligned} \quad (1.56)$$

by virtue of the spin commutation relations  $[S_i, S_j] = i\varepsilon_{ijk}S_k$ .<sup>2</sup> Thus  $\hat{\mathbf{S}}$  precesses about  $\hat{\mathbf{H}}(t)$ , which corresponds to the instantaneous angular velocity. Differential equations involving operators may make you uncomfortable, but this one is linear and first order, so

---

<sup>2</sup>The usual  $\hbar$  is missing because we defined  $\hat{\mathbf{S}} = \frac{1}{2}\hat{\boldsymbol{\sigma}}$

the solution must be expressible in the form of a matrix connecting the initial and final operators

$$\hat{\mathbf{S}}(t) = \mathbf{R}(t, t') \hat{\mathbf{S}}(t'). \quad (1.57)$$

$\mathbf{R}$  is a  $3 \times 3$  matrix describing the rotation of the spin from time  $t'$  to time  $t$ . The formal expression for  $\mathbf{R}(t, t')$  is

$$\mathbf{R}(t, t') = \mathcal{T} \exp \left( \int_{t'}^t \Omega(t_i) dt_i \right), \quad (1.58)$$

where the matrix  $\Omega(t)$  describing infinitesimal rotations has elements  $\Omega_{jk}(t) = -(1/\hbar)H_i(t)\varepsilon_{ijk}$ , i.e.

$$\Omega(t) = \begin{pmatrix} 0 & -H_z(t) & H_y(t) \\ H_z(t) & 0 & -H_x(t) \\ -H_y(t) & H_x(t) & 0 \end{pmatrix}. \quad (1.59)$$

$\mathbf{U}(t, t')$  and  $\mathbf{R}(t, t')$  contain the same information, of course. We'll return to the relationship between these two in Chapter 6 on Lie Groups.

To find the  $t \rightarrow t'$  time-evolution matrix for a magnetic field in the  $z$ -direction, and the matrix that maps the Heisenberg spin operator at time  $t'$  to the Heisenberg spin operator at  $t$ , we start from Eq. (1.53) to first yield the Hamiltonian

$$\hat{H} = \begin{pmatrix} H/2 & 0 \\ 0 & H/2 \end{pmatrix}. \quad (1.60)$$

This is independent of time, and therefore the time-ordered exponential is just an ordinary matrix exponential because the Hamiltonian cannot fail to commute with itself when evaluated at different times. Moreover it is diagonal in this basis and so the matrix exponential is also easily evaluated:

$$\begin{aligned} \hat{U}(t, t') &= \mathcal{T} \exp \left( -\frac{i}{\hbar} \int_{t'}^t H(t_i) dt_i \right) \\ &= \exp \left( -\frac{i}{\hbar} \begin{pmatrix} H(t-t')/2 & 0 \\ 0 & -H(t-t')/2 \end{pmatrix} \right) \\ &= \begin{pmatrix} \exp \left( -\frac{i}{\hbar} H(t-t')/2 \right) & 0 \\ 0 & \exp \left( \frac{i}{\hbar} H(t-t')/2 \right) \end{pmatrix}. \end{aligned} \quad (1.61)$$

Since  $\Omega_{jk} = -H_i \varepsilon_{ijk}$  and  $H_i = \delta_{iz}H$  we find that

$$\Omega(t) = \begin{pmatrix} 0 & -H & 0 \\ H & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (1.62)$$

which is not diagonal and therefore not trivial to exponentiate even though there is no problem time-ordering. We could perform a diagonalisation, but it is nicer to notice that  $\Omega = iH\sigma_y \oplus 0$ , and use  $e^{i\theta\sigma_a} = \cos \theta + i\sigma_a \sin \theta$  (to prove this, use  $\sigma_a^2 = 1$ ). Then

$$\begin{aligned} \mathbf{R}(t, t') &= \exp(iH(t-t')\sigma_y) \oplus \exp(0) \\ &= (\cos(H(t-t')) + i\sigma_y \sin(H(t-t'))) \oplus 1, \end{aligned} \quad (1.63)$$

with the matrix explicitly

$$\mathbf{R}(t, t') = \begin{pmatrix} \cos(H(t-t')) & -\sin(H(t-t')) & 0 \\ \sin(H(t-t')) & \cos(H(t-t')) & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (1.64)$$

### 1.2.1 Rabi Oscillations

One time dependent situation that we can describe exactly is the rotating field

$$\hat{\mathbf{H}}(t) = \begin{pmatrix} \hat{H}_R \cos(\omega t) \\ \hat{H}_r \sin(\omega t) \\ \hat{H}_0 \end{pmatrix}, \quad (1.65)$$

corresponding to the Hamiltonian

$$\hat{H}(t) = \hat{H}_0 \hat{S}_z + \frac{\hat{H}_R}{2} (\hat{S}_+ e^{-i\omega t} + \hat{S}_- e^{i\omega t}), \quad (1.66)$$

where  $\hat{S}_{\pm} = \hat{S}_x \pm i\hat{S}_y$ . The key to solving the problem is to transform the Schrödinger equation Eq. (1.54) by multiplying by  $\exp(i\omega t \hat{S}_z)$ . Define

$$|\Psi_R(t)\rangle \equiv \exp(i\omega t \hat{S}_z) |\Psi(t)\rangle. \quad (1.67)$$

This transformed state satisfies

$$\begin{aligned} i\hbar \frac{d}{dt} |\Psi_R\rangle &= i\hbar e^{i\omega t \hat{S}_z} \frac{d}{dt} |\Psi\rangle - \hbar\omega \hat{S}_z |\Psi_R\rangle \\ &= e^{i\omega t \hat{S}_z} \hat{H}(t) |\Psi\rangle - \hbar\omega \hat{S}_z |\Psi_R\rangle \\ &= e^{i\omega t \hat{S}_z} \hat{H}(t) e^{-i\omega t \hat{S}_z} |\Psi_R\rangle - \hbar\omega \hat{S}_z |\Psi_R\rangle \\ &= \hat{H}_{\text{Rabi}} |\Psi_R\rangle. \end{aligned} \quad (1.68)$$

In the last line we defined

$$\hat{H}_{\text{Rabi}} \equiv e^{i\omega t \hat{S}_z} \hat{H}(t) e^{-i\omega t \hat{S}_z} - \hbar\omega \hat{S}_z = (\hat{H}_0 - \hbar\omega) \hat{S}_z + \hat{H}_R \hat{S}_x. \quad (1.69)$$

To get the last equality you have to transform the Hamiltonian. You can use Eq. (??), or, since everything is a  $2 \times 2$  matrix, you can multiply the matrices explicitly.

Physically, this corresponds to viewing things in a frame rotating with the field, so the Hamiltonian is now time independent. In this new frame the system precesses about a fixed axis  $(\hat{H}_R, 0, \hat{H}_0 - \hbar\omega)$  at the **Rabi frequency**,

$$\omega_R = \frac{1}{\hbar} \sqrt{(\hat{H}_0 - \hbar\omega)^2 + \hat{H}_R^2}. \quad (1.70)$$

The amplitude of the oscillations in  $\hat{S}_z$  due to this precession is maximal when  $\hat{H}_0 = \hbar\omega$ . In this case the rotation frequency of the field matches the frequency of precession about the  $z$ -axis that would occur if  $\hat{H}_R = 0$ .

## 1.3 The Adiabatic Approximation

The idea of *separation of scales*, be they in length, time, or energy, is endemic in science. If we are interested in studying processes on one scale (such as the weather, say) we hope that they don't depend on the details of processes at another (the motion of molecules).

Rather, we hope that these latter processes can be described in an average way, involving only a few parameters and dynamical quantities (density, local velocity).

The adiabatic approximation is a special case of this idea. Let's suppose that in our two level system, the field  $\hat{\mathbf{H}}(t)$  is changing very slowly (we'll make this idea precise in a moment). If this motion is truly glacial, we'd expect to be able to forget about it altogether, and just solve the problem by finding the energy eigenstates and eigenvalues in the present epoch

$$\hat{H}(t) |\pm_t\rangle = E_{\pm}(t) |\pm_t\rangle. \quad (1.71)$$

We put the  $t$  in a subscript on the states to emphasise that they depend on time as a *parameter*. We refer to the  $|\pm_t\rangle$  as the **instantaneous energy eigenstates**. Although we can always define these states for any  $\hat{\mathbf{H}}(t)$ , we have no reason in general to expect that this  $t$ -dependence has anything to do with the other kind of  $t$ -dependence that arises by solving the time dependent Schrödinger equation.

The **adiabatic theorem** is roughly the statement that these two  $t$  dependences *do* in fact coincide, in the limit that  $\hat{H}(t)$  changes very slowly. To make this more precise, let's expand the state of the system, evolving in time *according to the Schrödinger equation*, in the instantaneous eigenbasis

$$|\Psi(t)\rangle = c_+(t) |+_t\rangle + c_-(t) |-_t\rangle. \quad (1.72)$$

Thus, some of the  $t$  dependence is “carried” by the  $|\pm_t\rangle$ , and by substituting into the Schrödinger equation we are going to find the time dependence of the  $c_{\pm}(t)$ . This involves finding  $d|\pm_t\rangle/dt$ .

Now the following idea you may find a bit odd. Since the time dependence of  $|\pm_t\rangle$  is parametric, we can view the problem of calculating  $d|\pm_t\rangle/dt$  as an exercise in *time independent* perturbation theory.<sup>3</sup> Going from  $t$  to  $t + \delta t$  changes the Hamiltonian by an amount

$$\delta \hat{H}(t) = \frac{d\hat{H}(t)}{dt} \delta t. \quad (1.73)$$

Treating this as a perturbation, the state  $|+_t\rangle$  changes by an amount

$$\delta |+_t\rangle = \frac{\langle -_t | \delta \hat{H}(t) | +_t \rangle}{E_+(t) - E_-(t)} |-_t\rangle, \quad (1.74)$$

so that

$$\frac{d|+_t\rangle}{dt} = \frac{\langle -_t | \dot{\hat{H}}(t) | +_t \rangle}{E_+(t) - E_-(t)} |-_t\rangle. \quad (1.75)$$

Using Eq. (1.75) and the corresponding result for  $d|-_t\rangle/dt$ , we find that the Schrödinger equation gives the following pair of equations for the  $c_{\pm}(t)$

$$i\hbar \frac{d}{dt} \begin{pmatrix} c_+ \\ c_- \end{pmatrix} = \begin{pmatrix} E_+(t) & i\hbar \frac{\langle +_t | \dot{\hat{H}}(t) | -_t \rangle}{E_+ - E_-} \\ i\hbar \frac{\langle -_t | \dot{\hat{H}}(t) | +_t \rangle}{E_- - E_+} & E_-(t) \end{pmatrix} \begin{pmatrix} c_+(t) \\ c_-(t) \end{pmatrix}. \quad (1.76)$$

---

<sup>3</sup>It's a bit like the interaction representation in time dependent perturbation theory.



If  $\hat{H}(t)$  is changing slowly enough, the off-diagonal terms can be neglected and the solution is<sup>4</sup>

$$c_{\pm}(t) = \exp\left(-\frac{i}{\hbar} \int_0^t E_{\pm}(t') dt'\right) c_{\pm}(0). \quad (1.77)$$

Thus the amplitudes evolve independently, and there are no transitions between the instantaneous eigenstates. The phase factor is a generalization of the familiar  $e^{-iE_{\pm}t/\hbar}$  for stationary states, which accounts for the slowly varying instantaneous eigenenergy.

When is this approximation valid? The off-diagonal matrix element in Eq. (1.76) must be small compared to  $E_1(t) - E_2(t)$ , which corresponds to the condition

$$\hbar \left| \langle -_t | \dot{\hat{H}} | +_t \rangle \right| \ll [E_+(t) - E_-(t)]^2. \quad (1.78)$$

*Degeneracy* must be avoided, because the eigenbasis becomes undefined within the degenerate subspace. You can't remain in an eigenstate if you don't know what it is. The approximation is a *semiclassical* one, meaning that it improves at smaller  $\hbar$ .

### 1.3.0.1 Adiabatic

**Adiabatic** is a peculiar term that appears in two related contexts in physics, both referring to slow changes to a system. In thermodynamics, it describes changes without a change in entropy. For reversible changes, this corresponds to no flow of heat, which is the origin of the name (from the Greek for ‘impassable’).

Later, the idea entered mechanics when it was realized that a mechanical system with one degree of freedom undergoing periodic motion, and subject to slow changes, has an **adiabatic invariant**. This turns out to be the **action**

$$S = \oint p dx. \quad (1.79)$$

( $\oint$  indicates that we integrate for one period of the motion) Largely due to the work of Paul Ehrenfest (1880-1933), the invariant played a major role in the “old” quantum theory that predated Schrödinger, Heisenberg, et al.. If the motion of a system is quantised, slow changes to the system's parameters presumably do not lead to sudden jumps. Thus the quantity that comes in quanta must be an adiabatic invariant – and conveniently Planck's constant has the right units. This line of reasoning eventually gave rise to the **Bohr–Sommerfeld quantisation condition**

$$\oint p dx = nh, \quad n \in \mathbb{N}. \quad (1.80)$$

### 1.3.1 Landau–Zener Tunnelling

The picture of adiabatic evolution described above is extremely simple, and it's natural to ask how it breaks down when the condition Eq. (1.78) is not satisfied. Let's consider

---

<sup>4</sup>Note the resemblance to the WKB wavefunction, with energy and time taking the roles of momentum and position. WKB is a kind of adiabatic approximation in space.

**Fig. 1.1:** The action of a periodic trajectory is equal to the area enclosed in the phase plane. For a simple harmonic oscillator the curve is an ellipse and the action is the product of the energy and the period. If the period of the oscillator is altered slowly (by changing the length of a pendulum, say) the ellipse will distort but the area will remain fixed.

time evolution with the Hamiltonian

$$\hat{H}(t) = \begin{pmatrix} \beta t & \Delta \\ \Delta & -\beta t \end{pmatrix}. \quad (1.81)$$

The instantaneous eigenvalues are

$$E_{\pm}(t) = \pm \sqrt{(\beta t)^2 + \Delta^2}. \quad (1.82)$$

We denote the corresponding eigenvalues  $|\pm_t\rangle$ . As a function of  $t$ , the eigenvalues show an **avoided crossing**. The adiabatic theorem tells us that if we start in the state corresponding to the lower energy  $E_-(t)$ , and  $\beta$  is sufficiently small, the state at time  $t$  is

$$\exp\left(-\frac{i}{\hbar} \int_0^t E_-(t') dt'\right) | -_t \rangle, \quad (1.83)$$

where  $| -_t \rangle$  is the corresponding eigenstate. We're integrating from  $t = 0$  because the integral diverges at  $-\infty$  as the phase whizzes faster and faster.

**Fig. 1.2:** Instantaneous eigenvalues of the Landau-Zener problem. The dotted line schematically illustrates what happens when we pass over the branch point.

How small should  $\beta$  be? We use the condition Eq. (1.78), and the fact that the minimum splitting of the energy levels is  $2\Delta$  to arrive at the requirement

$$\frac{\hbar\beta}{\Delta^2} \ll 1. \quad (1.84)$$

We are interested in the situation where this is not the case.

In fact, we can do better than this, via an ingenious excursion into the complex plane. The functions  $E_{\pm}(t)$  have branch cuts starting at  $t = \pm i\Delta/\beta$ . We can think of adiabaticity failing because the branch points are too close to the real axis.

**Fig. 1.3:** (Top) Branch cut and contour of time evolution in the complex  $t$  plane. (Bottom) Riemann surface of  $\sqrt{(\beta t)^2 + \Delta^2}$  (real part).

But who said  $t$  had to be real? There is nothing to stop us integrating the Schrödinger equation along an arbitrary contour. Then we can be as far away from the branch points as we like (Fig. 1.3), and the adiabatic approximation should be valid once more. We can use Eq. (1.83): the exponent now acquires a real part, which describes the decay of the amplitude. Having made the adiabatic approximation, we can deform the contour of integration in Eq. (1.83). The real part of the exponent arising during evolution from  $t = -\infty$  to  $t = +\infty$  can then be written

$$\frac{2i}{\hbar} \int_0^{i\Delta/\beta} \sqrt{(\beta t)^2 + \Delta^2} dt = -\frac{\pi\Delta^2}{2\hbar\beta}, \quad (1.85)$$

**Fig. 1.4:** Caption

giving the modulus of the amplitude

$$|c_-(-\infty \rightarrow +\infty)| = \exp\left(-\frac{\pi\Delta^2}{2\hbar\beta}\right). \quad (1.86)$$

Note, however, that our state is now evolving with an instantaneous energy  $E_+(t)$ , because we passed onto the other sheet of the Riemann surface. We are now in the *upper* state  $|+_t\rangle$ , see Fig. 1.2.

Thus the square of Eq. (1.86) actually gives the probability to make the transition to the upper state. The probability to remain in the lower state is therefore

$$P_{\text{ground}} = 1 - \exp\left(-\frac{\pi\Delta^2}{2\hbar\beta}\right). \quad (1.87)$$

To verify the Landau-Zener result to lowest non-vanishing order in time-dependent perturbation theory, since the result is perturbative in  $\Delta$ , so we take for our unperturbed Hamiltonian  $\hat{H}_0$

$$\hat{H}_0 = \begin{pmatrix} \beta t & 0 \\ 0 & \beta t \end{pmatrix}, \quad (1.88)$$

and our perturbation  $\hat{V}$

$$\hat{V} = \begin{pmatrix} 0 & \Delta \\ \Delta & 0 \end{pmatrix}. \quad (1.89)$$

Since our unperturbed Hamiltonian carries an explicit time dependence, we have to remember to integrate it up when converting between the interaction and Schrödinger pictures:

$$\hat{V}_I(t) = \exp\left(i \int_{t_0}^t \hat{H}_0(t') dt' / \hbar\right) \hat{V}_S(t_0) \exp\left(-i \int_{t_0}^t \hat{H}_0(t') dt' / \hbar\right). \quad (1.90)$$

The Landau-Zener tunnelling appears in this formulation as the transition from the  $-\beta t$  eigenstate to the  $+\beta t$  eigenstate. Using the result of first-order time dependent perturbation theory

$$c_+^{(1)} = -\frac{i}{\hbar} \int_{-\infty}^{\infty} \exp\left(i \int_0^{t'} (E_0^+(t'') - E_0^-(t'')) dt''\right) V_{+-}(t') dt', \quad (1.91)$$

we find

$$\begin{aligned} c_+^{(1)} &= -\frac{i\Delta}{\hbar} \int_{-\infty}^{\infty} e^{i(\beta - (-\beta))t'^2/2} dt' \\ &= -\frac{i\Delta}{\hbar} \sqrt{\frac{\pi\hbar}{i\beta}}. \end{aligned} \quad (1.92)$$

Therefore the transition probability  $P_{-\rightarrow+}$  is given by

$$|c_+^{(1)}|^2 = \frac{\pi\Delta^2}{\beta\hbar}, \quad (1.93)$$

in agreement with the Landau-Zener result to order  $\Delta^2$ .



## CHAPTER 2

# Introduction to Path Integrals

---



## CHAPTER 3

# Scattering Theory

---





## CHAPTER 4

# Identical Particles in Quantum Mechanics

---



## CHAPTER 5

# Density Matrices

---



## CHAPTER 6

# Lie Groups [*non-examinable*]

---



## CHAPTER 7

# Relativistic Quantum Physics [*non-examinable*]

---





# Appendix: Operator Algebra

---

## A.1 General Operator Algebra

Some properties of the **commutator**  $[A, B] = AB - BA$  of operators<sup>1</sup>  $A$  and  $B$

$$[A, BC] = [A, B]C + B[A, C] \tag{A.1}$$

---

<sup>1</sup>In AQP operators wore hats: we'll omit them unless there is a danger of ambiguity