

# Bayes Estimators & Ridge Regression

## Readings ISLR 6

STA 521 Duke University

Merlise Clyde

March 8, 2017

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (x_{ij}^o - \bar{x}_j^o)^2}$$

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (\mathbf{x}_{ij}^o - \bar{\mathbf{x}}_j^o)^2}$$

- ▶ Equivalent to using 'r scale(X)'

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (\mathbf{x}_{ij}^o - \bar{\mathbf{x}}_j^o)^2}$$

- ▶ Equivalent to using 'r scale(X)'
- ▶ Model:

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\beta + \epsilon$$

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (\mathbf{x}_{ij}^o - \bar{\mathbf{x}}_j^o)^2}$$

- ▶ Equivalent to using 'r scale(X)'
- ▶ Model:

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

- ▶  $\mathbf{X}^T \mathbf{X} = (n - 1)\text{Cor}(\mathbf{X})$  (correlation matrix of  $X$ )

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (\mathbf{x}_{ij}^o - \bar{\mathbf{x}}_j^o)^2}$$

- ▶ Equivalent to using 'r scale(X)'
- ▶ Model:

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

- ▶  $\mathbf{X}^T \mathbf{X} = (n - 1) \text{Cor}(\mathbf{X})$  (correlation matrix of  $X$ )
- ▶ Eigenvalue Decomposition  $\mathbf{X}^T \mathbf{X} = \mathbf{U} \boldsymbol{\Lambda} \mathbf{U}^T$

# Model

- ▶ Assume that we have centered (as before) and rescaled  $\mathbf{X}^o$  (original  $\mathbf{X}$ ) so that

$$\mathbf{x}_j = \frac{\mathbf{x}_j^o - \bar{\mathbf{x}}_j^o}{\sum_i (\mathbf{x}_{ij}^o - \bar{\mathbf{x}}_j^o)^2}$$

- ▶ Equivalent to using 'r scale(X)'
- ▶ Model:

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\beta + \epsilon$$

- ▶  $\mathbf{X}^T \mathbf{X} = (n - 1)\text{Cor}(\mathbf{X})$  (correlation matrix of  $X$ )
- ▶ Eigenvalue Decomposition  $\mathbf{X}^T \mathbf{X} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$
- ▶ if smallest eigen value is 0,  $\mathbf{X}$  has columns that are linearly dependent

# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$



# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$

- Consider our expected loss (before we see the data) of taking an “action”  $\mathbf{a}$

# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$

- ▶ Consider our expected loss (before we see the data) of taking an “action”  $\mathbf{a}$
- ▶ Under OLS or the Reference prior the Expected Mean Square Error

# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$

- ▶ Consider our expected loss (before we see the data) of taking an “action”  $\mathbf{a}$
- ▶ Under OLS or the Reference prior the Expected Mean Square Error

$$\mathbb{E}_{\mathbf{Y}}[(\beta - \hat{\beta})^T (\beta - \hat{\beta})] = \sigma^2 \text{tr}[(\mathbf{X}^T \mathbf{X})^{-1}]$$

# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$

- ▶ Consider our expected loss (before we see the data) of taking an “action”  $\mathbf{a}$
- ▶ Under OLS or the Reference prior the Expected Mean Square Error

$$\begin{aligned} \mathbb{E}_{\mathbf{Y}}[(\beta - \hat{\beta})^T (\beta - \hat{\beta})] &= \sigma^2 \text{tr}[(\mathbf{X}^T \mathbf{X})^{-1}] \\ &= \sigma^2 \sum_{j=1}^p \lambda_j^{-1} \end{aligned}$$

# How Good are Various Estimators

Quadratic loss for estimating  $\beta$  using estimator  $\mathbf{a}$

$$L(\beta, \mathbf{a}) = (\beta - \mathbf{a})^T (\beta - \mathbf{a})$$

- ▶ Consider our expected loss (before we see the data) of taking an “action”  $\mathbf{a}$
- ▶ Under OLS or the Reference prior the Expected Mean Square Error

$$\begin{aligned} E_{\mathbf{Y}}[(\beta - \hat{\beta})^T (\beta - \hat{\beta})] &= \sigma^2 \text{tr}[(\mathbf{X}^T \mathbf{X})^{-1}] \\ &= \sigma^2 \sum_{j=1}^p \lambda_j^{-1} \end{aligned}$$

- ▶ If smallest  $\lambda_j \rightarrow 0$  then  $\text{MSE} \rightarrow \infty$

# Problems

Estimates:

$$\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$$

or with  $g$ -prior

$$\hat{\beta} = \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$$

may be unstable

Solutions:

- ▶ remove redundant variables (model selection) (AIC, BIC, other approaches)  $2^p$  models combinatorial hard problem even with Stochastic Search
- ▶ add constant to  $\mathbf{X}^T \mathbf{X}$ :  $\tilde{\beta} = (\mathbf{X}^T \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \mathbf{Y}$  to stabilise eigenvalues - alternative shrinkage estimator

# Independent Prior

- ▶ Reference prior  $p(\beta_0, \phi) \propto \phi^{-1}$
- ▶ Prior Distribution on

$$\beta \mid \phi, \beta_0, k \sim \text{N}(\mathbf{0}_p, \frac{1}{\phi k} \mathbf{I}_p)$$

# Independent Prior

- ▶ Reference prior  $p(\beta_0, \phi) \propto \phi^{-1}$
- ▶ Prior Distribution on

$$\beta \mid \phi, \beta_0, k \sim N(\mathbf{0}_p, \frac{1}{\phi k} \mathbf{I}_p)$$

- ▶ log likelihood (integrated) for  $\beta$  plus prior

$$-\frac{\phi}{2} (\|\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta\|^2 + k\|\beta\|^2)$$

- ▶ Posterior mean

$$\mathbf{b}_n = (\mathbf{X}^T \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \mathbf{X} \hat{\beta}$$



# Independent Prior

- ▶ Reference prior  $p(\beta_0, \phi) \propto \phi^{-1}$
- ▶ Prior Distribution on

$$\beta \mid \phi, \beta_0, k \sim N(\mathbf{0}_p, \frac{1}{\phi k} \mathbf{I}_p)$$

- ▶ log likelihood (integrated) for  $\beta$  plus prior

$$-\frac{\phi}{2} (\|\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta\|^2 + k\|\beta\|^2)$$

- ▶ Posterior mean

$$\mathbf{b}_n = (\mathbf{X}^T \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \mathbf{X} \hat{\beta}$$

- ▶ importance of standardizing

# Independent Prior

- ▶ Reference prior  $p(\beta_0, \phi) \propto \phi^{-1}$
- ▶ Prior Distribution on

$$\beta \mid \phi, \beta_0, k \sim N(\mathbf{0}_p, \frac{1}{\phi k} \mathbf{I}_p)$$

- ▶ log likelihood (integrated) for  $\beta$  plus prior

$$-\frac{\phi}{2} (\|\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta\|^2 + k\|\beta\|^2)$$

- ▶ Posterior mean

$$\mathbf{b}_n = (\mathbf{X}^T \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \mathbf{X} \hat{\beta}$$

- ▶ importance of standardizing
- ▶ Choice of  $k$  in practice?
- ▶  $k = 0$  OLS
- ▶  $k = \infty$  estimates are  $\mathbf{0}$  (intercept only)

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular **X**

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular  $\mathbf{X}$
- ▶ Control how large coefficients may grow

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular  $\mathbf{X}$
- ▶ Control how large coefficients may grow

$$\min_{\beta} (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)^T (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)$$

subject to

$$\sum \beta_j^2 \leq t$$

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular  $\mathbf{X}$
- ▶ Control how large coefficients may grow

$$\min_{\beta} (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)^T (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)$$

subject to

$$\sum \beta_j^2 \leq t$$

- ▶ Equivalent Quadratic Programming Problem

$$\min_{\beta} \|\mathbf{Y}^c - \mathbf{X}^c\beta\|^2 + k\|\beta\|^2$$

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular  $\mathbf{X}$
- ▶ Control how large coefficients may grow

$$\min_{\beta} (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)^T (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)$$

subject to

$$\sum \beta_j^2 \leq t$$

- ▶ Equivalent Quadratic Programming Problem

$$\min_{\beta} \|\mathbf{Y}^c - \mathbf{X}^c\beta\|^2 + k\|\beta\|^2$$

- ▶ “penalized” likelihood

## Alternative Motivation

- ▶ If  $\hat{\beta}$  is unconstrained expect high variance with nearly singular  $\mathbf{X}$
- ▶ Control how large coefficients may grow

$$\min_{\beta} (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)^T (\mathbf{Y} - \mathbf{1}\bar{Y} - \mathbf{X}\beta)$$

subject to

$$\sum \beta_j^2 \leq t$$


- ▶ Equivalent Quadratic Programming Problem

$$\min_{\beta} \|\mathbf{Y}^c - \mathbf{X}^c\beta\|^2 + k\|\beta\|^2$$


- ▶ “penalized” likelihood
- ▶ Ridge Regression



# Geometry



Longley Data: `library(MASS); data(longley)`



# OLS

```
> longley.lm = lm(Employed ~ ., data=longley)
> summary(longley.lm)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	-3.482e+03	8.904e+02	-3.911	0.003560	**
GNP.deflator	1.506e-02	8.492e-02	0.177	0.863141	
GNP	-3.582e-02	3.349e-02	-1.070	0.312681	
Unemployed	-2.020e-02	4.884e-03	-4.136	0.002535	**
Armed.Forces	-1.033e-02	2.143e-03	-4.822	0.000944	***
Population	-5.110e-02	2.261e-01	-0.226	0.826212	
Year	1.829e+00	4.555e-01	4.016	0.003037	**

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.3049 on 9 degrees of freedom  
Multiple R-squared: 0.9955, Adjusted R-squared: 0.9925  
F-statistic: 330.3 on 6 and 9 DF, p-value: 4.984e-10

# Ridge Regression


```
# from library MASS
longley.ridge = lm.ridge(Employed ~ ., data=longley,
                        lambda=seq(0, 0.1, 0.0001))

# lambda = k in notes

summary(longley.ridge)
```

##	Length	Class	Mode
## coef	6006	-none-	numeric
## scales	6	-none-	numeric
## Inter	1	-none-	numeric
## lambda	1001	-none-	numeric
## ym	1	-none-	numeric
## xm	6	-none-	numeric
## GCV	1001	-none-	numeric
## kHKB	1	-none-	numeric
## kLW	1	-none-	numeric

# Ridge Trace Plot



## Choice of $k$


```
k = seq(0, 0.1, 0.0001)
n.k = length(k); n = nrow(longley)
cv.lambda = matrix(NA, n, n.k)

rmse.ridge = function(data, i, j, k) {
  m.ridge = lm.ridge(Employed ~ ., data = data, lambda=k[j],
                     subset = -i)
  yhat = scale(data[i,1:6, drop=F], center = m.ridge$xm,
               scale = m.ridge$scales) %*%
           m.ridge$coef + m.ridge$ym
  (yhat - data$Employed[i])^2
}

for (i in 1:n) {
  for (j in 1:n.k) {
    cv.lambda[i,j] = rmse.ridge(longley, i, j, k)
  }
}
```

# Cross Validation Error

```
cv.error = apply(cv.lambda, 2, mean)  
plot(k, cv.error, type="l")
```



Best  $k = 0.0028$

# Generalized Cross-validation

```
select(lm.ridge(Employed ~ ., data=longley,  
              lambda=seq(0, 0.1, 0.0001)))  
  
## modified HKB estimator is 0.004275357  
## modified L-W estimator is 0.03229531  
## smallest value of GCV   at 0.0028  
  
best.k = longley.ridge$lambda[which.min(longley.ridge$GCV)]  
longley.RReg = lm.ridge(Employed ~ ., data=longley,  
                       lambda=best.k)  
coef(longley.RReg)  
  
##                GNP.deflator                GNP      Unemployed  Arme  
## -2.950348e+03 -5.381450e-04 -1.822639e-02 -1.761107e-02 -9.60  
##      Population                Year  
## -1.185103e-01  1.557856e+00
```



## Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

# Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

- ▶  $p(\beta_0, \phi \mid \boldsymbol{\beta}, \kappa) \propto \phi^{-1}$

# Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

- ▶  $p(\beta_0, \phi \mid \boldsymbol{\beta}, \kappa) \propto \phi^{-1}$
- ▶  $\boldsymbol{\beta} \mid \phi, \kappa \sim \mathbf{N}(\mathbf{0}, \mathbf{I}(\phi\kappa)^{-1})$

# Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

- ▶  $p(\beta_0, \phi \mid \boldsymbol{\beta}, \kappa) \propto \phi^{-1}$
- ▶  $\boldsymbol{\beta} \mid \phi, \kappa \sim \mathbf{N}(\mathbf{0}, \mathbf{I}(\phi\kappa)^{-1})$
- ▶ prior on  $\kappa$ ?

# Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

- ▶  $p(\beta_0, \phi \mid \boldsymbol{\beta}, \kappa) \propto \phi^{-1}$
- ▶  $\boldsymbol{\beta} \mid \phi, \kappa \sim \mathbf{N}(\mathbf{0}, \mathbf{I}(\phi\kappa)^{-1})$
- ▶ prior on  $\kappa$ ?
- ▶ Take

$$\kappa \mid \phi \sim \text{Gamma}(1/2, 1/2)$$

# Priors on $k$

$\mathbf{X}$  is centered and standardized

$$\mathbf{Y} = \mathbf{1}\beta_0 + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Hierarchical prior

- ▶  $p(\beta_0, \phi \mid \boldsymbol{\beta}, \kappa) \propto \phi^{-1}$
- ▶  $\boldsymbol{\beta} \mid \phi, \kappa \sim \mathcal{N}(\mathbf{0}, \mathbf{I}(\phi\kappa)^{-1})$
- ▶ prior on  $\kappa$ ?
- ▶ Take

$$\kappa \mid \phi \sim \text{Gamma}(1/2, 1/2)$$

- ▶ What is induced prior on  $\boldsymbol{\beta} \mid \phi$ ?

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable



# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

Pick initial values  $\beta_0^{(0)}, \beta^{(0)}, \phi^{(0)}$ ,

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

Pick initial values  $\beta_0^{(0)}, \beta^{(0)}, \phi^{(0)}$ ,

Set  $t = 1$

1. Sample  $\kappa^{(t)} \sim p(\kappa \mid \beta_0^{(t-1)}, \beta^{(t-1)}, \phi^{(t-1)}, \mathbf{Y})$

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

Pick initial values  $\beta_0^{(0)}, \beta^{(0)}, \phi^{(0)}$ ,

Set  $t = 1$

1. Sample  $\kappa^{(t)} \sim p(\kappa \mid \beta_0^{(t-1)}, \beta^{(t-1)}, \phi^{(t-1)}, \mathbf{Y})$
2. Sample  $\beta_0^{(t)}, \beta^{(t)}, \phi^{(t)} \mid \kappa^{(t)}, \mathbf{Y}$

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

Pick initial values  $\beta_0^{(0)}, \beta^{(0)}, \phi^{(0)}$ ,

Set  $t = 1$

1. Sample  $\kappa^{(t)} \sim p(\kappa \mid \beta_0^{(t-1)}, \beta^{(t-1)}, \phi^{(t-1)}, \mathbf{Y})$
2. Sample  $\beta_0^{(t)}, \beta^{(t)}, \phi^{(t)} \mid \kappa^{(t)}, \mathbf{Y}$
3. Set  $t = t + 1$  and repeat until  $t > T$

# Posterior Distributions

## Joint Distribution

- ▶  $\beta_0, \beta, \phi \mid \kappa, \mathbf{Y}$  Normal-Gamma family given  $\mathbf{Y}$  and  $\kappa$
- ▶  $\kappa \mid \mathbf{Y}$  not tractable

Obtain marginal for  $\beta$  via MCMC

Pick initial values  $\beta_0^{(0)}, \beta^{(0)}, \phi^{(0)}$ ,

Set  $t = 1$

1. Sample  $\kappa^{(t)} \sim p(\kappa \mid \beta_0^{(t-1)}, \beta^{(t-1)}, \phi^{(t-1)}, \mathbf{Y})$
2. Sample  $\beta_0^{(t)}, \beta^{(t)}, \phi^{(t)} \mid \kappa^{(t)}, \mathbf{Y}$
3. Set  $t = t + 1$  and repeat until  $t > T$

Use Samples  $\beta_0^{(t)}, \beta^{(t)}, \phi^{(t)}, \kappa^{(t)}$  for  $t = B, \dots, T$  for inference

# JAGS

JAGS = Just Another Gibbs Sampler

# JAGS

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors



# JAGS

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors
- ▶ "derives" full conditional distributions

# JAGS

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors
- ▶ "derives" full conditional distributions
- ▶ integrates with R

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors
- ▶ "derives" full conditional distributions
- ▶ integrates with R
- ▶ typically faster than interpreted R code

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors
- ▶ "derives" full conditional distributions
- ▶ integrates with R
- ▶ typically faster than interpreted R code
- ▶ accounts for uncertainty about  $k$

JAGS = Just Another Gibbs Sampler

- ▶ scripting language to express sampling models and priors
- ▶ "derives" full conditional distributions
- ▶ integrates with R
- ▶ typically faster than interpreted R code
- ▶ accounts for uncertainty about  $k$

How would you compare Bayes predictions with Ridge with Cross-validation?