



Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

# Computational Strategies in Lévy Adaptive Regression Kernels

Merlise Clyde

Department of Statistical Science  
Duke University

BNPski Jan 9, 2014



# Outline

## Computational Strategies

M. Clyde

## Outline

### Motivation

### Nonparametric Regression

### Examples

### Summary

## Motivation

Mass Spectroscopy

Ovarian Cancer Risk Prediction

## Nonparametric Regression

Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation

Moving to Higher Dimensions

## Examples

## Summary



# MALDI-TOF Mass Spectroscopy

Computational  
Strategies

M. Clyde

Outline


Motivation

Mass  
Spectroscopy  
Ovarian Cancer  
Risk Prediction

Nonparametric  
Regression

Examples

Summary



Location of peaks/proteins in spectra in the presence of noise  
Non-Gaussian Noise



# Multiple Spectra

## Computational Strategies

M. Clyde

## Outline

## Motivation


## Mass Spectroscopy

Ovarian Cancer  
Risk Prediction

## Nonparametric Regression

## Examples

## Summary



Learning “features” that are common versus those that separate groups of spectra but do not know a priori the number of ions



# Predict Ovarian Cancer Risk

## Computational Strategies

M. Clyde

## Outline

## Motivation

Mass Spectroscopy  
Ovarian Cancer Risk Prediction

## Nonparametric Regression

## Examples

## Summary

- ▶ age
- ▶ family history of ovarian and breast cancer
- ▶ oral contraceptive use (duration)
- ▶ hormone replacement therapy
- ▶ hysterectomy
- ▶ number of pregnancies
- ▶ breastfeeding
- ▶ genetic single nucleotide polymorphisms (SNPs)

potential nonlinear functions and interactions?



# Problem Setting

## Computational Strategies

M. Clyde

## Outline

## Motivation

## Nonparametric Regression

Stochastic Expansions  
Lévy Random Field Priors - LARK  
Poisson Representation  
Moving to Higher Dimensions

## Examples

## Summary

## Regression problem

$$E[Y | \mathbf{x}] = f(\mathbf{x}), \quad \mathbf{x} \in \mathcal{X}$$

with unknown function  $f(\mathbf{x}) : \mathcal{X} \rightarrow \mathbb{R}$

Nonparametric Bayesian would place prior distribution on functions

- ▶ Gaussian Process Priors
- ▶ Dirichlet Process priors
- ▶ Lévy Processes

latter related to GP & DP and completely randomized measures



# Expansions

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK

Poisson  
Representation  
Moving to  
Higher  
Dimensions

Examples

Summary

## Expand

$$f(\mathbf{x}_i) = \sum_{j=0}^J \psi(\mathbf{x}_i, \omega_j) \beta_j$$

in terms of an (over-complete) dictionary where

- ▶  $\{\beta_j\}$ : unknown coefficients
- ▶  $J$ : number of terms in expansion (finite and infinite)
- ▶  $\psi(\mathbf{x}, \omega_j)$  Dictionary elements from a “generator function  $g$ ”
  - ▶ translation and scaling wavelet families
  - ▶ multivariate kernels (Gaussian, Cauchy, Exponential, e.g.)

$$g(\Lambda_j(\mathbf{x} - \chi_j)) = \exp \left\{ -\frac{1}{2}(\mathbf{x} - \chi_j)^T \Lambda_j(\mathbf{x} - \chi_j) \right\}$$

- ▶ Problem driven choices - Nualet, Barat & Rousseau
- ▶ Need not be symmetric!



# Kernel Convolution

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression





Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation  
Moving to Higher Dimensions

Examples

Summary



Easy to generate non-stationarity processes





# Lévy Adaptive Regression Kernels

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK

Poisson  
Representation  
Moving to  
Higher  
Dimensions

Examples

Summary

## Stochastic Integral Representation

$$f(\mathbf{x}) = \sum_{j \leq J} \psi(\mathbf{x}; \omega_j) \beta_j \equiv \int_{\Omega} \psi(\mathbf{x}; \omega) \mathcal{L}(d\omega)$$

$$\psi(\mathbf{x}, \omega_j) \equiv g(\Lambda_j(\mathbf{x} - \chi_j)) \quad \text{"generator"}$$

$\mathcal{L}$  is a Signed Measure (generalization of Completely Random Measures):

$$\mathcal{L}(d\omega) = \sum_{j \leq J} \beta_j \delta_{\omega_j}(d\omega)$$

- ▶ support points of measure  $\mathcal{L}$ :  $\{\omega_j\} = \{\chi_j, \lambda_j\}$ 
  - ▶ "location" parameters:  $\chi_j \in \mathcal{X}$
  - ▶ "scaling" parameters:  $\lambda_j \in \mathbb{R}^+$
- ▶ measure:  $\beta_j$
- ▶ number of support points  $J$



# Lévy Random Fields

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK

Poisson  
Representation

Moving to  
Higher  
Dimensions

Examples

Summary

- ▶  $\mathcal{L}(d\omega)$  is a **random (signed) measure** on  $\Omega$
- ▶ Convenient to think of a random measure as stochastic process where  $\mathcal{L}$  assigns random variables to sets  $A \in \Omega$
- ▶ Take

$$\mathcal{L} \sim \text{Lv}(\nu) \text{ with Lévy measure } \nu(d\beta, d\omega)$$

where  $\nu$  satisfies integrability condition:

$$\int_{\mathbb{R} \times \Omega} \min(1, \beta^2) \nu(d\beta, d\omega) < \infty \quad (1)$$

Choose  $\nu$  so the resulting  $f$  is well defined with desired smoothness properties or covariance function inherited from  $g$  (Wolpert, Clyde and Tu - 2011)



# Poisson Representation of Lévy Random Fields

## Computational Strategies

M. Clyde

## Outline

## Motivation

## Nonparametric Regression

## Stochastic Expansions

## Lévy Random Field Priors - LARK

## Poisson Representation

## Moving to Higher Dimensions

## Examples

## Summary

$$\begin{aligned} f(x) &= \int_{\Omega} \psi(x; \omega) \mathcal{L}(d\omega) \\ &= \int_{\mathbb{R} \times \Omega} \beta \psi(x; \omega) N(d\beta d\omega), \\ N(d\beta d\omega) &\sim P(\nu) \end{aligned}$$

when

$$\int_{\mathbb{R} \times \Omega} (1 \wedge |\beta|) \nu(d\beta, d\omega) < \infty$$

is satisfied and no need for “drift” term that arises with compensation under (1).

If  $\nu_+ \equiv \nu(\mathbb{R} \times \Theta) = \infty$ , must approximate to implement.



# Lévy- Khinchine Poisson Representation

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions  
Lévy Random  
Field Priors -  
LARK  
Poisson  
Representation  
Moving to  
Higher  
Dimensions

Examples

Summary

Goal:  $f(x) = \sum_{j < J} \psi(\mathbf{x}, \omega_j) \beta_j = \sum_{j < J} g(\Lambda_j(\mathbf{x} - \chi_j))$

**Sufficient condition** for bounded  $g$ :

$$\int_{\mathbb{R} \times \Omega} \min(1, |\beta|) \nu(d\beta, d\omega) < \infty \quad (2)$$

$$\Rightarrow J \sim P(\nu_+), \quad \nu_+ \equiv \nu(\mathbb{R} \times \Omega)$$

$$\Rightarrow \beta_j, \omega_j \mid J \stackrel{iid}{\sim} \pi(d\beta, d\omega) \propto \nu(d\beta, d\omega).$$

- ▶ Finite number of “big” coefficients  $|\beta_j|$
- ▶ Possibly infinite number of  $\beta \in [-\epsilon, \epsilon]$
- ▶ Coefficients  $|\beta_j|$  are absolutely summable<sup>1</sup>

---

<sup>1</sup>need to add a deterministic drift term to “compensate” for the infinite number of tiny jumps that are not absolutely summable under the more general integrability condition Equation (1)



# Existence Theorem

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK

Poisson  
Representation

Moving to  
Higher  
Dimensions

Examples

Summary

## Theorem

Let  $\nu$  be a Lévy measure on  $\mathbb{R} \times \Omega$  satisfying (1). Then  $f(\mathbf{x})$  is well-defined if  $\psi(\omega) \equiv g(\Lambda(\cdot - \chi))$  satisfies

$$\iint_{[-1,1]^c \times \Omega} (1 \wedge |\beta \psi(\omega)|) \nu(d\beta d\omega) < \infty \quad (3a)$$

$$\iint_{[-1,1] \times \Omega} (|\beta \psi(\omega)| \wedge |\beta \psi(\omega)|^2) \nu(d\beta d\omega) < \infty. \quad (3b)$$

For  $\nu$  satisfying Equation (2) the condition simplifies

$$\iint_{\mathbb{R} \times \Omega} (1 \wedge |\beta \phi(\omega)|) \nu(d\beta d\omega) < \infty \quad (3c)$$

$g$  is in a “Musielak-Orlicz space”



# $\alpha$ -Stable Lévy Measures

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions  
Lévy Random  
Field Priors -  
LARK

Poisson  
Representation  
Moving to  
Higher  
Dimensions

Examples

Summary

Lévy measure:  $\nu(d\beta, d\omega) = c_\alpha |\beta|^{-(\alpha+1)} \gamma(d\omega) \quad 0 < \alpha < 2$

For  $\alpha$ -Stable  $\nu^+(\mathbb{R}, \Omega) = \infty$

Fine in theory, but not in practice for MCMC!

Truncate measure to obtain a finite expansion:

- ▶ Finite number of support points  $\omega$  with  $\beta$  in  $[-\epsilon, \epsilon]^c$
- ▶ Fix  $\epsilon$  (for given prior approximation error)
- ▶ Use approximate Lévy measure

$$\nu_\epsilon(d\beta, d\omega) \equiv \nu(d\beta, d\omega) \mathbf{1}(|\beta| > \epsilon)$$

$$\Rightarrow J \sim P(\nu_\epsilon^+) \text{ where } \nu_\epsilon^+ = \nu([-\epsilon, \epsilon]^c, \Omega)$$

$$\Rightarrow \beta_j, \omega_j \stackrel{iid}{\sim} \pi(d\beta, d\omega) \equiv \nu_\epsilon(d\beta, d\omega) / \nu_\epsilon^+ \quad (\beta_j \text{ distributed as Pareto})$$

- ▶ need term for compensation for  $\alpha \geq 1$  for non-symmetric kernels



# Computation - Reversible Jump MCMC

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression

Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation

Moving to Higher Dimensions

Examples

Summary

- ▶ Birth: generate coefficients  $\beta_j$  near  $\epsilon$  in absolute value and generate kernel parameters  $\omega_j$  given increase in  $J \rightarrow J + 1$
- ▶ Death:  $\beta_j = 0$  drop dictionary element when  $J$  decrements by 1.
- ▶ Update: Random-Walk update, but also leads to deaths with coefficients that wander out of bounds and cross  $\epsilon$  boundary!
- ▶ Merge-Split: allow “neighboring points” to merge into one dictionary element (an alternative death) or a split into new dictionary elements

Advantage over fixed dimensional over-complete methods (frames)



# Wavelet Test Functions (SNR = 7)

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK


Poisson  
Representation

Moving to  
Higher  
Dimensions


Examples

Summary


blocks




bumps



doppler



heavysine







# Kernel Functions

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK


Poisson  
Representation

Moving to  
Higher  
Dimensions


Examples

Summary


**blocks**




**bumps**



**doppler**



**heavysine**





# Comparisons of OCD Methods

## Computational Strategies

M. Clyde

## Outline

## Motivation

## Nonparametric Regression

## Stochastic Expansions

## Lévy Random Field Priors - LARK

## Poisson Representation

## Moving to Higher Dimensions

## Examples

## Summary

- ▶ Translational Invariant Wavelets – Laplace Priors (Johnstone & Silverman 2005)
- ▶ Continuous Wavelet Dictionary – Compound Poisson with Gaussian Priors (Chu, Clyde, Liang 2007)
- ▶ LARK Symmetric Gamma
- ▶ LARK Cauchy

Range of Over-complete Dictionaries and Priors



# Comparison of Mean Square Error w/ OCDs

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK





Poisson  
Representation

Moving to  
Higher  
Dimensions

Examples

Summary

100 realizations of each function





# Motorcycle Crash Data: A Real Example

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK


Poisson  
Representation

Moving to  
Higher  
Dimensions

Examples

Summary

On average, only  $E[J \mid Y] \approx 4$  jumps are needed for fit:





# Form of Kernel

## Computational Strategies

M. Clyde

## Outline

## Motivation

## Nonparametric Regression

## Stochastic Expansions

## Lévy Random Field Priors - LARK


## Poisson Representation

## Moving to Higher Dimensions

## Examples

## Summary

$$k(t_i; \tau_j, \lambda_j) = e^{-\lambda_j |t_i - \tau_j|^\rho}$$





# Higher Dimensional $\mathcal{X}$

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression

Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation

Moving to Higher Dimensions

Examples

Summary

R-MCMC is too slow in higher dimensional space to allow

- ▶  $\chi$  to be completely arbitrary; restrict support to observed  $\{\mathbf{x}_i\}$
- ▶ use diagonal  $\Lambda$

Kernels take form:

$$\begin{aligned}\psi(\mathbf{x}, \omega_j) &= \prod_d \exp\left\{-\frac{1}{2}\lambda_d(x_d - \chi_d)^2\right\} \\ f(\mathbf{x}) &= \sum_j \psi(\mathbf{x}, \omega_j)\beta_j\end{aligned}$$



# Approximate Lévy Prior II

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions

Lévy Random  
Field Priors -  
LARK

Poisson  
Representation

Moving to  
Higher  
Dimensions

Examples

Summary

Continuous Approximation Student  $t(\alpha, 0, \epsilon)$  approximation:

$$\nu_\epsilon(d\beta, d\omega) = c_\alpha(\beta^2 + \alpha\epsilon^2)^{-(\alpha+1)/2} d\beta \gamma(d\omega)$$

Based on the following hierarchical prior

$$\begin{aligned}\beta_j \mid \phi_j &\stackrel{\text{ind}}{\sim} \text{N}(0, \varphi_j^{-1}) \\ \phi_j &\stackrel{\text{ind}}{\sim} \text{G}\left(\frac{\alpha}{2}, \frac{\alpha\epsilon^2}{2}\right) \\ J &\sim \text{P}(\nu_\epsilon^+)\end{aligned}$$

$$\text{where } \nu_\epsilon^+ = \nu_\epsilon(\mathbb{R}, \boldsymbol{\Omega}) = \frac{\alpha^{1-\alpha/2} \Gamma(\alpha) \Gamma(\alpha/2)}{\epsilon^\alpha \pi^{1/2} \Gamma(\frac{\alpha+1}{2})} \sin(\frac{\pi\alpha}{2}) \gamma(\boldsymbol{\Omega})$$



# Limiting Case

## Computational Strategies

M. Clyde

## Outline

## Motivation

## Nonparametric Regression

Stochastic Expansions  
Lévy Random Field Priors - LARK  
Poisson Representation  
Moving to Higher Dimensions

## Examples

## Summary

$$\begin{aligned}\beta_j \mid \varphi_j &\stackrel{\text{ind}}{\sim} \mathcal{N}(0, 1/\varphi_j) \\ \varphi_j &\stackrel{\text{iid}}{\sim} \mathcal{G}(\alpha/2, "0")\end{aligned}$$

## Notes:

- ▶ Require  $0 < \alpha < 2$  for characteristic function for  $\mathcal{L}$  and functionals to exist. Additional restrictions on  $\omega$
- ▶ Cauchy process corresponds to  $\alpha = 1$
- ▶ Tipping's choice corresponds to  $\alpha = 0$
- ▶ Provides an extension of [Generalized Ridge Priors](#) to infinite dimensional
- ▶ Infinite dimensional analog of Cauchy priors





# Further Simplification in Case with $\alpha = 1$

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Stochastic  
Expansions  
Lévy Random  
Field Priors -  
LARK

Poisson  
Representation  
Moving to  
Higher  
Dimensions

Examples

Summary

- Poisson number of points  $J_\epsilon \sim P(\nu_\epsilon^+(\alpha, \gamma))$  with

$$\nu_\epsilon^+(\alpha, \gamma) = \frac{\gamma \alpha^{1-\alpha/2}}{2^{1-\alpha} \epsilon^\alpha} \frac{\Gamma(\alpha/2)}{\Gamma(1-\alpha/2)}$$

- Given  $J, [n_1 : n_n] \sim MN(J, 1/(n+1))$  points supported at each kernel located at  $\mathbf{x}_j$

The regression mean function can be rewritten as

$$f(\mathbf{x}) = \sum_{i=0}^n \tilde{\beta}_i k(\mathbf{x}, \omega_i), \quad \tilde{\beta}_i = \sum_{\{j | \mathbf{x}_j = \mathbf{x}_i\}} \beta_j.$$

In particular, if  $\alpha = 1$ , not only the Cauchy process is infinitely divisible, the approximated Cauchy prior distributions on the regression coefficients are also infinitely divisible:

$$\tilde{\beta}_i \stackrel{iid}{\sim} N(0, n_i^2 \tilde{\varphi}_i^{-1}), \quad \tilde{\varphi}_i \stackrel{iid}{\sim} G(1/2, \epsilon^2/2)$$

At most  $n$  non-zero coefficients!



# Collapsed Sampler

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression

Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation

Moving to Higher Dimensions

Examples

Summary

**Advantage:** Gaussian prior so  $\beta$  can be integrated out for MCMC under Gaussian error model in the  $n$  dimensional problem

- ▶ Integrate out  $\beta$  vector in Normal regression leaving kernel parameters  $\Lambda$  and  $\varphi$  with multinomial weights  $n_i$
- ▶  $n_i$  may be 0 which drops dictionary elements from representation in finite representation for fixed  $\epsilon$  - Holmes approximate models?
- ▶ still trans-dimensional in  $\varphi$  but much simpler RJMCMC now



# Feature Selection in Kernel

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression

Stochastic Expansions

Lévy Random Field Priors - LARK

Poisson Representation

Moving to Higher Dimensions

Examples

Summary

- ▶ Product structure allows interactions between variables
- ▶ Many input variables may be irrelevant
- ▶ Feature selection; if  $\lambda_d = 0$  variable  $x_d$  is removed from all kernels
- ▶ Allow point mass on  $\lambda_h = 0$  with probability  $p_\lambda \sim B(a, b)$  (in practice have used  $a = b = 1$ )

## Consider 3 Scenarios

- ▶  $D$  Different  $\lambda$  –  $D$  parameters in each dimension
- ▶  $S + D$  Different  $\lambda_d$  parameters + Selection
- ▶  $S + E$  Selection + Equal for Remaining  $\lambda_d = \lambda$



# Regression Out of Sample Prediction

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

## Average Relative MSE to best procedure

Data Sets	BARK			SVM	BART
	D	S + E	S + D		
Friedman1	1.22	2.26	1.93	5.36	1.97
Friedman2	1.07	1.09	1.04	4.36	3.64
Friedman3	1.46	2.30	1.44	2.70	1.00
Boston Housing	1.09	1.23	1.20	1.56	1.01
Body Fat	1.81	1.01	2.19	4.04	1.68
Basketball	1.01	1.01	1.02	1.16	1.10

D: dimension specific scale  $\lambda_d$

E: equal scales  $\lambda_d = \lambda \forall d$

S: selection  $\lambda_d = 0$  with probability  $\rho$



# Feature Selection in Boston Housing Data

Computational  
Strategies

M. Clyde

Outline

Motivation


Nonparametric  
Regression

Examples

Summary

## Posterior Distribution of $\lambda_d$

**Boston Housing in BARK with different weights**





# Classification Examples

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

Name	$d$	data type	$n$ (train/test)
Circle	2	simulation	200/1000
Circle (3 null)	5	simulation	200/1000
Circle (18 null)	20	simulation	200/1000
Swiss Bank Notes	6	real data	200 (5 cv)
Breast Cancer	30	real data	569 (5 cv)
Ionosphere	33	real data	351 (5 cv)

- ▶ Add latent Gaussian  $Z_i$  for probit regression (as in Albert & Chib)
- ▶ Same model as before conditional on  $\mathbf{Z}$
- ▶ Advantage: Draw  $\beta$  in a block from full conditional
- ▶ Can extend to Logistic



# Predictive Error Rate for Classification

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

Data Sets	BARK			SVM	BART
	D	S + E	S + D		
Circle 2	4.91%	1.88%	1.93%	5.03%	3.97%
Circle 5	4.70%	1.47%	1.65%	10.99%	6.51%
Circle 20	4.84%	2.09%	3.69%	44.10%	15.10%
Bank	1.25%	0.55%	0.88%	1.12%	0.50%
BC	4.02%	2.49%	6.09%	2.70%	3.36%
Ionosphere	8.59%	5.78%	10.87%	5.17%	7.34%

D: dimension specific scale  $\lambda_d$

E: equal scales  $\lambda_d = \lambda \forall d$

S: selection  $\lambda_d = 0$  with probability  $\rho$



# Needs & Limitations

## Computational Strategies

M. Clyde

Outline

Motivation

Nonparametric Regression

Examples

Summary

- ▶ NP Bayes of many flavors often does better than frequentist methods (LARK, BART, Treed GP, more)
- ▶ Hyper-parameter specification - theory & computational approximation
- ▶ need faster code that is easier for users (BART & TGP are great!)
- ▶ Can these models be added to JAGS, STAN, etc instead of stand-alone R packages
- ▶ With availability of code what are caveats for users?





# Summary

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

## Lévy Random Field Priors & LARK models:

- ▶ Provide limit of finite dimensional priors (GRP & SVSS) to infinite dimensional setting
- ▶ Adaptive bandwidth for kernel regression
- ▶ Allow flexible generating functions
- ▶ Provide sparser representations compared to SVM & RVM, with coherent Bayesian interpretation
- ▶ Incorporation of prior knowledge if available
- ▶ Relax assumptions of equally spaced data and Gaussian likelihood
- ▶ Hierarchical Extensions
- ▶ Formulation allows one to define stochastic processes on arbitrary spaces (spheres, manifolds)



# Thanks!

Computational  
Strategies

M. Clyde

Outline

Motivation

Nonparametric  
Regression

Examples

Summary

This would not be possible without coauthors:

- ▶ Robert Wolpert
- ▶ Leanna House
- ▶ Chong Tu
- ▶ Zhi Ouyang

and support from the National Science Foundation

For papers see <http://stat.duke.edu/~clyde>