


# Drought Index Comparison Portal

*A tool to help compare the outputs of various commonly used drought indices and aid in the exploration of historical drought in the Contiguous United States.*

## Why

Drought can represent various forms of water shortage and is often categorized according to the type of Earth or social system it affects (hydrological, meteorological, agricultural, etc.). In turn, there are many different indices designed to reflect different forms of drought. There is no

Study Period Year Range: 1985 - 2019



Map Type

Dark × ▾


Function

Percentiles Index Values  
Mean × ▾



Color Gradient

Reversed Not Reversed  
Default × ▾


SUBMIT OPTIONS



Palmer Drought Severity Index  
Moffat County, CO



Standardized Precipitation-Evapotranspiration Index - 1 month  
Moffat County, CO



**Figure 1.** A comparison of the Palmer Drought Severity Index and the 1-month Standardized Precipitation-Evapotranspiration Index as average values across the Contiguous United States and as a time-series of monthly values in Routt County, Colorado.


singular index that can be used for all applications and, even within one application, indices will vary across space in terms of performance.

It has been suggested by the research community to apply a multi-index approach to complex water management problems, or at least to explore multiple indices before basing decisions on any singular index. Such an exploration can be tedious without extensive efforts to acquire, calculate, and manipulate index data. There are some services that automate the process, provide data for a suite of indices, and allow for comparisons such as this (see the [Drought Risk Atlas](#) of the [NDMC](#), the [US Water Watcher](#), or the [WestWide Drought Tracker](#)). However, while these are invaluable resources, they do not provide the basic ability to quickly compare both spatial and temporal patterns of drought according to different indices. Therefore, this online tool was created to provide a singular portal in which to perform quick visual comparisons across time and space between a collection of drought index products.

## What


The application currently displays two side-by-side panels of drought indices over the Contiguous United States. For now, there are variations on four common drought indices available: The Palmer Drought Severity Index (PDSI), the Standardized Precipitation Index (SPI), the Standardized Precipitation-Evapotranspiration Index (SPEI), and the Evaporative Demand Drought Index (EDDI). The SPI, SPEI, and EDDI are multi-scalar indices for which 1-, 2-, 3-, and 6-month time periods are available. In addition to the PDSI, the Self-Calibrating PDSI and Palmer Z Index are also available, bringing the total number of available indices to fifteen.

For each panel, you may select any index from above to display a map of values along with a study area and time period, as demonstrated in **Figure 1**. Values may be presented as mean, min, or maximum index or percentile values. When using the original index values, information may also be presented as the percent of the selected area in various drought categories. For most indices, excepting EDDI, the study period can include any number of months since 1900. EDDI is only



**Figure 2.** The spatial extent of drought in the 1930's, as exhibited by average SPEI and SPI values at the 6-month time-scale, with an example time series in Sanborn County, South Dakota.

available since 1980 because it uses a more sophisticated method of calculating evapotranspiration and requires data that was not available at earlier times.



**Figure 3.** The “RdWhBu (NOAA PSD scale)” color scheme allows for the identification of categorical drought events. It also incorporates a slight gradient within each category to allow for further differentiation of severity. This option works best for a single month because colors quickly converge on white color as values are averaged over longer time periods.

## Colors

We feel that it is important to provide as many perspectives of the drought as possible, and that the color schemes used to display this information affects the perception of the phenomenon. Therefore, eighteen different color scales are available, along with the ability to reverse scales for each. Defaults are set to a red, white, and blue scale for percentiles and a brown, white, green scale for original index values. The “NOAA PSD scale” is based on the National Oceanic and Atmospheric Administration Physical Science Division’s

percentile color scale used for EDDI and highlights particularly wet and dry periods. An example of this scale is displayed in **Figure 3**. The color scales are kept relative to the chosen time period because they quickly homogenize over time as values approach the mean (particularly for percentiles).


Data was acquired from the WestWide Drought Tracker of the Desert Research Institute and [NOAA's Physical Sciences Division](#).

## How

To generate outputs, select a function to apply to the chosen index, adjust the time period over which the function is calculated, and change the base map and color scale. Each point in the map represents the southwest corner of a 0.25 by 0.25 decimal degree grid cell. Hovering over these points shows the index value and county of each location. Hovering over the map and scrolling allows you to zoom and search the base map for a particular location. Clicking and dragging the mouse allows you to pan the map. Holding control, clicking, and dragging the mouse allows you to change the viewing aspect.

Clicking on a point renders a time series of monthly index values as a bar graph. If you know the county of a desired location you may select it with a drop-down menu, though this will only select roughly 15 square miles of the northwest portion of that county. To select more specific locations, scroll the mouse and zoom into the base map. Because grid cells are represented as points, it is possible to zoom in beyond them, view geographic labels, and click the closest point.


It is also possible to choose multiple grid cells with the box or lasso selection tool. Doing so will render a time series of monthly values averaged within the selection. By clicking and dragging the box selection tool directly up or down (or left or right) you can select a range of latitudes (or



**Figure 4.** The Drought Comparison Portal allows for a high degree of flexibility when selecting areas over which to aggregate various metrics of drought. Displayed here are averages from the selection of a range of longitudes and latitudes of PDSI percentile values from 1985 to 2019.

longitudes (**Fig. 4** ).

The resulting time series is highly manipulatable, as per the functionality of Plotly's DASH module. Using the toolbar at the top right of the graph, the user may change the y-scale, change the x-scale, zoom into any segment, and pan along either axis for an up-close view of the




**Figure 5.** Time series values may be highlighted using the box selection tool, which is particularly useful for identifying patterns such as those buried within short-term index data.

information. Double clicking on the graph resets the view or expand the range to the extent of the panel. The user may also use the select tool to highlight bar values above or below chosen y- or x-axis values (**Fig. 5**).

## Identifying extreme events

Certain color scales combined with the minimum value (maximum for EDDI) option make it simple to find where and when extreme drought events/anomalies occurred. Because color schemes are relative to each map, these extreme values contrast highly with those around them, allowing the user to select a location, generate a time series, zoom into the general time period surrounding the event and hover over the bar graph to find the exact month and year, as is demonstrated in **Figure 6**.



**Figure 6.** Examples of using the minimum value option to identify extreme events. In this case, the “RdYlGnBu” color scale creates a field of blue values with pockets of orange/red if extreme events occurred at any point within the study period.

## The “Drought Severity Area” option


You may calculate the percentage of any selected area that was in drought according to classifications found on the United States Drought Monitor (USDM) [data information page](#). This is done by projecting the selected coordinates to the North American Albers Equal Area Conic coordinate projection system, filtering the data for values within the specified categorical ranges, and calculating the percentage of grid cells within each category. The information generated from this process is not the same as that shown in the USDM, but instead represents analogous estimations of drought severity. **Table 1** shows the categorical thresholds used for each index.



Palmer index are each set according to that of the PDSI, and thresholds for the EDDI, SPEI, and SPI are set to that of the SPI (the original multi-scalar index). The percentages displayed are inclusive of overlapping categories such that an area in the “severe” drought category is also counted in the “moderate” and “dry” categories.

**Table 1.** The drought severity categories for various indices used in the Drought Comparison Portal. Derived from the drought classifications used by the National Drought Mitigation Center.

Category	Description	Palmer Indices	SPI, SPEI, & EDDI
D0	Abnormally Dry	-1.0 to -2.0	-0.5 to -0.8
D1	Moderate Drought	-2.0 to -3.0	-0.8 to -1.3
D2	Severe Drought	-3.0 to -4.0	-1.3 to -1.6
D3	Extreme Drought	-4.0 to -5.0	-1.6 to -2.0
D4	Exceptional Drought	less than -5.0	less than -2.0




**Figure 7.** Percent of area in drought according to the Palmer Drought Severity Index and the Standardized Precipitation-Evapotranspiration Index in Montana, South Dakota, and North Dakota. 1895 to 2019.

The Drought Severity Coverage Index (DSCI), which is described by the NDMC [here](#), is also available. The DSCI takes each drought category into account by weighting the percentage of area covered by each according to their respective levels of severity, as described below. This value is not calculated with inclusive categories.

$$DSCI = 1*(\%D0) + 2*(\%D2) + 3*(\%D2) + 4*(\%D3) + 5*(\%D4)$$

Hovering over this graph will update monthly values of percent coverage for each category in a box at the bottom of the page.



**Figure 8.** A line representing the Drought Severity Coverage Index (DSCI) may be toggled on and off to display a singular metric of drought severity by area.

## Downloading Information

There are currently three ways of downloading the information you generate with this application:

- 1) Hover over the top right corner of a graph and click the camera icon in the toolbar to download the image as a portable network graphic (png) file.
- 2) Click the computer disk icon to edit the chart in Plotly's chart studio. Visit this page for tutorials on how to use this service: <https://help.plot.ly/tutorials/#fundamentals>.
- 3) Click the "Download Time Series Data" link at the bottom of each time series graph to download a csv of the data underlying that graph.

This tool is available at <https://climate-scatterplot.space>. It is in development and we appreciate your comments. Feel free to report any bugs or glitches:

Travis Williams: [travis.williams@colorado.edu](mailto:travis.williams@colorado.edu)

William Travis: [william.travis@colorado.edu](mailto:william.travis@colorado.edu)