

Calculus Cheat Sheet

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1.$$

$$\frac{dx^n}{dx} = nx^{n-1}.$$

$$\frac{d \ln(x)}{dx} = \frac{1}{x}.$$

$$\frac{da^x}{dx} = \ln(a)a^x.$$

$$\frac{d \sin(x)}{dx} = \cos(x).$$

$$\frac{d \cos(x)}{dx} = -\sin(x).$$

$$\frac{d \tan(x)}{dx} = \sec^2(x).$$

$$\frac{d \cot(x)}{dx} = -\csc^2(x).$$

$$\frac{d \sec(x)}{dx} = \tan(x) \sec(x).$$

$$\frac{d \csc(x)}{dx} = -\cot(x) \csc(x).$$

$$\frac{d \arcsin(x)}{dx} = \frac{1}{\sqrt{1-x^2}}.$$

$$\frac{d \arccos(x)}{dx} = -\frac{1}{\sqrt{1-x^2}}.$$

$$\frac{d \arctan(x)}{dx} = \frac{1}{1+x^2}.$$

$$\frac{d \operatorname{arccot}(x)}{dx} = -\frac{1}{1+x^2}.$$

$$\frac{d \operatorname{arcsec}(x)}{dx} = \frac{1}{|x| \sqrt{x^2-1}}.$$

$$\frac{d \operatorname{arccsc}(x)}{dx} = -\frac{1}{|x| \sqrt{x^2-1}}.$$

$$\frac{d \sinh(x)}{dx} = \cosh(x).$$

$$\frac{d \cosh(x)}{dx} = \sinh(x).$$

$$\frac{d \tanh(x)}{dx} = \operatorname{sech}^2(x).$$

$$\frac{d \coth(x)}{dx} = -\operatorname{csch}^2(x), \quad x \neq 0.$$

$$\frac{d \operatorname{sech}(x)}{dx} = -\tanh(x) \operatorname{sech}(x).$$

$$\frac{d \operatorname{csch}(x)}{dx} = -\coth(x) \operatorname{csch}(x), \quad x \neq 0.$$

$$\frac{d \operatorname{arcsinh}(x)}{dx} = \frac{1}{\sqrt{x^2 + 1}}.$$

$$\frac{d \operatorname{arccosh}(x)}{dx} = \frac{1}{\sqrt{x^2 - 1}}, \quad x > 1.$$

$$\frac{d \operatorname{arctanh}(x)}{dx} = \frac{1}{1 - x^2}, \quad |x| < 1.$$

$$\frac{d \operatorname{arcoth}(x)}{dx} = \frac{1}{1 - x^2}, \quad |x| > 1.$$

$$\frac{d \operatorname{arcsech}(x)}{dx} = -\frac{1}{x\sqrt{1 - x^2}}, \quad 0 < x < 1.$$

$$\frac{d \operatorname{arccsch}(x)}{dx} = -\frac{1}{|x|\sqrt{1 + x^2}}, \quad x \neq 0.$$

$$\int x^n dx = \begin{cases} \frac{1}{n+1} x^{n+1} + C, & n \neq -1 \\ \ln |x| + C, & n = -1 \end{cases}.$$

$$\int \ln(x) dx = x \ln(x) - x + C.$$

$$\int (\ln(x))^n dx = x (\ln(x))^n - n \int (\ln(x))^{n-1} dx, \quad n \in \mathbb{N}.$$

$$\int e^{nx} dx = \frac{1}{n} e^{nx} + C, \quad n \neq 0.$$

$$\int \sin(x) dx = -\cos(x) + C.$$

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(2x)}{4} + C.$$

$$\int \sin^n(x) dx = -\frac{1}{n} \sin^{n-1}(x) \cos(x) + \frac{n-1}{n} \int \sin^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cos(x) dx = \sin(x) + C.$$

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(2x)}{4} + C.$$

$$\int \cos^n(x) dx = \frac{1}{n} \cos^{n-1}(x) \sin(x) + \frac{n-1}{n} \int \cos^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \tan(x) dx = -\ln |\cos(x)| + C = \ln |\sec(x)| + C.$$

$$\int \tan^2(x) dx = \tan(x) - x + C.$$

$$\int \tan^n(x) dx = \frac{1}{n-1} \tan^{n-1}(x) - \int \tan^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cot(x) dx = \ln |\sin(x)| + C.$$

$$\int \cot^2(x) dx = -\cot(x) - x + C.$$

$$\int \cot^n(x) dx = -\frac{1}{n-1} \cot^{n-1}(x) - \int \cot^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \sec(x) \, dx = \ln |\sec(x) + \tan(x)| + C.$$

$$\int \sec^2(x) \, dx = \tan(x) + C.$$

$$\int \sec^n(x) \, dx = \frac{1}{n-1} \sec^{n-2}(x) \tan(x) + \frac{n-2}{n-1} \int \sec^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \csc(x) \, dx = -\ln |\csc(x) + \cot(x)| + C.$$

$$\int \csc^2(x) \, dx = -\cot(x) + C.$$

$$\int \csc^n(x) \, dx = -\frac{1}{n-1} \csc^{n-2}(x) \cot(x) + \frac{n-2}{n-1} \int \csc^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \arcsin(x) \, dx = x \arcsin(x) + \sqrt{1-x^2} + C.$$

$$\int \arccos(x) \, dx = x \arccos(x) - \sqrt{1-x^2} + C.$$

$$\int \arctan(x) \, dx = x \arctan(x) - \frac{1}{2} \ln(1+x^2) + C.$$

$$\int \operatorname{arccot}(x) \, dx = x \operatorname{arccot}(x) + \frac{1}{2} \ln(1+x^2) + C.$$

$$\int \operatorname{arcsec}(x) \, dx = x \operatorname{arcsec}(x) - \operatorname{sgn}(x) \ln \left| x + \sqrt{x^2-1} \right| + C, \quad |x| \geq 1.$$

$$\int \operatorname{arccsc}(x) \, dx = x \operatorname{arccsc}(x) + \operatorname{sgn}(x) \ln \left| x + \sqrt{x^2-1} \right| + C, \quad |x| \geq 1.$$

$$\int \sinh(x) \, dx = \cosh(x) + C.$$

$$\int \sinh^2(x) \, dx = \frac{\sinh(2x)}{4} - \frac{x}{2} + C.$$

$$\int \sinh^n(x) \, dx = \frac{1}{n} \sinh^{n-1}(x) \cosh(x) - \frac{n-1}{n} \int \sinh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cosh(x) \, dx = \sinh(x) + C.$$

$$\int \cosh^2(x) \, dx = \frac{\sinh(2x)}{4} + \frac{x}{2} + C.$$

$$\int \cosh^n(x) \, dx = \frac{1}{n} \cosh^{n-1}(x) \sinh(x) + \frac{n-1}{n} \int \cosh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \tanh(x) \, dx = \ln \cosh(x) + C.$$

$$\int \tanh^2(x) \, dx = x - \tanh(x) + C.$$

$$\int \tanh^n(x) \, dx = \frac{1}{n-1} \tanh^{n-1}(x) - \int \tanh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \coth(x) \, dx = \ln |\sinh(x)| + C.$$

$$\int \coth^2(x) dx = x - \coth(x) + C.$$

$$\int \coth^n(x) dx = -\frac{1}{n-1} \coth^{n-1}(x) - \int \coth^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{sech}(x) dx = \operatorname{arctanh} \sinh(x) + C.$$

$$\int \operatorname{sech}^2(x) dx = \tanh(x) + C.$$

$$\int \operatorname{sech}^n(x) dx = \frac{1}{n-1} \operatorname{sech}^{n-1}(x) \tanh(x) + \frac{n-2}{n-1} \int \operatorname{sech}^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{csch}(x) dx = \ln \left| \tanh \frac{x}{2} \right| + C.$$

$$\int \operatorname{csch}^2(x) dx = -\coth(x) + C.$$

$$\int \operatorname{csch}^n(x) dx = -\frac{1}{n-1} \operatorname{csch}^{n-1}(x) \coth(x) + \frac{n-2}{n-1} \int \operatorname{csch}^{n-2}(x) dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{arcsinh}(x) dx = x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1} + C.$$

$$\int \operatorname{arccosh}(x) dx = x \operatorname{arccosh}(x) - \sqrt{x^2 - 1} + C, \quad (x) \geq 1.$$

$$\int \operatorname{arctanh}(x) dx = x \operatorname{arctanh}(x) + \frac{1}{2} \ln(1 - x^2) + C, \quad |x| < 1.$$

$$\int \operatorname{arccoth}(x) dx = x \operatorname{arccoth}(x) + \frac{1}{2} \ln(x^2 - 1) + C, \quad |x| > 1.$$

$$\int \operatorname{arcsech}(x) dx = x \operatorname{arcsech}(x) + \arctan \left(\frac{\sqrt{1-x^2}}{x} \right) + C, \quad 0 < x \leq 1.$$

$$\int \operatorname{arccsch}(x) dx = x \operatorname{arccsch}(x) + \ln \left| x + \sqrt{x^2 + 1} \right| + C, \quad x \neq 0.$$

$$\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \arctan \frac{x}{a} + C, \quad a \neq 0.$$

$$\int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \ln \left(\frac{x-a}{x+a} \right) + C, \quad a \neq 0.$$

$$\int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \ln \left(\frac{a-x}{a+x} \right) + C, \quad a \neq 0.$$

$$\int \frac{1}{ax + b} dx = \frac{1}{a} \ln |ax + b| + C, \quad a \neq 0.$$

$$\int (ax + b)^n dx = \frac{1}{a(n+1)} (ax + b)^{n+1} + C, \quad a \neq 0 \wedge n \neq -1.$$

$$\int \frac{x}{ax + b} dx = \frac{x}{a} - \frac{b}{a^2} \ln |ax + b| + C, \quad a \neq 0.$$

$$\int \frac{x}{(ax + b)^2} dx = \frac{1}{a^2} \ln |ax + b| + \frac{b}{a^2(ax + b)} + C, \quad a \neq 0.$$

$$\int x(ax + b)^n dx = \frac{a(n+1)x - b}{a^2(n+1)(n+2)} (ax + b)^{n+1} + C, \quad a \neq 0 \wedge n \notin \{-1, -2\}.$$

$$\int \sqrt{a^2 + x^2} \, dx = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \ln(x + \sqrt{a^2 + x^2}) + C.$$

$$\int x \sqrt{a^2 + x^2} \, dx = \frac{1}{3} (a^2 + x^2)^{3/2} + C.$$

$$\int \frac{1}{\sqrt{a^2 + x^2}} \, dx = \operatorname{arcsinh} \frac{x}{|a|} + C = \ln(x + \sqrt{a^2 + x^2}) + C.$$

$$\int \frac{x}{\sqrt{a^2 + x^2}} \, dx = \sqrt{a^2 + x^2} + C.$$

$$\int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \ln(x + \sqrt{x^2 - a^2}) + C, \quad x^2 > a^2.$$

$$\int x \sqrt{x^2 - a^2} \, dx = \frac{1}{3} (x^2 - a^2)^{3/2} + C, \quad x^2 > a^2.$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \arcsin\left(\frac{x}{|a|}\right) + C, \quad x^2 > a^2.$$

$$\int \frac{x}{\sqrt{a^2 - x^2}} \, dx = -\sqrt{a^2 - x^2} + C, \quad x^2 > a^2.$$

$$\int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{|a|} + C, \quad a^2 \geq x^2.$$

$$\int x \sqrt{a^2 - x^2} \, dx = \frac{1}{3} (a^2 - x^2)^{3/2} + C, \quad a^2 \geq x^2.$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \arcsin\left(\frac{x}{|a|}\right) + C, \quad a^2 \geq x^2.$$

$$\int \frac{x}{\sqrt{a^2 - x^2}} \, dx = \sqrt{a^2 - x^2} + C, \quad a^2 \geq x^2.$$

$$\int \sqrt{ax + b} \, dx = \frac{2}{3a} (ax + b)^{3/2} + C, \quad a \neq 0.$$

$$\int x^n \sqrt{ax + b} \, dx = \frac{2}{a(2n+3)} \left(x^n (ax + b)^{3/2} - bn \int x^{n-1} \sqrt{ax + b} \, dx \right), \quad a \neq 0 \wedge n \in \mathbb{N}.$$

$$\int \frac{1}{\sqrt{ax + b}} \, dx = \frac{2\sqrt{ax + b}}{a} + C, \quad a \neq 0.$$

$$\int \frac{x^n}{\sqrt{ax + b}} \, dx = \frac{2}{a} \left(x^n \sqrt{ax + b} - n \int x^{n-1} \sqrt{ax + b} \, dx \right), \quad a \neq 0 \wedge n \in \mathbb{N},$$

$$= \frac{2}{a(2n+1)} \left(x^n \sqrt{ax + b} - bn \int \frac{x^{n-1}}{\sqrt{ax + b}} \, dx \right).$$

$$\int_{-\infty}^{\infty} e^{-t^2} \, dt = \sqrt{\pi}.$$

$$\int_0^{\infty} e^{-t^2} \, dt = \frac{\sqrt{\pi}}{2}.$$