

Calculus Cheat Sheet

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1.$$

$$\frac{dx^n}{dx} = nx^{n-1}.$$

$$\frac{d \ln(x)}{dx} = \frac{1}{x}.$$

$$\frac{da^x}{dx} = \ln(a)a^x.$$

$$\frac{d \sin(x)}{dx} = \cos(x).$$

$$\frac{d \cos(x)}{dx} = -\sin(x).$$

$$\frac{d \tan(x)}{dx} = \sec^2(x).$$

$$\frac{d \cot(x)}{dx} = -\csc^2(x).$$

$$\frac{d \sec(x)}{dx} = \tan(x) \sec(x).$$

$$\frac{d \csc(x)}{dx} = -\cot(x) \csc(x).$$

$$\frac{d \arcsin(x)}{dx} = \frac{1}{\sqrt{1-x^2}}.$$

$$\frac{d \arccos(x)}{dx} = -\frac{1}{\sqrt{1-x^2}}.$$

$$\frac{d \arctan(x)}{dx} = \frac{1}{1+x^2}.$$

$$\frac{d \operatorname{arccot}(x)}{dx} = -\frac{1}{1+x^2}.$$

$$\frac{d \operatorname{arcsec}(x)}{dx} = \frac{1}{|x| \sqrt{x^2-1}}.$$

$$\frac{d \operatorname{arccsc}(x)}{dx} = -\frac{1}{|x| \sqrt{x^2-1}}.$$

$$\frac{d \sinh(x)}{dx} = \cosh(x).$$

$$\frac{d \cosh(x)}{dx} = \sinh(x).$$

$$\frac{d \tanh(x)}{dx} = \operatorname{sech}^2(x).$$

$$\frac{d \coth(x)}{dx} = -\operatorname{csch}^2(x).$$

$$\frac{d \operatorname{sech}(x)}{dx} = -\tanh(x) \operatorname{sech}(x).$$

$$\frac{d \operatorname{csch}(x)}{dx} = -\coth(x) \operatorname{csch}(x).$$

$$\frac{\mathrm{d} \operatorname{arcsinh}(x)}{\mathrm{d} x} = \frac{1}{\sqrt{x^2 + 1}}.$$

$$\frac{\mathrm{d} \operatorname{arccosh}(x)}{\mathrm{d} x} = \frac{1}{\sqrt{x^2 - 1}}, \quad x > 1.$$

$$\frac{\mathrm{d} \operatorname{arctanh}(x)}{\mathrm{d} x} = \frac{1}{1 - x^2}, \quad |x| < 1.$$

$$\frac{\mathrm{d} \operatorname{arcoth}(x)}{\mathrm{d} x} = \frac{1}{1 - x^2}, \quad |x| > 1.$$

$$\frac{\mathrm{d} \operatorname{arcsech}(x)}{\mathrm{d} x} = -\frac{1}{x\sqrt{1 - x^2}}, \quad 0 < x \leq 1.$$

$$\frac{\mathrm{d} \operatorname{arccsch}(x)}{\mathrm{d} x} = -\frac{1}{|x|\sqrt{1 + x^2}}.$$

$$x^n = \sum_{k=0}^{\infty} \binom{n}{k} a^{n-k} (x - a)^k, \quad |x - a| < |a|.$$

$$\ln(x) = \ln(a) + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(x - a)^n}{na^n}, \quad 0 < x \leq a.$$

$$a^x = \sum_{n=0}^{\infty} \frac{a^b (\ln(a))^n}{n!} (x - b)^n, \quad a \in \mathbb{R}_{>0}.$$

$$\sin(x) = \sum_{n=0}^{\infty} \frac{\sin\left(a + \frac{n\pi}{2}\right)}{n!} (x - a)^n.$$

$$\cos(x) = \sum_{n=0}^{\infty} \frac{\cos\left(a + \frac{n\pi}{2}\right)}{n!} (x - a)^n.$$

$$\int x^n \mathrm{d} x = \begin{cases} \frac{1}{n+1} x^{n+1} + C, & n \neq -1 \\ \ln|x| + C, & n = -1 \end{cases}.$$

$$\int \ln(x) \mathrm{d} x = x \ln(x) - x + C.$$

$$\int (\ln(x))^n \mathrm{d} x = x (\ln(x))^n - n \int (\ln(x))^{n-1} \mathrm{d} x, \quad n \in \mathbb{N}.$$

$$\int e^{nx} \mathrm{d} x = \frac{1}{n} e^{nx} + C, \quad n \neq 0.$$

$$\int \sin(x) \mathrm{d} x = -\cos(x) + C.$$

$$\int \sin^2(x) \mathrm{d} x = \frac{x}{2} - \frac{\sin(2x)}{4} + C.$$

$$\int \sin^n(x) \mathrm{d} x = -\frac{1}{n} \sin^{n-1}(x) \cos(x) + \frac{n-1}{n} \int \sin^{n-2}(x) \mathrm{d} x, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cos(x) \mathrm{d} x = \sin(x) + C.$$

$$\int \cos^2(x) \mathrm{d} x = \frac{x}{2} + \frac{\sin(2x)}{4} + C.$$

$$\int \cos^n(x) \, dx = \frac{1}{n} \cos^{n-1}(x) \sin(x) + \frac{n-1}{n} \int \cos^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \tan(x) \, dx = -\ln |\cos(x)| + C = \ln |\sec(x)| + C.$$

$$\int \tan^2(x) \, dx = \tan(x) - x + C.$$

$$\int \tan^n(x) \, dx = \frac{1}{n-1} \tan^{n-1}(x) - \int \tan^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cot(x) \, dx = \ln |\sin(x)| + C.$$

$$\int \cot^2(x) \, dx = -\cot(x) - x + C.$$

$$\int \cot^n(x) \, dx = -\frac{1}{n-1} \cot^{n-1}(x) - \int \cot^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \sec(x) \, dx = \ln |\tan(x) + \sec(x)| + C = \ln \left| \tan\left(\frac{x}{2} + \frac{\pi}{4}\right) \right| + C.$$

$$\int \sec^2(x) \, dx = \tan(x) + C.$$

$$\int \sec^n(x) \, dx = \frac{1}{n-1} \tan(x) \sec^{n-2}(x) + \frac{n-2}{n-1} \int \sec^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \csc(x) \, dx = -\ln |\cot(x) + \csc(x)| + C = -\ln \left| \cot\left(\frac{x}{2}\right) \right| + C.$$

$$\int \csc^2(x) \, dx = -\cot(x) + C.$$

$$\int \csc^n(x) \, dx = -\frac{1}{n-1} \cot(x) \csc^{n-2}(x) + \frac{n-2}{n-1} \int \csc^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \arcsin(x) \, dx = x \arcsin(x) + \sqrt{1-x^2} + C.$$

$$\int \arccos(x) \, dx = x \arccos(x) - \sqrt{1-x^2} + C.$$

$$\int \arctan(x) \, dx = x \arctan(x) - \frac{1}{2} \ln(1+x^2) + C.$$

$$\int \operatorname{arccot}(x) \, dx = x \operatorname{arccot}(x) + \frac{1}{2} \ln(1+x^2) + C.$$

$$\int \operatorname{arcsec}(x) \, dx = x \operatorname{arcsec}(x) - \operatorname{sgn}(x) \ln \left| x + \sqrt{x^2-1} \right| + C, \quad |x| \geq 1.$$

$$\int \operatorname{arccsc}(x) \, dx = x \operatorname{arccsc}(x) + \operatorname{sgn}(x) \ln \left| x + \sqrt{x^2-1} \right| + C, \quad |x| \geq 1.$$

$$\int \sinh(x) \, dx = \cosh(x) + C.$$

$$\int \sinh^2(x) \, dx = -\frac{x}{2} + \frac{\sinh(2x)}{4} + C.$$

$$\int \sinh^n(x) \, dx = \frac{1}{n} \sinh^{n-1}(x) \cosh(x) - \frac{n-1}{n} \int \sinh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \cosh(x) \, dx = \sinh(x) + C.$$

$$\int \cosh^2(x) \, dx = \frac{x}{2} + \frac{\sinh(2x)}{4} + C.$$

$$\int \cosh^n(x) \, dx = \frac{1}{n} \cosh^{n-1}(x) \sinh(x) + \frac{n-1}{n} \int \cosh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \tanh(x) \, dx = \ln(\cosh(x)) + C.$$

$$\int \tanh^2(x) \, dx = x - \tanh(x) + C.$$

$$\int \tanh^n(x) \, dx = \frac{1}{n-1} \tanh^{n-1}(x) - \int \tanh^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \coth(x) \, dx = \ln |\sinh(x)| + C.$$

$$\int \coth^2(x) \, dx = x - \coth(x) + C.$$

$$\int \coth^n(x) \, dx = -\frac{1}{n-1} \coth^{n-1}(x) - \int \coth^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{sech}(x) \, dx = 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + C = \arctan(\sinh(x)) + C = \arcsin(\tanh(x)) + C.$$

$$\int \operatorname{sech}^2(x) \, dx = \tanh(x) + C.$$

$$\int \operatorname{sech}^n(x) \, dx = \frac{1}{n-1} \tanh(x) \operatorname{sech}^{n-1}(x) + \frac{n-2}{n-1} \int \operatorname{sech}^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{csch}(x) \, dx = \ln \left| \tanh\left(\frac{x}{2}\right) \right| + C = \ln |\coth(x) - \operatorname{csch}(x)| + C.$$

$$\int \operatorname{csch}^2(x) \, dx = -\coth(x) + C.$$

$$\int \operatorname{csch}^n(x) \, dx = -\frac{1}{n-1} \coth(x) \operatorname{csch}^{n-1}(x) + \frac{n-2}{n-1} \int \operatorname{csch}^{n-2}(x) \, dx, \quad n \in \mathbb{N}_{>2}.$$

$$\int \operatorname{arcsinh}(x) \, dx = x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1} + C.$$

$$\int \operatorname{arccosh}(x) \, dx = x \operatorname{arccosh}(x) - \sqrt{x^2 - 1} + C.$$

$$\int \operatorname{arctanh}(x) \, dx = x \operatorname{arctanh}(x) + \frac{1}{2} \ln(1 - x^2) + C.$$

$$\int \operatorname{arcoth}(x) \, dx = x \operatorname{arcoth}(x) + \frac{1}{2} \ln(x^2 - 1) + C.$$

$$\int \operatorname{arcsech}(x) \, dx = x \operatorname{arcsech}(x) + \arctan\left(\frac{\sqrt{1-x^2}}{x}\right) + C.$$

$$\int \operatorname{arccsch}(x) \, dx = x \operatorname{arccsch}(x) + \ln \left| x + \sqrt{x^2 + 1} \right| + C.$$

$$\int \frac{1}{x^2 + a^2} \, dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C, \quad a \neq 0.$$

$$\int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \ln \left(\frac{x-a}{x+a} \right) + C, \quad a \neq 0.$$

$$\int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \ln \left(\frac{a-x}{a+x} \right) + C, \quad a \neq 0.$$

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln |ax+b| + C, \quad a \neq 0.$$

$$\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + C, \quad a \neq 0 \wedge n \neq -1.$$

$$\int \frac{x}{ax+b} dx = \frac{x}{a} - \frac{b}{a^2} \ln |ax+b| + C, \quad a \neq 0.$$

$$\int \frac{x}{(ax+b)^2} dx = \frac{1}{a^2} \ln |ax+b| + \frac{b}{a^2(ax+b)} + C, \quad a \neq 0.$$

$$\int x(ax+b)^n dx = \frac{a(n+1)x-b}{a^2(n+1)(n+2)} (ax+b)^{n+1} + C, \quad a \neq 0 \wedge n \notin \{-1, -2\}.$$

Let $r = \sqrt{a^2 + x^2}$.

$$\int r dx = \frac{1}{2} (xr + a^2 \ln(x+r)) + C.$$

$$\int xr dx = \frac{1}{3} r^3 + C.$$

$$\int \frac{1}{r} dx = \operatorname{arcsinh} \left(\frac{x}{|a|} \right) + C = \ln(x+r) + C.$$

$$\int \frac{x}{r} dx = r + C.$$

Let $s = \sqrt{x^2 - a^2}$, $x^2 > a^2$.

$$\int s dx = \frac{1}{2} (xs - a^2 \ln(x+s)) + C.$$

$$\int xs dx = \frac{1}{3} s^3 + C.$$

$$\int \frac{1}{s} dx = \arcsin \left(\frac{x}{|a|} \right) + C.$$

$$\int \frac{x}{s} dx = -s + C.$$

Let $q = \sqrt{a^2 - x^2}$, $a^2 \geq x^2$.

$$\int q dx = \frac{1}{2} \left(q + a^2 \arcsin \left(\frac{x}{|a|} \right) \right) + C.$$

$$\int xq dx = \frac{1}{3} q^3 + C.$$

$$\int \frac{1}{q} dx = \arcsin \left(\frac{x}{|a|} \right) + C.$$

$$\int \frac{x}{q} dx = q + C.$$

Let $R = \sqrt{ax+b}$, $a \neq 0$.

$$\int R dx = \frac{2}{3a} R^3 + C.$$

$$\int x^n R \, \mathrm{d}x = \frac{2}{a(2n+3)} \left(x^n R^3 - bn \int x^{n-1} R \, \mathrm{d}x \right), \quad n \in \mathbb{N}.$$

$$\int \frac{1}{R} \, \mathrm{d}x = \frac{2R}{a} + C.$$

$$\int \frac{x^n}{R} \, \mathrm{d}x = \frac{2}{a} \left(x^n R - n \int x^{n-1} R \, \mathrm{d}x \right) = \frac{2}{a(2n+1)} \left(x^n R - bn \int \frac{x^{n-1}}{R} \, \mathrm{d}x \right), \quad n \in \mathbb{N}.$$