Logarithm and Exponentiation

沈威宇

November 1, 2026

Contents

1 Logarithm (對數) and Exponentiation (指數) 1

I Definitions 1

i Principal branch (主分支) 1

ii Positive integer exponent (正整數指數) 1

iii Natural logarithm (自然對數) in real numbers 1

iv Natural logarithm in complex numbers 1

v Logarithm (對數) in real numbers 1

vi Logarithm in complex numbers 1

vii Common logarithm (常用對數) 1

viii Exponent (指數) in real numbers 2

ix Euler's number (歐拉數/尤拉數) 2

x Euler's formula (歐拉公式) 3

xi Exponent in complex numbers 3

xii Root (根號) 4

xiii Scientific notation (科學記號) 4

xiv Characteristic (首數) and mantissa (尾數) 4

xv Logarithmic function (對數函數) 4

xvi Exponential function (指數函數) 4

xvii Natural exponential function (自然指數函數) 4

xviii Exponential growth (指數成長) 4

II Laws 5

i Logarithmic laws (對數律) 5

ii Exponential laws (指數律) 5

iii Inverse 5

iv Tangent 5

v Self-power function 5

III Values 6

1 Logarithm (對數) and Exponentiation (指數)

I Definitions

i Principal branch (主分支)

In the following definitions, $\arg\left(x \in \mathbb{C}_{\neq 0}\right)$ represents the argument (輻角) of x. If we replace $\arg(x)$ with the principal argument (輻角主值) of x, we get the definition of the principal branch of them.

ii Positive integer exponent (正整數指數)

The exponent of $a \in \mathbb{R}$ to the power of $n \in \mathbb{N}$, denoted as a^n, is defined as

$$a^n \coloneqq \prod_{k=1}^{n} a.$$

iii Natural logarithm (自然對數) in real numbers

The natural logarithm of x, denoted as $\ln(x)$ or $\log_e(x)$, is defined in real numbers as:

$$\ln(x) \coloneqq \int_1^x \frac{1}{t}\,\mathrm{d}t, \quad x \in \mathbb{R}_{>0}.$$

iv Natural logarithm in complex numbers

The natural logarithm of x, denoted as $\ln(x)$ or $\log_e(x)$, is defined in complex numbers as:

$$\ln|x| + i\arg(x), \quad x \in \mathbb{C}_{\neq 0}.$$

v Logarithm (對數) in real numbers

The logarithm of x, called argument (真數), to the a, called base (底數), denoted as $\log_a(x)$, is defined in real numbers as

$$\log_a(x) \coloneqq \frac{\ln(x)}{\ln(a)}, \quad a, x \in \mathbb{R}_{\neq 0} \wedge a \neq 1.$$

vi Logarithm in complex numbers

The logarithm of x, called argument, to the a, called base, denoted as $\log_a(x)$, is defined in complex numbers as

$$\log_a(x) \coloneqq \frac{\ln(x)}{\ln(a)}, \quad a, x \in \mathbb{C}_{\neq 0} \wedge a \neq 1.$$

vii Common logarithm (常用對數)

$\log_{10}(x)$ is called the common logarithm of x, also denoted as $\log(x)$.

viii Exponent (指數) in real numbers

The exponent of a, called the base (底數), to the exponent (指數) or power (冪次) of n, denoted as a^n, is defined in real numbers as

$$a^n \coloneqq \begin{cases} e^{n\ln(a)}, & a \in \mathbb{R}_{>0} \land n \in \mathbb{R}, \\ -|a|^n, & a \in \mathbb{R}_{<0} \land \exists p, q \in \mathbb{N} \text{ s.t. } n = \dfrac{2p-1}{2q-1}, \\ |a|^n, & a \in \mathbb{R}_{<0} \land \exists p, q \in \mathbb{N} \text{ s.t. } n = \dfrac{2p-2}{2q-1}, \\ 0, & a = 0 \land n \in \mathbb{R}_{>0}. \end{cases}$$

Exponentiation is right-associative, that is,

$$a^{b^c} = a^{(b^c)}, \quad \text{not } (a^b)^c.$$

ix Euler's number (歐拉數/尤拉數)

The Euler's number, denoted as e, is defined as

$$e = \lim_{n\to\infty} \left(1 + \frac{1}{n}\right)^n.$$

Some other equivalent definitions include:

$$e = \sum_{n=0}^{\infty} \frac{1}{n!}.$$

Proof.

$$\begin{aligned} \left(1 + \frac{1}{n}\right)^n &= \sum_{k=0}^{n} \binom{n}{k} \frac{1}{n^k} \\ &= \sum_{k=0}^{n} \frac{\prod_{j=0}^{k-1}(n-j)}{n^k k!} \\ &= \sum_{k=0}^{n} \prod_{j=0}^{k-1} \left(1 - \frac{j}{n}\right) \frac{1}{k!} \end{aligned}$$

For fixed k:

$$\lim_{n\to\infty} \prod_{j=0}^{k-1} \left(1 - \frac{j}{n}\right) = 1$$

$$\lim_{n\to\infty} \sum_{k=0}^{n} \prod_{j=0}^{k-1} \left(1 - \frac{j}{n}\right) \frac{1}{k!} = \sum_{k=0}^{\infty} \frac{1}{k!}.$$

□

$$e^x \coloneqq \text{ Solution of } f'(x) = f(x), \quad f(0) = 1,$$

that is,

$$\lim_{h\to 0} \frac{e^h - 1}{h} = 1.$$

Proof. By separation of variables:

$$\frac{\mathrm{d}y}{\mathrm{d}x} = y$$
$$\frac{\mathrm{d}y}{y} = \mathrm{d}x\,.$$
$$\ln|y| = x + C$$
$$y(x) = Ce^x$$
$$C = 1$$
$$y(x) = e^x.$$

□

$$\int_1^e \frac{\mathrm{d}x}{x} = 1.$$

Proof.

$$f(x) \coloneqq \sum_{n=0}^{\infty} \frac{x^n}{n!}.$$

$$f'(x) = \sum_{n=1}^{\infty} \frac{x^{n-1}}{(n-1)!} = f(x).$$

By uniqueness of solution of $f'(x) = f(x)$ and $f(0) = 1$,

$$f(x) = e^x.$$

Threrefore,

$$e = f(1).$$

By the definition of natural logarithm:

$$\ln y = \int_1^y \frac{\mathrm{d}x}{x}.$$
$$\int_1^e \frac{\mathrm{d}x}{x} = 1.$$

□

x Euler's formula (歐拉公式)

For any real number θ:

$$e^{i\theta} = \cos\theta + i\sin\theta.$$

xi Exponent in complex numbers

The exponent of a, called the base, to the exponent or power of n, denoted as a^n, is defined in complex numbers as

$$a^n \coloneqq \begin{cases} e^{n\ln(a)}, & a \in \mathbb{C}_{\neq 0} \land n \in \mathbb{C}, \\ 0, & a = 0 \land n \in \mathbb{R}_{>0}. \end{cases}$$

Exponentiation is right-associative, that is,

$$a^{b^c} = a^{(b^c)}, \quad \text{not } (a^b)^c.$$

xii Root (根號)

The yth root (y 次方根) of $w \in \mathbb{C}$ and y such that $w^{\frac{1}{y}}$ is defined, denoted as $\sqrt[y]{w}$ is defined as the principal branch of $w^{\frac{1}{y}}$, where $\sqrt{\ }$ is called the radical symbol, radical sign, root symbol, or surd (根號).

xiii Scientific notation (科學記號)

Scientific notation refers to the representation of a real number in the form of $a \times 10^n$, where $1 \leq |a| < 10 \wedge n \in \mathbb{Z}$.

xiv Characteristic (首數) and mantissa (尾數)

Given a common logarithm $\log(x)$, the characteristic of it is defined as $\lfloor \log(x) \rfloor$, and the mantissa of it is defined as $\log(x) - \lfloor \log(x) \rfloor$.

xv Logarithmic function (對數函數)

$f(x) = k \log_a(x)$ where $a > 0 \wedge a \neq 1 \wedge k \in \mathbb{R}_{\neq 0}$ is called a logarithmic function with base a, of which the domain is $\mathbb{R}_{>0}$ and the range is $\mathbb{R}$.

xvi Exponential function (指數函數)

$f(x) = ka^x$ where $a \neq 0 \wedge k \in \mathbb{R}_{\neq 0}$ is called an exponential function with base a, of which the domain is $\mathbb{R}$ and the range is $\mathbb{R}_{>0}$ when $k > 0$ and $\mathbb{R}_{<0}$ when $k < 0$.

xvii Natural exponential function (自然指數函數)

e^x is called the natural exponential function.

xviii Exponential growth (指數成長)

Exponential growth refers to a real function such that $f'(x) = kf(x)$ where $k \in \mathbb{R}_{\neq 0}$ (or $k \in \mathbb{R}_{>0}$ in some contexts), that is, $f(x) = e^{kx} + c$ where $c \in \mathbb{R}$. And the equation $f'(x) = kf(x)$ is sometimes called the law of natural growth.

Exponential decay refers to a real function such that $f'(x) = kf(x)$ where $k \in \mathbb{R}_{<0}$, that is, $f(x) = e^{kx} + c$ where $c \in \mathbb{R}$. And the equation $f'(x) = kf(x)$ is sometimes called the law of natural decay.

II Laws

i Logarithmic laws (對數律)

$$\log_a(r) + \log_a(s) = \log_a(rs)$$
$$\log_a(r) - \log_a(s) = \log_a\left(\frac{r}{s}\right)$$
$$\log_{a^m}\left(r^n\right) = \frac{n}{m}\log_a(r)$$
$$\log_a(b) = \frac{\log_c(b)}{\log_c\{a\}}$$
$$a^{\log_c(b)} = b^{\log_c(a)}$$

ii Exponential laws (指數律)

$$a^r \cdot a^s = a^{r+s},$$
$$(a^r)^s = a^{rs},$$
$$(a \cdot b)^r = a^r \cdot b^r$$

iii Inverse

For any $a > 0 \wedge a \neq 1 \wedge b \neq 0$, the logarithmic function $k\log_a\left(\frac{x}{b}\right)$ and the exponential function $ba^{\frac{x}{k}}$ are inverses of each other, that is, they are symmetric about $y = x$.

iv Tangent

In the xy plane, $a = e^{\left(e^{-1}\right)}$ if and only if any two of $y = a^x$, $y = x$, and $y = \log_a x$ are tangent to each other. If $a = e^{\left(e^{-1}\right)}$, $y = a^x$, $y = x$, and $y = \log_a x$ are tangent at (e, e), and any two of them do not meet at any point other than (e, e).

Proof.

$$\frac{\mathrm{d}a^x}{\mathrm{d}x} = \ln(a)a^x = 1$$
$$x = a^x = \frac{1}{\ln(a)}$$
$$a^{\frac{1}{\ln(a)}} = \frac{1}{\ln(a)}$$
$$\ln(a)\frac{1}{\ln(a)} = 1 = -\ln\left(\ln(a)\right)$$
$$a = e^{\left(e^{-1}\right)}$$

□

v Self-power function

$$\arg\min\left(x^x\right) = \frac{1}{e}, \quad \min\left(x^x\right) = \left(e^{-1}\right)^{\left(e^{-1}\right)}$$

III Values

$\log 2 \approx 0.3010$, $\log e \approx 0.4343$, $\log 3 \approx 0.4771$, $\log \pi \approx 0.4971$, $\log 4 \approx 0.6021$, $\log 5 \approx 0.6990$,

$\log 6 \approx 0.7782$, $\log 7 \approx 0.8451$, $\log 8 \approx 0.9031$, $\log 9 \approx 0.9542$, $\log 11 \approx 1.0414$, $\log 12 \approx 1.0792$.

$$\ln 2 \approx 0.6931,\quad \ln 3 \approx 1.0986,\quad \ln \pi \approx 1.1447,\quad \ln 4 \approx 1.3863,\quad \ln 5 \approx 1.6094,\quad \ln 6 \approx 1.7918,$$

$$\ln 7 \approx 1.9459,\quad \ln 10 \approx 2.3026,\quad \ln 11 \approx 2.3980,\quad \ln 12 \approx 2.4849,\quad \frac{1}{\ln 2} \approx 1.4427.$$

$$\sqrt{2} \approx 1.4142,\quad \sqrt{e} = 1.6487,\quad \sqrt{3} \approx 1.7321,\quad \sqrt{\pi} \approx 1.7725,\quad \sqrt{5} \approx 2.2361,$$

$$\sqrt{6} \approx 2.4495,\quad \sqrt{7} \approx 2.6458,\quad \sqrt{10} \approx 3.1623,\quad \sqrt{11} \approx 3.3166,\quad \sqrt{13} \approx 3.6056,$$

$$\sqrt{14} \approx 3.7417,\quad \sqrt{15} \approx 3.8730,\quad \sqrt{17} \approx 4.1231,\quad \sqrt{19} \approx 4.3590,\quad \sqrt{21} \approx 4.5826,$$

$$\frac{\sqrt{5}+1}{2} \approx 1.6180,\quad \frac{\sqrt{5}-1}{2} \approx 0.6180,\quad \frac{\sqrt{6}+\sqrt{2}}{4} \approx 0.9659,\quad \frac{\sqrt{6}-\sqrt{2}}{4} \approx 0.2588,$$

$$\frac{\sqrt{10+2\sqrt{5}}}{4} \approx 0.9511,\quad \frac{\sqrt{10-2\sqrt{5}}}{4} \approx 0.5878.$$

$$\langle a_n = n^2 \rangle_{n=1}^{20} = 1, 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, 144, 169, 196, 225, 256, 289, 324, 361, 400.$$

$$\langle a_n = n^2 \rangle_{n=21}^{30} = 441, 484, 529, 576, 625, 676, 729, 784, 841, 900.$$

$$\langle a_n = n^2 \rangle_{n=31}^{40} = 961, 1024, 1089, 1156, 1225, 1296, 1369, 1444, 1521, 1600.$$

$$\langle a_n = n^{-1} \rangle_{n=1}^{15} = 1, 0.5, 0.\bar{3}, 0.25, 0.2, 0.1\bar{6}, 0.\overline{142857}, 0.125, 0.\bar{1}, 0.1, 0.\overline{09}, 0.08\bar{3}, 0.\overline{076923}, 0.0\overline{714285}, 0.0\bar{6}.$$

$$\frac{1}{\sqrt{2}} \approx 0.7071,\quad \frac{1}{\sqrt{3}} \approx 0.5774,\quad \frac{1}{\sqrt{5}} \approx 0.4772,\quad \frac{1}{2\sqrt{2}} \approx 0.3536,\quad \frac{1}{2\sqrt{3}} \approx 0.2887,\quad \frac{1}{2\sqrt{5}} \approx 0.2236.$$

$$e \approx 2.7183,\quad \frac{1}{e} \approx 0.3680,\quad e^e \approx 15.1543,\quad e^{\left(e^{-1}\right)} \approx 1.4447,\quad \left(e^{-1}\right)^e \approx 0.0660,\quad \left(e^{-1}\right)^{\left(e^{-1}\right)} \approx 0.6922.$$