

Linear Algebra and Functional Analysis

沈威宇

June 9, 2026

Contents

Chapter 1	Linear Algebra (線性代數) and Functional Analysis (泛函分析)	1
1	Vector Space and Functional	1
I	Spaces and maps	1
i	Vector space (向量空間) or Linear space (線性空間)	1
ii	Isomorphism (同構)	2
iii	Linear combination (線性組合)	2
iv	Division	2
v	Affine space (仿射空間)	2
vi	Linear independence (線性獨立)	3
vii	Linear map (線性映射) or linear operator	3
viii	Linear transformation (線性變換)	3
ix	Invertible map (可逆映射)	3
x	Basis (基)	3
xi	Orientation	3
xii	Dimension (維度)	4
xiii	Bilinear form (雙線性形式)	4
xiv	Symmetric bilinear form (對稱雙線性形式)	4
xv	Topological vector space (TVS) (拓撲向量空間)	4
xvi	Functional (泛函)	4
xvii	Seminorm (半範數)	4
xviii	Norm (範數/模長)	5
xix	Sublinear function (亞線性函數) or quasi-seminorm (準半範數)	5
xx	Operator norm or induced norm	5
xxi	Absolutely continuous function (絕對連續函數)	5
xxii	Bounded linear operator (有界線性運算子)	6

xxiii Hahn–Banach theorem (哈恩-巴拿赫定理)	6
xxiv p-norm (p-範數)	6
xxv Locally convex topological vector space (LCTVS) or locally convex space .	6
xxvi Banach space (巴拿赫空間)	7
xxvii Dot product (點積)	7
xxviii Inner product space (內積空間)	7
xxix Inner product is bilinear form	7
xxx Orthonormal basis (正交基).	7
xxxi Balanced set (平衡集), circled set, or disk set	8
xxxii Well-defined (定義良好)/Unambiguous (不模糊的)	8
xxxiii Affine map (仿射映射).	8
xxxiv Affine transformation (仿射變換).	8
xxxv Euclidean vector space	8
xxxvi Von Neumann bounded or bounded.	8
xxxvii Euclidean affine space (歐幾里德仿射空間) or Euclidean space (歐幾里得空間)	9
xxxviii Convex set (凸集)	9
xxxix (Algebraic) dual (對偶) (vector) (space)	9
xl Continuous dual (space)	9
xli Transpose, dual map, or adjoint	9
xlII Strong dual (space).	9
xlIII Orthogonal complement (正交補)	10
xlIV Affine hull (仿射包).	10
xlV Convex hull (凸包)	10
xlVI Coordinate system (座標/坐標系統)	10
xlVII Cartesian coordinate system (笛卡爾座標系統) or rectangular coordinate system (直角座標系統)	10
xlVIII Lattice points (格子點).	10
xlIX Quadrant (象限)	11
I Right-hand rule (右手定則).	11
II Octant (卦限)	11
II Curvilinear coordinate system	11
IIII Projection (投影)	11

liv Orthographic projection (正射影)	11
lv Hilbert space (希爾伯特空間)	11
lvi Angle (夾角) between two vectors	11
lvii Principal angles (主夾角) between two subspaces	12
lviii Flat, hyperplane, or affine subspace (仿射子空間)	12
lix Hypersurface.	12
lx Parallel (平行)	12
lxi Coplanar.	12
lxii Elementwise multiplication	12
lxiii Unit vector (單位向量) or normalized vector	12
lxiv Affine independence.	13
lxv Simplex	13
lxvi Standard, unit, or probability simplex	13
lxvii Gram matrix, Gramian matrix, or Gramian	13
lxviii Support of functional	13
lxix Rectangle	14
lxx Tensor	14
lxxi Upper and lower indices	14
II In three dimensions	15
i Cross product (叉積), external product (外積), or vector product (向量積).	15
ii Basic properties of cross product	15
iii Triple product identity (三重積恆等式)/Jacobi identity (雅可比恆等式)	16
iv Scalar triple product.	16
2 Matrix (矩陣)	16
I Matrix	16
i Size	16
ii Notation	16
II Matrix equality.	17
III Basic operations	17
i Matrix addition.	17
ii Scalar multiplication	17
iii Substraction	17

iv Matrix multiplication (矩陣乘法).	18
v Positive integer exponent	18
vi Zero matrix (零矩陣)	18
vii Identity matrix (單位矩陣)	18
viii Kronecker product (克羅內克積)	18
ix Hadamard product (哈達瑪乘積), Schur product (舒爾乘積), elementwise product, or entrywise product (逐項乘積)	18
x Property	19
xi Zero exponent	19
IV Matrix row and column operations	20
i Row operations (zh-hant: 列/zh-hans: 行運算).	20
ii (General) row echelon form ((廣義) zh-hant: 列/zh-hans: 行階梯形式)	20
iii Column operations (zh-hant: 行/zh-hans: 列運算)	20
iv Transpose (轉置).	20
v Symmetric matrix (對稱矩陣)	20
vi Skew-symmetric matrix/antisymmetric matrix (反對稱矩陣)	20
vii Transpose identities.	21
V Trace (跡)	21
i Conjugate transpose (共軛轉置)/Hermitian conjugate (埃爾米特共軛)/Hermitian transpose (埃爾米特轉置)	21
VI Cofactor matrix, adjugate matrix, determinant, and inverse.	21
i Determinant (行列式或判別式)	21
ii Determinant identities	21
iii Cofactor matrix (餘因子矩陣)	22
iv Adjugate matrix (伴隨矩陣)/classical adjoint (經典伴隨矩陣)	22
v Inverse (反方陣、逆方陣、反矩陣或逆矩陣), multiplicative inverse (乘法反方陣), or reciprocal.	22
vi Negative integer exponent.	22
vii Invertible matrix (可逆矩陣), orthogonal matrix (正交矩陣), orthonormal matrix (正則矩陣), non-singular matrix (非奇異矩陣), or non-degenerate matrix (非退化矩陣)	22
viii Non-invertible matrix (不可逆矩陣), non-orthogonal matrix (非正交矩陣), non-orthonormal matrix (非正則矩陣), singular matrix (奇異矩陣), or degenerate matrix (退化矩陣).	22

ix	Inverse of a two-by-two matrix	22
x	Computing inverse with row operations	23
xi	Unitary matrix (么正矩陣或酉矩陣).	23
xii	Determinant property.	23
xiii	Inverse identities.	23
xiv	Vandermonde matrix (范德蒙矩陣).	23
VII	Diagonal matrix (對角矩陣).	23
i	Diagonal matrix	23
ii	Diagonal matrix identities	24
VIII	System of Linear Equations (線性方程組)	24
i	System of equations (方程組)	24
ii	Linear Equation (線性方程)	24
iii	System of Linear Equations (線性方程組).	24
iv	Coefficient matrix (係數矩陣)	25
v	Augmented matrix (增廣矩陣).	25
vi	Gaussian Elimination (高斯消去法) or row reduction (zh-hant: 列/zh-hans: 行簡化)	25
vii	Cramer Rule or Cramer formula (克拉瑪公式)	25
3	Linear Subspace (線性子空間)	26
I	Linear subspace	26
II	Four fundamental subspaces of a matrix (矩陣的四大子空間).	26
III	Column space (zh-hant: 行/zh-hans: 列空間)	26
IV	Row space (zh-hant: 列/zh-hans: 行空間).	26
V	Null space (零空間), kernel (核), or kernel space (核空間)	26
VI	Left null space (左零空間), left kernel (左核), or left kernel space (左核空間)	26
VII	Image space (像空間) or range (值域)	26
VIII	Cokernel (餘核).	26
IX	Constant rank theorem (常秩定理).	27
X	Rank (秩)	27
XI	Rank-nullity theorem (秩—零化度定理).	27
XII	Full rank (滿秩) and rank deficiency (秩虧)	29
XIII	Property of rank	29

4	Matrix Analysis (矩陣分析)	29
I	Eigenvector or characteristic vector (特徵向量) and characteristic polynomial (特徵多項式)	29
i	Eigenvector or characteristic vector (特徵向量) and eigenvalue (本徵值、固有值、特徵值或特徵根)	29
ii	Characteristic polynomial	29
iii	Property	30
iv	Cayley–Hamilton theorem (凱萊–哈密頓定理)	30
v	Diagonalization (對角化)	30
vi	Definiteness (正定性)	30
vii	Eigenvalues Test of Definiteness for Symmetric Hessians of Scalar Fields	31
II	Generalized Definition of Inverse	31
i	One-sided inverse (單邊逆)	31
ii	Generalized inverse or g-inverse (廣義逆)	32
iii	Reflexive generalized inverse (自反廣義逆)	32
iv	Moore–Penrose inverse (摩爾–彭若斯廣義逆) or pseudoinverse (偽逆或擬反)	32
5	Transformation Matrices	33
I	Groups	33
i	General linear group	33
ii	Orthogonal group or general orthogonal group	33
iii	Special orthogonal group or rotation group	33
II	Transformation matrices in two-dimensional space	33
i	Degenerate matrices	33
ii	Stretching (伸縮/縮放/拉伸)	34
iii	Shearing (推移)	34
iv	Rotation (旋轉)	34
v	Reflection (鏡射/反射)	34
vi	Orthogona projection (正射影)	34
vii	Exmample 1	35
viii	Exmample 2	35
ix	Example 3	35
x	Example 4	36

III	Transformation matrices in three-dimensional space	36
i	Stretching (伸縮/縮放/拉伸)	36
ii	Shearing (推移)	36
iii	Rotation (旋轉)	37
iv	Rodrigues' rotation formula	37
v	Reflection (鏡射/反射).	38
vi	Orthogona projection (正射影)	38

Chapter 1 Linear Algebra (線性代數) and Functional Analysis (泛函分析)

1 Vector Space and Functional

I Spaces and maps

i Vector space (向量空間) or Linear space (線性空間)

A vector space over a field F is a non-empty set V together with a binary operation and a binary function that satisfy the eight axioms listed below. In this context, the elements of V are commonly called vectors, and the elements of F are called scalars.

The binary operation, called vector addition or simply addition assigns to any two vectors v and w in V a third vector in V which is commonly written as $v + w$, and called the sum of these two vectors.

The binary function, called scalar multiplication, assigns to any scalar a in F and any vector v in V another vector in V , which is denoted av .

To have a vector space, the eight following axioms must be satisfied for every u, v and w in V , and a and b in F .

1. Associativity of vector addition: $u + (v + w) = (u + v) + w$.
2. Commutativity of vector addition: $u + v = v + u$.
3. Identity element of vector addition: There exists an element $0 \in V$, called the zero vector, such that $\forall v \in V : v + 0 = v$.
4. Inverse elements of vector addition: $\forall v \in V : \exists$ an element $-v \in V$, called the additive inverse or inverse of v , such that $v + (-v) = 0$.
5. Compatibility of scalar multiplication with field multiplication: $a(bv) = (ab)v$.
6. Identity element of scalar multiplication: $1v = v$, where 1 denotes the multiplicative identity in F .
7. Distributivity of scalar multiplication with respect to vector addition: $a(u + v) = au + av$.
8. Distributivity of scalar multiplication with respect to field addition: $(a + b)v = av + bv$.

When the scalar field is the real numbers, the vector space is called a real vector space, and when the scalar field is the complex numbers, the vector space is called a complex vector space. These two cases are the most common ones, but vector spaces with scalars in an arbitrary field F are also commonly considered. Such a vector space is called an F -vector space or a vector space over F .

When a vector consists of n scalar components, we typically denote it as

$$(x_1, x_2, \dots, x_n)$$

or

$$\langle x_1, x_2, \dots, x_n \rangle$$

ii Isomorphism (同構)

Vector spaces X and Y are called isomorphic if there exists a mapping $f : X \rightarrow Y$ between them such that f is bijective and continuous and f^{-1} is continuous, written as $X \cong Y$, and f is called an isomorphism between them.

iii Linear combination (線性組合)

A vector a in a vector space V over a field F is a linear combination of a series of vectors $(a_1, a_2, \dots, a_n)$ in a vector space V over a field F if there exists a series of scalars $(\alpha_1, \alpha_2, \dots, \alpha_n)$ in F such that:

$$a = \sum_{i=1}^n \alpha_i a_i.$$

iv Division

For nonzero vector a and vector b in a vector space such that for some scalar s , $sa = b$, $\frac{b}{a} := s$.

v Affine space (仿射空間)

An affine space is a point set A together with a vector space $\vec{A}$, and a transitive and free action of the additive group of $\vec{A}$ on the set A . The vector space $\vec{A}$ is said to be associated to the affine space, and its elements are called vectors, translations, or sometimes free vectors.

Explicitly, the definition above means that the action is a mapping, generally denoted as an addition,

$$\begin{aligned} A \times \vec{A} &\rightarrow A \\ (a, v) &\mapsto a + v \end{aligned}$$

that has the following properties.

1. Right identity:

$$\forall a \in A : a + 0 = a, \quad \text{where } 0 \text{ is the zero vector in } \vec{A}$$

2. Associativity:

$$\forall v, w \in \vec{A}, \forall a \in A, (a + v) + w = a + (v + w)$$

3. Free and transitive action:

$\forall a \in A$: the mapping $\vec{A} \rightarrow A : v \mapsto a + v$ is a bijection.

4. Existence of one-to-one translations

$\forall v \in \vec{A}$: the mapping $\vec{A} \rightarrow A : v \mapsto a + v$ is a bijection.

vi Linear independence (線性獨立)

For a set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$, if the equation:

$$\sum_{i=1}^n c_i \mathbf{v}_i = 0$$

is true only if all coefficients $c_1, c_2, \dots, c_n$ are zero, then the set of vectors is linearly independent.

vii Linear map (線性映射) or linear operator

For vector spaces V, W , a map $T : V \rightarrow W$ between vector spaces V and W over field F is linear if

$$\forall a, b \in F, v, w \in V : T(av + bw) = aT(v) + bT(w),$$

also called linear operator.

The space of linear operators from V to W is a vector space, denoted as $L(V, W)$.

viii Linear transformation (線性變換)

A linear transformation is a bijection from an vector space onto itself that is an linear map.

ix Invertible map (可逆映射)

A map $T : V \rightarrow W$ is Invertible if and only if it is bijective. The unique inverse of T , called T^{-1} , is defined to be:

$$T^{-1} : W \rightarrow V; \forall v \in V : T(v) \mapsto v = T^{-1}(T(v)).$$

x Basis (基)

The basis of a vector space X over a field F is a set of linearly independent vectors $\mathbf{v} = \{v_1, v_2, \dots\} \subseteq X$ such that for any vector $\mathbf{P} \in X$, there exists a unique coefficient vector $(c_1, c_2, \dots) \subseteq F$ such that:

$$\sum_{i=1} c_i v_i = \mathbf{P}.$$

xi Orientation

The orientation of a vector space is the arbitrary choice of which ordered bases are "positively" oriented and which are "negatively" oriented.

xii Dimension (維度)

The dimension of a vector space is defined as the cardinality of a basis of it.

The dimension of an affine space is defined as the dimension of its associated vector space.

The n -dimensional vector space over field F is usually written as F^n .

xiii Bilinear form (雙線性形式)

Let V be a vector space of dimension n over a field $\mathbb{K}$. A map $B : V \times V \rightarrow \mathbb{K}$ is a bilinear form on the space if:

1. $\forall u, v, w \in V : B(u + v, w) = B(u, w) + B(v, w).$
2. $\forall u, v \in V, \lambda \in \mathbb{K} : B(\lambda u, v) = \lambda B(u, v).$
3. $\forall u, v, w \in V : B(u, v + w) = B(u, v) + B(u, w).$
4. $\forall u, v \in V, \lambda \in \mathbb{K} : B(u, \lambda v) = \lambda B(u, v).$

xiv Symmetric bilinear form (對稱雙線性形式)

Let V be a vector space of dimension n over a field $\mathbb{K}$. A map $B : V \times V \rightarrow \mathbb{K}$ is a symmetric bilinear form on the space if:

1. $\forall u, v \in V : B(u, v) = B(v, u).$
2. $\forall u, v, w \in V : B(u + v, w) = B(u, w) + B(v, w).$
3. $\forall u, v \in V, \lambda \in \mathbb{K} : B(\lambda u, v) = \lambda B(u, v).$

xv Topological vector space (TVS) (拓樸向量空間)

A topological vector space X is a vector space over a topological field $\mathbb{K}$, such that vector addition $X \times X \rightarrow X$ and scalar multiplication $\mathbb{K} \times X \rightarrow X$ are continuous functions.

xvi Functional (泛函)

A functional is a function from a vector space into the field of real or complex numbers. For a vector space X , $x \in X$ and a functional on f on X , $f(x)$ is also denoted as $\langle f, x \rangle$ or $f x$.

xvii Seminorm (半範數)

Given a vector space X over a ordered field F , a seminorm on X is a real-valued function $p : X \rightarrow \mathbb{R}$ with the following conditions, where $|s|$ denotes the usual absolute value of a scalar s :

1. Subadditivity/Triangle inequality:

$$\forall x, y \in X : p(x + y) \leq p(x) + p(y).$$

2. Absolute homogeneity:

$$\forall s \in \mathbb{R}, x \in X : p(sx) = |s|p(x).$$

These conditions implies that:

1. Non-negativity: $\forall x \in X : p(x) \geq 0$.
2. $p(0) = 0$.

xviii Norm (範數/模長)

A norm on X is a seminorm $p : X \rightarrow \mathbb{R}$ with the following properties:

- Positive definiteness/Positiveness/Point-separating:

$$\forall x \in X : p(x) = 0 \implies x = 0$$

A vector x such that $p(x) = 1$ is called an unit vector.

xix Sublinear function (亞線性函數) or quasi-seminorm (準半範數)

Let $p : X \rightarrow \mathbb{R}$ be a function on a vector space X over the field $\mathbb{K}$, which is either $\mathbb{R}$ or $\mathbb{C}$. p is called a sublinear function or quasi-seminorm if it satisfies the following conditions:

1. Subadditivity/Triangle inequality:

$$\forall x, y \in X : p(x + y) \leq p(x) + p(y).$$

2. Nonnegative homogeneity:

$$\forall x \in X, r \geq 0 : p(rx) = rp(x).$$

xx Operator norm or induced norm

For normed vector spaces V, W , the vector space of linear operators $L(V, W)$ is equipped with a norm $\| \cdot \|_L$ called operator norm or induced norm defined as, for $A \in L(V, W)$,

$$\|A\|_L = \sup_{\|v\|_V \leq 1} \|A(v)\|_W.$$

xxi Absolutely continuous function (絕對連續函數)

Function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is absolutely continuous on I if for every positive number ε , there is a positive number δ such that, for every finite sequence of pairwise disjoint sub-intervals (x_k, y_k) of I with $x_k < y_k \in I$ and cardinality N ,

$$\sum_{k=1}^N (y_k - x_k) < \delta \implies \sum_{k=1}^N |f(y_k) - f(x_k)| < \varepsilon.$$

The collection of all absolutely continuous functions on I is denoted $AC(I)$.

xxii Bounded linear operator (有界線性運算子)

A bounded linear operator is a linear transformation $L : X \rightarrow Y$ between topological vector spaces X and Y that maps bounded subsets of X to bounded subsets of Y . If X and Y are normed vector spaces, then L is bounded if and only if there exists some $M > 0$ such that:

$$\forall x \in X : \|Lx\|_Y \leq M\|x\|_X.$$

The smallest such M is called the operator norm of L and denoted by $\|L\|$. A bounded operator between normed spaces is continuous and vice versa.

xxiii Hahn-Banach theorem (哈恩-巴拿赫定理)

Let $p : X \rightarrow \mathbb{R}$ be a sublinear functional on a vector space X over the field $\mathbb{K}$, which is either $\mathbb{R}$ or $\mathbb{C}$. If $f : M \rightarrow \mathbb{K}$ is a linear functional on a vector subspace M such that

$$\forall m \in M : f(m) \leq p(m),$$

then there exists a linear functional $F : X \rightarrow \mathbb{K}$ such that

$$\forall m \in M : F(m) = f(m),$$

$$\forall x \in X : F(x) \leq p(x).$$

xxiv p-norm (p-範數)

The p -norm of a n -dimensional vector $x = ((x_i)_{i=1}^n)$ is defined to be

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}, \quad p \in \mathbb{N}_{\infty},$$

where the ∞ -norm is

$$\|x\|_{\infty} = \max_{i=1}^n |x_i|.$$

The 1-norm is also called Manhattan norm (曼哈頓範數); the 2-norm is also called Euclidean norm (歐幾里德範數).

xxv Locally convex topological vector space (LCTVS) or locally convex space

A locally convex topological vector space (LCTVS) $(X, \mathcal{T})$ is a TVS whose topology is generated by a family of seminorms P on it such that:

$$\bigcap_{p \in P} \{x : p(x) = 0\} = \{0\}$$

That is, $\mathcal{T}$ is generated by a basis $\{U_{\epsilon, x}\}_{\epsilon \in \mathbb{R}_{\geq 0}, x \in X}$ of neighborhoods defined to be:

$$U_{\epsilon, 0} = \{y \in X : \forall p \in P : p(y) < \epsilon\}.$$

$$U_{\epsilon, x} = \{y \in X : y - x \in U_{\epsilon, 0}\}.$$

This definition implies that a LCTVS is necessarily a Hausdorff space (T2 space).

xxvi Banach space (巴拿赫空間)

A Banach space is a normed vector space $(X, \|\cdot\|)$ with induced metric d

$$d(x, y) = \|x - y\|, \quad \forall x, y \in X$$

that is a Complete metric space (aka Cauchy space).

xxvii Dot product (點積)

The dot product of two vectors $\mathbf{a} = (a_1, a_2, \dots, a_n)$ and $\mathbf{b} = (b_1, b_2, \dots, b_n)$ is defined as:

$$\mathbf{a} \cdot \mathbf{b} = \sum_{i=1}^n a_i b_i.$$

Two nonzero vector are called orthogonal or perpendicular iff their dot product is zero.

xxviii Inner product space (內積空間)

A bar over an expression representing a scalar denotes the complex conjugate of this scalar.

An inner product space is a vector space V over the field F together with an inner product, that is, a map

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow F$$

that satisfies the following three properties for all vectors $x, y, z \in V$ and all scalars $a, b \in F$.

- Conjugate symmetry:

$$\langle x, y \rangle = \overline{\langle y, x \rangle}.$$

This implies that $\langle x, x \rangle$ is a real number.

- Linearity in the first argument:

$$\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle.$$

- Positive-definiteness: if x is not zero, then $\langle x, x \rangle > 0$.

xxix Inner product is bilinear form

An inner product is a bilinear form. PLACEHOLDER

Proof. PLACEHOLDER

□

xxx Orthonormal basis (正交基)

An orthonormal basis for an inner product space V with finite dimension is a basis for V such that for e, f in it:

$$\langle e, f \rangle = \begin{cases} 1, & \text{if } e = f, \\ 0, & \text{if } e \neq f. \end{cases}$$

xxxi **Balanced set (平衡集), circled set, or disk set**

Let X be a vector space over a field $\mathbb{K}$ with an absolute value function $|\cdot|$. A subset S of X is called a balanced set or balanced if:

$$\forall a \in \mathbb{K} \text{ s.t. } |a| \leq 1 : aS \subseteq S.$$

xxxii **Well-defined (定義良好)/Unambiguous (不模糊的)**

A well-defined expression or unambiguous expression is an expression whose definition assigns it a unique interpretation or value. Otherwise, the expression is said to be not well defined, ill defined or ambiguous.

xxxiii **Affine map (仿射映射)**

Given two affine spaces A and B whose associated vector spaces are $\mathcal{A}$ and $\mathcal{B}$, an affine map from A to B is a map $f : A \rightarrow B$ such that

$$\begin{aligned} f &: \mathcal{A} \rightarrow \mathcal{B} \\ b - a &\mapsto f(b) - f(a) \end{aligned}$$

is a well-defined linear map.

xxxiv **Affine transformation (仿射變換)**

An affine transformation is a bijection from an affine space onto itself that is an affine map.

xxxv **Euclidean vector space**

A Euclidean vector space E is a finite-dimensional vector space over the real numbers equipped with an inner product

$$\langle \cdot, \cdot \rangle : E \times E \rightarrow \mathbb{R},$$

and a norm $\|\cdot\|$ defined as:

$$\|v\| = \sqrt{\langle v, v \rangle},$$

and a metric, called the Euclidean distance $d(\cdot, \cdot) : E \times E \rightarrow \mathbb{R}$, defined as:

$$d(v, w) = \|v - w\|.$$

xxxvi **Von Neumann bounded or bounded**

Suppose X is a topological vector space over a field $\mathbb{K}$. A subset B of X is called von Neumann bounded (or bounded) in X if for every neighborhood V of the origin, there exists a real $r > 0$ such that $B \subseteq sV$ for all scalars s satisfying $|s| \geq r$.

xxxvii Euclidean affine space (歐幾里德仿射空間) or Euclidean space (歐幾里得空間)

A Euclidean space, also known as Euclidean affine space, is an affine space over $\mathbb{R}$ such that the associated vector space is a Euclidean vector space.

A Euclidean space is equipped with a metric, called the Euclidean distance $d(\cdot, \cdot) : E \times E \rightarrow \mathbb{R}$, defined as:

$$d(v, w) = \|v - w\|.$$

xxxviii Convex set (凸集)

Let S be a vector space or an affine space over some ordered field. A subset C of S is convex if, for all x and y in C , the line segment connecting x and y is included in C .

xxxix (Algebraic) dual (對偶) (vector) (space)

Given any vector space V over a field F , the dual space V^* is defined as the set of all linear maps $\varphi : V \rightarrow F$. The dual space V^* itself thus becomes a vector space over F when equipped with an addition and scalar multiplication satisfying that for any $\varphi, \psi \in V^*$, for any $x \in V$, for any $a \in F$,

$$(\varphi + \psi)(x) = \varphi(x) + \psi(x),$$

$$(a\varphi)(x) = a(\varphi(x)).$$

Element in dual is called covector.

xl Continuous dual (space)

Given any vector space V over a field F , the continuous dual space V' is defined as the set of all continuous linear maps $\varphi : V \rightarrow F$. The continuous dual space V' itself thus becomes a vector space over F when equipped with an addition and scalar multiplication satisfying that for any $\varphi, \psi \in V'$, for any $x \in V$, for any $a \in F$,

$$(\varphi + \psi)(x) = \varphi(x) + \psi(x),$$

$$(a\varphi)(x) = a(\varphi(x)).$$

xli Transpose, dual map, or adjoint

For two vector spaces X, Y and a continuous linear map $A : X \rightarrow Y$, the transpose, dual map, or adjoint of A is defined as the unique map ${}^tA : Y' \rightarrow X'$ such that for all $x \in X$, for all $y \in Y'$,

$$\langle y, A(x) \rangle = \langle {}^tA(y), x \rangle.$$

xlii Strong dual (space)

A strong dual space is a continuous dual space of a TVS equipped with the strong (dual) topology, also called topology of uniform convergence on bounded subsets. Suppose X is

a TVS over field $\mathbb{F}$ and $\mathcal{B}$ is a family of bounded subsets of X such that every bounded subset of X is a subset of some $B \in \mathcal{B}$. Define the family T of seminorms $\|\cdot\|_B$ on X' for all $x' \in X'$ as

$$T = \{\|x'\|_B := \sup_{x \in B} |x'(x)| \mid B \in \mathcal{B}\}.$$

The strong dual topology is defined as the topology generated by T , and thus B' is a LCTVS.

xlili Orthogonal complement (正交補)

An orthogonal complement of a subspace W of a vector space V equipped with a bilinear form B is the set $W^\perp$ of all vectors in V that are orthogonal to every vector in W , of which $v \in V$ is orthogonal to $w \in V$ if and only if $B(v, w) = 0$.

xliv Affine hull (仿射包)

An affine combination is a linear combination of points where all coefficients sum up to 1.

The affine hull $\text{aff}(S)$ of a set of points S in a Euclidean space is the smallest affine space that contains all the points in S , namely, the set of all affine combinations of the points in S , i.e.:

$$\text{aff}(S) = \left\{ \sum_{i=1}^k \alpha_i x_i : k > 0, x_i \in S, \alpha_i \in \mathbb{R}, \sum_{i=1}^k \alpha_i = 1 \right\}$$

xlv Convex hull (凸包)

A convex combination is a linear combination of points where all coefficients are non-negative and sum up to 1.

The convex hull $\text{conv}(S)$ of a set of points S in a Euclidean space is the smallest convex set that contains all the points in S , namely, the set of all convex combinations of points in S , i.e.:

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \alpha_i x_i : k > 0, x_i \in S, \alpha_i \in [0, 1], \sum_{i=1}^k \alpha_i = 1 \right\}$$

xlvi Coordinate system (座標/坐標系統)

A coordinate system is a system that uses one or more numbers, or coordinates, to uniquely determine the points on a manifold.

xlvi Cartesian coordinate system (笛卡爾座標系統) or rectangular coordinate system (直角座標系統)

The Cartesian or rectangular coordinate system is a coordinate system in a Euclidean vector space E determined by an orthogonal basis of unit vectors such that the inner product of two vectors in E equals to the dot product of them, and that the norm of E is 2-norm.

xlvi Lattice points (格子點)

In a Cartesian coordinate system, vectors whose components are all integers are called lattice points.

xlix Quadrant (象限)

In two dimensions, I(+, +), II(-, +), III(-, -), IV(+, -).

I Right-hand rule (右手定則)

Right-hand rule is a common used orientation for a three-dimensional space. In a right-handed Cartesian coordinate system, if you point the thumb of your right hand in the positive x -axis direction and your index finger in the positive y -axis direction, then your middle finger (extended perpendicularly from the palm) will point in the positive z -axis direction. If not specified otherwise, right-handed Cartesian coordinate system is usually used.

li Octant (卦限)

In three dimensions, I(+, +, +), II(-, +, +), III(-, -, +), IV(+, -, +), V(+, +, -), VI(-, +, -), VII(-, -, -), VIII(+, -, -).

lii Curvilinear coordinate system

A curvilinear coordinate system is a coordinate system for Euclidean space such that there exists an invertible map from a Cartesian coordinate system to it.

liii Projection (投影)

A (vector) projection onto a subset U of a vector space V is a linear operator $P : V \rightarrow U$ such that $P^2 = P$, denoted as $\text{proj}_U P$. The property $P^2 = P$ is called idempotency (冪等性). Its norm is called scalar projection denoted as $\text{comp}_U P$.

liv Orthographic projection (正射影)

An orthographic on a subset U of a vector space V is a projection $P : V \rightarrow U$ such that

$$\forall v \in V : v - P(v) \in U^*$$

where U^* is the orthogonal complement of U .

lv Hilbert space (希爾伯特空間)

A Hilbert space is a real or complex inner product space that is also a complete metric space with respect to the distance function induced by the inner product.

lvi Angle (夾角) between two vectors

Given two vectors u, v in an inner product space, the angle θ between them is defined as

$$\theta = \arccos(u \cdot v).$$

Ivii Principal angles (主夾角) between two subspaces

Given two subspaces U, W of an inner product space with $\dim(U) = k \leq \dim(W) = m$, there exists then a sequence of k angles $0 \leq \theta_1 \leq \theta_2 \leq \dots \leq \theta_k \leq \frac{\pi}{2}$ called the principal angles between U and W , the first one defined as

$$\theta_1 := \min \left\{ \arccos \left(\frac{|\langle u, w \rangle|}{\|u\| \|w\|} \right) \middle| u \in U \wedge w \in W \right\} = \angle(u_1 \in U, w_1 \in W),$$

The vectors u_1 and w_1 are the corresponding principal vectors.

The other principal angles and vectors are then defined recursively via

$$\theta_i := \min \left\{ \arccos \left(\frac{|\langle u, w \rangle|}{\|u\| \|w\|} \right) \middle| u \in U \wedge w \in W \wedge \forall j \in \mathbb{N} \wedge j < i : \langle u_j, u \rangle = \langle w_j, w \rangle = 0 \right\} = \angle(u_i \in U, w_i \in W)$$

Iviii Flat, hyperplane, or affine subspace (仿射子空間)

A flat or an affine subspace of an affine space is a subspace of it that is an affine space.

lix Hypersurface

A hypersurface is a $(n - 1)$ -dimensional manifold that is a subspace of an n -dimensional manifold.

Ix Parallel (平行)

Two non-zero vectors u, v in a vector space over field F are called to be parallel if there exists a scalar $x \in F$ such that $xu = v$, denoted as $u \parallel v$.

Two affine subspaces are called to be parallel if the direction spaces of them D, E satisfies:

$$D \subseteq E \vee E \subseteq D,$$

denoted as $D \parallel E$.

Ixi Coplanar

Multiple elements of a vector or affine space are called coplanar if they are in the same 2D affine subspace.

Ixii Elementwise multiplication

For two vectors $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$, their elementwise multiplication, denoted as $\mathbf{u} \odot \mathbf{v}$, is defined as

$$\mathbf{u} \odot \mathbf{v} = (u_1 v_1, \dots, u_n v_n).$$

Ixiii Unit vector (單位向量) or normalized vector

A unit vector or a normalized vector in a normed vector space is a vector of norm 1, denoted by a lowercase letter with a circumflex or hat, as in $\hat{\mathbf{v}}$ (pronounced "v-hat"). The unit vector or

normalized vector $\hat{\mathbf{u}}$ of a non-zero vector $\mathbf{u}$ in a normed vector space equipped with a norm $\|\cdot\|$ is defined as:

$$\hat{\mathbf{u}} = \frac{\mathbf{u}}{\|\mathbf{u}\|}.$$

Ixiv Affine independence

For an affine space U associated with vector space V , $u_0, u_1, \dots, u_n \in U$ are affinely independent iff $(u_1 - u_0), (u_2 - u_0), \dots, (u_n - u_0)$ are linearly independent in V .

Ixv Simplex

A k -simplex or k -dimensional simplex in an affine space U is a set

$$\left\{ \sum_{i=0}^k \theta_i u_i \mid \sum_{i=0}^k \theta_i = 1 \wedge \theta_0, \theta_1, \dots, \theta_k \in \mathbb{R}_{\geq 0} \right\},$$

where

$$u_0, u_1, \dots, u_k \in U$$

are affine independent and called vertices of the simplex.

Ixvi Standard, unit, or probability simplex

The standard, unit, or probability k -simplex is the simplex in $\mathbb{R}^{k+1}$ whose vertices are the unit vectors of each axis, that is, the set

$$\{(p_0, p_1, \dots, p_k) \mid \sum_{i=0}^k p_i = 1 \wedge p_0, p_1, \dots, p_k \in \mathbb{R}_{\geq 0}\}.$$

Ixvii Gram matrix, Gramian matrix, or Gramian

For n vectors $v_1, v_2, \dots, v_n$ in an inner product space, the Gram matrix, Gramian matrix, or Gramian G is defined as an $n \times n$ matrix where

$$G_{ij} = \langle v_i, v_j \rangle.$$

The Gram determinant or Gramian is the determinant of the Gram matrix.

Ixviii Support of functional

For a function $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ (or $\mathbb{C}$), the support of it, denoted as $\text{supp}(f)$, is defined as

$$\text{supp}(f) = \text{cl}(\{x \in U \mid f(x) \neq 0\}).$$

If $\text{supp}(f)$ is a compact set, f is called with compact support. If $\text{supp}(f)$ is a compact subset of a set $V \subseteq U$, f is called with compact support on V .

Ixix Rectangle

An open rectangle in $\mathbb{R}^n$ is the product of n open intervals.

An closed rectangle in $\mathbb{R}^n$ is the product of n closed intervals.

Rectangle may refer to either, or product of n intervals, or product of n intervals satisfying certain constraints, depending on context, and often the case they make no non-trivial difference if not specified.

Ixx Tensor

Let V be a vector space and V^* be its dual. A tensor of type (r, s) is a multilinear map

$$T : (V^*)^r \times V^s \rightarrow \mathbb{R},$$

where r is the number of contravariant slots, which take covectors in V^* , and s is the number of covariant slots, which take vectors in V . The order or rank of it is $r + s$.

This map

$$T : (V^*)^r \times V^s \rightarrow \mathbb{R}$$

is also a map

$$T : (V^*)^q \times V^t \rightarrow \{S : (S : (V^*)^{r-q} \times V^{s-t} \rightarrow \mathbb{R})\}$$

for any $q \in \mathbb{Z} \cap [0, r]$ and $t \in \mathbb{Z} \cap [0, s]$.

A scalar is a tensor of type $(0, 0)$. A vector is a tensor of type $(1, 0)$, which maps a covector to a real number, thus also called a contravariant tensor. A covector is a tensor of type $(0, 1)$, which maps a vector to a real number, thus also called a covariant tensor. A matrix as a linear map $V \rightarrow V$ is a tensor of type $(1, 1)$. A matrix as a bilinear form $V \times V \rightarrow \mathbb{R}$ is a tensor of type $(0, 2)$. A matrix as $V^* \times V^* \rightarrow \mathbb{R}$ is a tensor of type $\mathbb{R}$.

It's conventional that a vector is a column vector and a covector is a row vector. Tensor convention follows.

A tensor can be represented as an array whose dimension is its rank when a basis is chosen.

Ixxi Upper and lower indices

Dummy indices such as μ and ν are used to indicate contravariant and covariant vectors and tensors.

We use a superscript of r dummy indices and a subscript of s dummy indices to indicate a tensor of type (r, s) , i.e., $(V^*)^r \times V^s \rightarrow \mathbb{R}$ for vector space V , such as $T_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r}$. Each i th component of each upper index μ is denoted by replacing μ with i . Each i th component of each lower index μ is denoted by replacing μ with i . For vectors, that is, $A^\mu = (A^0, A^1, A^2, A^3)$. For covectors, that is, $A_\mu = (A_0, A_1, A_2, A_3)$.

Each upper index μ is fixed to a summation $\sum_{\mu=0}^{\dim(V)}$ when it takes an input of the same upper index μ , i.e.

$$T^{\mu \dots 1} U^{\mu \dots 3} = \sum_{\mu=0}^{\dim(V)} T^{\mu \dots 1} U^{\mu \dots 3}.$$

Each lower index μ is fixed to a summation $\sum_{\mu=0}^{\dim(V)}$ when it takes an input of the same lower index μ , i.e.

$$T^{\dots 1}_{\mu \dots 2} U^{\dots 3}_{\mu \dots 4} = \sum_{\mu=0}^{\dim(V)} T^{\dots 1}_{\mu \dots 2} U^{\dots 3}_{\mu \dots 4}.$$

The indexing replaces the order of them in a formula and makes the multiplication notation of tensors work without extra notations.

II In three dimensions

i Cross product (叉積), external product (外積), or vector product (向量積)

The cross product of vector $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$ is defined as:

$$\mathbf{a} \times \mathbf{b} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{pmatrix},$$

where $\mathbf{i} = (1, 0, 0)$, $\mathbf{j} = (0, 1, 0)$, $\mathbf{k} = (0, 0, 1)$.

ii Basic properties of cross product

- Self cross product is the zero vector:

$$\mathbf{a} \times \mathbf{a} = \mathbf{0}.$$

- Anticommutative law (反交換律):

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}.$$

- Left and right distributive over addition and subtraction:

$$\mathbf{a} \times (\mathbf{b} \pm \mathbf{c}) = \mathbf{a} \times \mathbf{b} \pm \mathbf{a} \times \mathbf{c}.$$

$$(\mathbf{b} \pm \mathbf{c}) \times \mathbf{a} = \mathbf{b} \times \mathbf{a} \pm \mathbf{c} \times \mathbf{a}.$$

- Commutative and associative over scalar multiplication:

$$(r\mathbf{a}) \times \mathbf{b} = \mathbf{a} \times (r\mathbf{b}) = r(\mathbf{a} \times \mathbf{b}).$$

- Two vector in $\mathbb{R}^3$ are parallel iff their cross product is the zero vector.

iii Triple product identity (三重積恆等式)/Jacobi identity (雅可比恆等式)

$$\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) + \mathbf{B} \times (\mathbf{C} \times \mathbf{A}) + \mathbf{C} \times (\mathbf{A} \times \mathbf{B}) = \mathbf{0}.$$

$$\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \mathbf{B} \cdot (\mathbf{C} \times \mathbf{A}) = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B}) = \begin{vmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{vmatrix}$$

$$(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = (\mathbf{A} \cdot \mathbf{C})\mathbf{B} - (\mathbf{B} \cdot \mathbf{C})\mathbf{A}.$$

iv Scalar triple product

$$[\mathbf{a}, \mathbf{b}, \mathbf{c}] = \det(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$$

2 Matrix (矩陣)

I Matrix

A matrix is a rectangular array of numbers (or other mathematical objects), called the entries (元或元素) of the matrix. Most commonly, a matrix over a field F is a rectangular array of elements of F .

i Size

The size of a matrix is defined by the number of rows (zh-hant: 列/zh-hans: 行) and columns (zh-hant: 行/zh-hans: 列) it contains. A matrix with m rows and n columns is called an $m \times n$ matrix, or m -by- n matrix, where m and n are called its dimensions (階). A matrix with the same number of rows and columns n is called a n square matrix (方陣). A matrix with an infinite number of rows or/and columns is called an infinite matrix.

The set of all $m \times n$ matrices over $\mathbb{F}$ is commonly denoted as $\mathbb{F}^{m \times n}$.

ii Notation

The entry in the i -th row and j -th column, called the i, j th entry (第 i, j 元) or ij th entry (第 ij 元), is commonly denoted as $a_{i,j}$ or a_{ij} .

An $m \times n$ matrix $\mathbf{A}$ is commonly denoted as:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix},$$

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix},$$

$$\mathbf{A} = [a_{ij}]_{1 \leq i \leq m, 1 \leq j \leq n},$$

$$\mathbf{A} = (a_{ij})_{1 \leq i \leq m, 1 \leq j \leq n},$$

$$\mathbf{A} = [a_{ij}]_{m \times n},$$

or

$$\mathbf{A} = (a_{ij})_{m \times n}.$$

The i, i th entries are called diagonal entries (對角線元).

A matrix with only one row is called a row matrix or row vector, and a matrix with only one column is called a column matrix or column vector.

The i th row of the matrix $\mathbf{A}$ is commonly denoted as $\mathbf{A}_{i,:}$, and the j th column is commonly denoted as $\mathbf{A}_{:,j}$.

The matrix formed by merging two matrices $\mathbf{A}$ and $\mathbf{B}$ with the same number of rows left and right is denoted as:

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \end{bmatrix}$$

or

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} \end{pmatrix}$$

The matrix formed by merging two matrices $\mathbf{A}$ and $\mathbf{B}$ with the same number of columns up and down is denoted as:

$$\begin{bmatrix} \mathbf{A} \\ \mathbf{B} \end{bmatrix}$$

or

$$\begin{pmatrix} \mathbf{A} \\ \mathbf{B} \end{pmatrix}$$

II Matrix equality

We say that two matrices $\mathbf{A} = [a_{ij}]_{m \times n}$ and $\mathbf{B} = [b_{ij}]_{m \times n}$ are equal if

$$\forall 1 \leq i \leq m, 1 \leq j \leq n : a_{ij} = b_{ij}.$$

III Basic operations

i Matrix addition

The sum of two $m \times n$ matrices $\mathbf{A}$ and $\mathbf{B}$ is calculated entrywise.

ii Scalar multiplication

The product $c\mathbf{A}$ of a scalar c is calculated by multiplying every entry of $\mathbf{A}$ by c .

iii Subtraction

The subtraction of two $m \times n$ matrices $\mathbf{A}$ and $\mathbf{B}$ is defined as $\mathbf{A} - \mathbf{B} = \mathbf{A} + (-1)\mathbf{B}$.

iv Matrix multiplication (矩陣乘法)

Multiplication of two matrices is defined only if the number of columns of the left matrix is the same as the number of rows of the right matrix. The matrix product $\mathbf{AB}$ of an $m \times n$ matrix $\mathbf{A}$ and $n \times p$ matrix $\mathbf{B}$ is the $m \times p$ matrix whose ij th entry is the dot product of the i th row of $\mathbf{A}$ and the j th column of $\mathbf{B}$.

v Positive integer exponent

Given a square matrix $\mathbf{A}$ and a positive integer n , $\mathbf{A}^n$ is defined as $\prod_{i=1}^n \mathbf{A}$.

vi Zero matrix (零矩陣)

A zero matrix $\mathbf{O}$ over a field F is a matrix of which all entries are the additive identity of F , that is, let $m \times n$ zero matrix over F be $\mathbf{O}_{m \times n}$, for any matrices $\mathbf{A}_{m \times n}$ and $\mathbf{B}_{n \times p}$ over F , it satisfies:

$$\begin{aligned}\mathbf{A}_{m \times n} + \mathbf{O}_{m \times n} &= \mathbf{A}_{m \times n} \\ \mathbf{A}_{m \times n} - \mathbf{A}_{m \times n} &= \mathbf{O}_{m \times n} \\ \mathbf{O}_{m \times n} \mathbf{B}_{n \times p} &= \mathbf{O}_{m \times p} \\ \mathbf{A}_{m \times n} \mathbf{O}_{n \times p} &= \mathbf{O}_{m \times p}\end{aligned}$$

vii Identity matrix (單位矩陣)

An identity matrix $\mathbf{I}$ over a field F is a square matrix of which all diagonal entries are the multiplicative identity of F and all off-diagonal entries are the multiplicative absorbing element 0_F of F (i.e. $\forall x \in F: x \times 0_F = 0_F$), that is, let $n \times n$ identity matrix over F be $\mathbf{I}_n$, for any $m \times n$ matrices $\mathbf{A}$ over F , it satisfies:

$$\mathbf{A} \mathbf{I}_n = \mathbf{I}_m \mathbf{A} = \mathbf{A}$$

viii Kronecker product (克羅內克積)

If $\mathbf{A}$ is an $m \times n$ matrix and $\mathbf{B}$ is a $p \times q$ matrix, then the Kronecker product $\mathbf{A} \otimes \mathbf{B}$ is the $pm \times qn$ block matrix:

$$(\mathbf{A} \otimes \mathbf{B})_{i,j} = a_{[i/p], [j/q]} b_{(i-1) \bmod p+1, (j-1) \bmod q+1}$$

ix Hadamard product (哈達瑪乘積), Schur product (舒爾乘積), elementwise product, or entrywise product (逐項乘積)

For two $m \times n$ matrices $\mathbf{A}$ and $\mathbf{B}$, the Hadamard product $\mathbf{A} \odot \mathbf{B}$ or $\mathbf{A} \circ \mathbf{B}$ is an $m \times n$ matrix defined as:

$$(\mathbf{A} \odot \mathbf{B})_{ij} = (\mathbf{A})_{ij} (\mathbf{B})_{ij}.$$

x Property

For scalars r and s , matrices A, B, C, D , zero matrix O , and identity matrix I , assuming that all of the following operations hold, $\neq$ means not necessarily equals:

$$A + B = B + A$$

$$A + (B + C) = (A + B) + C$$

$$r(A + B) = rA + rB$$

$$rA + sA = (r + s)A$$

$$(rs)A = r(sA) = s(rA)$$

$$(AB)C = A(BC)$$

$$r(AB) = (rA)B = A(rB)$$

$$A(B + C) = AB + AC$$

$$(A + B)C = AC + BC$$

$$(A + B)(C + D) = AC + AD + BC + BD$$

$$AB \neq BA$$

$$(AB = AC \wedge A \neq O) \nRightarrow B = C$$

$$A^2 = O \nRightarrow A = O$$

$$AB = O \nRightarrow (A = O \vee B = O)$$

$$A^2 = I \iff A = A^{-1} \nRightarrow A = \pm I$$

$$AB = I \nRightarrow (\exists r \in \mathbb{R} : A = rI \wedge B = \frac{I}{r})$$

$$AB = B \nRightarrow A = I$$

$$ACB = ADB \nRightarrow C = D$$

$$A \odot B = B \odot A$$

$$A \odot (B \odot C) = (A \odot B) \odot C$$

$$A \odot (B + C) = A \odot B + A \odot C$$

xi Zero exponent

Given an $n \times n$ matrix $\mathbf{A}$ and $n \times n$ identity matrix $\mathbf{I}$, $\mathbf{A}^0$ is defined as $\mathbf{I}$.

IV Matrix row and column operations

i Row operations (zh-hant: 列/zh-hans: 行運算)

Including:

- Row addition (zh-hant: 列/zh-hans: 行相加): Adding a row to another.
- Row multiplication (zh-hant: 列/zh-hans: 行加倍): Multiplying all entries of a row by a non-zero constant.
- Row switching (zh-hant: 列/zh-hans: 行交換): Interchanging two rows of a matrix.

If one matrix can be transformed to another through a sequence of row operations, the two matrices are called row-equivalent.

ii (General) row echelon form ((廣義) zh-hant: 列/zh-hans: 行階梯形式)

A matrix is in row echelon form if:

- All rows having only zero entries are at the bottom.
- The leading entry, that is, the left-most non-zero entry, of every non-zero row is on the right of the leading entry of every row above.
- Some texts add the condition that the leading coefficient must be 1.

iii Column operations (zh-hant: 行/zh-hans: 列運算)

Including:

- Column addition (zh-hant: 行/zh-hans: 列相加): Adding a column to another.
- Column multiplication (zh-hant: 行/zh-hans: 列加倍): Multiplying all entries of a column by a non-zero constant.
- Column switching (zh-hant: 行/zh-hans: 列交換): Interchanging two columns of a matrix.

If one matrix can be transformed to another through a sequence of column operations, the two matrices are called column-equivalent.

iv Transpose (轉置)

The transpose of an $m \times n$ matrix $\mathbf{A}$ is the $n \times m$ matrix $\mathbf{A}^T$:

$$(\mathbf{A}^T)_{i,j} = a_{j,i}$$

v Symmetric matrix (對稱矩陣)

$\mathbf{A}$ is called a symmetric matrix if $\mathbf{A} = \mathbf{A}^T$.

vi Skew-symmetric matrix/antisymmetric matrix (反對稱矩陣)

$\mathbf{A}$ is called a skew-symmetric matrix or an antisymmetric matrix if $\mathbf{A} = -\mathbf{A}^T$.

vii Transpose identities

$$\left(\sum_{i=1}^n \mathbf{A}_i \right)^\top = \sum_{i=1}^n \mathbf{A}_i^\top$$
$$\left(\prod_{i=1}^n \mathbf{A}_i \right)^\top = \prod_{i=1}^n \mathbf{A}_{n-i+1}^\top$$

V Trace (跡)

The trace of a matrix $\mathbf{A}$ is the sum of all diagonal entries of $\mathbf{A}$, denoted as $\text{tr}(\mathbf{A})$.

i Conjugate transpose (共軛轉置)/Hermitian conjugate (埃爾米特共軛)/Hermitian transpose (埃爾米特轉置)

The conjugate transpose of an $m \times n$ matrix $\mathbf{A}$ is the $n \times m$ matrix $\mathbf{A}^*$:

$$(\mathbf{A}^*)_{i,j} = \overline{a_{j,i}}$$

VI Cofactor matrix, adjugate matrix, determinant, and inverse

Let $\mathbf{M}_{ij}$ of $\mathbf{A}$ be the determinant of the submatrix of $\mathbf{A}$ without the i th row and j th column.

i Determinant (行列式或判別式)

The determinant of a 1×1 matrix is the only element of it; the determinant of an $n \times n$ matrix $\mathbf{A}$, denoted $\det(\mathbf{A})$ or

$$\begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{vmatrix},$$

is

$$\det(\mathbf{A}) = \sum_{i=1}^n a_{ij}(-1)^{i+j} \mathbf{M}_{ij} \quad j \leq n \in \mathbb{N}$$

$$\det(\mathbf{A}) = \sum_{j=1}^n a_{ij}(-1)^{i+j} \mathbf{M}_{ij} \quad i \leq n \in \mathbb{N}$$

ii Determinant identities

Let $\mathbf{A}$ and $\mathbf{A}_i$ be square matrices.

$$\det\left(\sum_{i=1}^n \mathbf{A}_i\right) = \sum_{i=1}^n \det(\mathbf{A}_i)$$

$$\det\left(\prod_{i=1}^n \mathbf{A}_i\right) = \prod_{i=1}^n \det(\mathbf{A}_i)$$

$$\det(\mathbf{A}) \neq 0 \implies \det(\mathbf{A}^n) \neq 0$$

iii Cofactor matrix (餘因子矩陣)

The cofactor matrix of an $n \times n$ matrix $\mathbf{A}$ is the $n \times n$ matrix $\text{cof}(\mathbf{A})$:

$$[\text{cof}(\mathbf{A})]_{ij} = ((-1)^{i+j} \mathbf{M}_{ij})$$

iv Adjugate matrix (伴隨矩陣)/classical adjoint (經典伴隨矩陣)

The adjugate matrix or classical adjoint matrix of an $n \times n$ matrix $\mathbf{A}$ is the $n \times n$ matrix $\text{adj}(\mathbf{A})$:

$$\text{adj}(\mathbf{A}) = \text{cof}(\mathbf{A})^T$$

v Inverse (反方陣、逆方陣、反矩陣或逆矩陣), multiplicative inverse (乘法反方陣), or reciprocal

The inverse of an $n \times n$ matrix $\mathbf{A}$ is an $n \times n$ matrix $\mathbf{A}^{-1}$ such that:

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$$

vi Negative integer exponent

Given an invertible matrix $\mathbf{A}$ and a negative integer n , $\mathbf{A}^n$ is defined as $(\mathbf{A}^{-1})^n$.

vii Invertible matrix (可逆矩陣), orthogonal matrix (正交矩陣), orthonormal matrix (正則矩陣), non-singular matrix (非奇異矩陣), or non-degenerate matrix (非退化矩陣)

An invertible matrix, orthogonal matrix, orthonormal matrix, non-singular matrix, or non-degenerate matrix is a matrix such that its inverse exists.

Let $\mathbf{A}$ be an $n \times n$ matrix, the following prepositions are equivalent to " $\mathbf{A}$ is an invertible matrix":

$$\exists \mathbf{A}^{-1}$$

$$\exists (\mathbf{A}^T)^{-1}$$

$$\exists (\mathbf{A}^T \mathbf{A})^{-1}$$

$$\text{rank}(\mathbf{A}) = n$$

$\mathbf{A}$ 存在非零的特徵值

viii Non-invertible matrix (不可逆矩陣), non-orthogonal matrix (非正交矩陣), non-orthonormal matrix (非正則矩陣), singular matrix (奇異矩陣), or degenerate matrix (退化矩陣)

A non-invertible matrix, non-orthogonal matrix, non-orthonormal matrix, singular matrix, or degenerate matrix is a matrix that is not an invertible matrix.

ix Inverse of a two-by-two matrix

For an invertible 2×2 matrix $\mathbf{A}$:

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \Rightarrow \mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

x Computing inverse with row operations

Let $\mathbf{A}$ be an $n \times n$ matrix and $\mathbf{I}$ be the $n \times n$ identity matrix. If $(\mathbf{A} \ \mathbf{I})$ can become $(\mathbf{I} \ \mathbf{B})$ after row operations, then

$$\mathbf{B} = \mathbf{A}^{-1}$$

xi Unitary matrix (么正矩陣或酉矩陣)

An unitary matrix is a square matrix whose inverse equals to its conjugate transpose.

xii Determinant property

- When a row (or column) is added to another, the determinant remains unchanged.
- When all entries of a row (or column) is multiplied by a real number n , the determinant is multiplied by n .
- When two rows (or columns) are swapped, the determinant is multiplied by -1 .
- Given matrix $\mathbf{A}$ and its inverse $\mathbf{A}^{-1}$, $\det(\mathbf{A}^{-1}) = \frac{1}{\det(\mathbf{A})}$.

xiii Inverse identities

Let $\mathbf{A}$ and $\mathbf{A}_i$ be invertible matrices, $\mathbf{B}$ be a matrix of the same size of $\mathbf{A}$.

$$\mathbf{A}^{-1} = \frac{\text{adj}(\mathbf{A})}{\det(\mathbf{A})}$$

$$(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$$

$$\left(\prod_{i=1}^n \mathbf{A}_i \right)^{-1} = \prod_{i=1}^n (\mathbf{A}_{n-i+1})^{-1}$$

$$(\mathbf{A}^{-1} \mathbf{B} \mathbf{A})^n = \mathbf{A}^{-1} \mathbf{B}^n \mathbf{A}, \quad n \in \mathbb{N}$$

xiv Vandermonde matrix (范德蒙矩陣)

A Vandermonde matrix $\mathbf{V}$ is:

$$V_{i,j} = \alpha_i^{j-1}$$

An $n \times n$ Vandermonde matrix $\mathbf{V}$ satisfies:

$$\det(V) = \prod_{1 \leq i < j \leq n} (\alpha_j - \alpha_i)$$

VII Diagonal matrix (對角矩陣)

i Diagonal matrix

A diagonal matrix over a field F is a square matrix over F of which all non-diagonal entries are the additive identity of F .

ii Diagonal matrix identities

If:

$$\mathbf{D} = \begin{bmatrix} d_1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_n \end{bmatrix}$$

where all $d_i \neq 0$, then:

$$\mathbf{D}^{-1} = \begin{bmatrix} \frac{1}{d_1} & 0 & \cdots & 0 \\ 0 & \frac{1}{d_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{1}{d_n} \end{bmatrix}$$

Let $\mathbf{E}, \mathbf{F}$ be two $n \times n$ diagonal matrices:

$$\mathbf{EF} = [e_{ij}f_{ij}]_{1 \leq i,j \leq n}$$

VIII System of Linear Equations (線性方程組)

i System of equations (方程組)

- **Consistent system (相容方程組):** A system of equations is considered consistent if it has at least one solution.
- **Inconsistent/contradictory system (不相容方程組):** A system of equations is considered inconsistent or contradictory if it has no solution.
- **Dependent system (相依方程組):** A consistent system of equations is considered dependent if it has infinite solutions.
- **Independent system (獨立方程組):** A consistent system of equations is considered independent if it has finite solutions.

ii Linear Equation (線性方程)

A linear equation is an equation that may be put in the form:

$$\sum_{k=1}^n a_k x_k = c,$$

where a_k and c are called coefficients.

If $c = 0$, the linear equation is called a homogeneous linear equation; otherwise, it is called a heterogeneous linear equation.

iii System of Linear Equations (線性方程組)

A system of linear equations is a collection of two or more linear equations involving the same tuple of variables.

iv Coefficient matrix (係數矩陣)

The coefficients of a system of linear equations written as a matrix, with one row for each equation.

v Augmented matrix (增廣矩陣)

Coefficient matrix with the constants on the right side of the equal signs of a system of linear equations augmented to the last column of the matrix.

vi Gaussian Elimination (高斯消去法) or row reduction (zh-hant: 列/zh-hans: 行簡化)

An algorithm to solve a system of linear equations by using row operations to modify its augmented matrix to row echelon form. Two systems of linear equations have the same solution if and only if their augmented matrices are row-equivalent.

vii Cramer Rule or Cramer formula (克拉瑪公式)

Given a system of linear equations that can be written in the form:

$$\mathbf{A}\mathbf{X} = \mathbf{C}$$

where $\mathbf{A}$ is an $n \times n$ coefficient matrix, $\mathbf{X}$ is a variable matrix:

$$\mathbf{X} = (x_1, x_2, x_3, \dots, x_n)^T,$$

and $\mathbf{C}$ is a coefficient matrix:

$$\mathbf{C} = (c_1, c_2, c_3, \dots, c_n)^T.$$

Let $\mathbf{A}_i$ be $\mathbf{A}$ with the i th column replaced by $\mathbf{C}$. The solution to the system of linear equations is given by:

$$\begin{cases} x_i = \frac{\det(\mathbf{A}_i)}{\det(\mathbf{A})}, & \det(\mathbf{A}) \neq 0 \\ \text{inconsistent,} & \det(\mathbf{A}) = 0 \wedge \bigvee_{i=1}^n \det(\mathbf{A}_i) \neq 0 \\ \text{dependent,} & \det(\mathbf{A}) = 0 \wedge \bigvee_{i=1}^n \det(\mathbf{A}_i) = 0 \end{cases}$$

where $x_i = \frac{\det(\mathbf{A}_i)}{\det(\mathbf{A})}$ can be geometrically viewed as the volume of the parallelogram formed by the column vectors of $\mathbf{A}_i$ divided by the volume of the parallelogram formed by the column vectors of $\mathbf{A}$.

Particularly, when $\mathbf{C}$ is a zero matrix, that is,

$$\mathbf{A}\mathbf{X} = \mathbf{0},$$

the solution to the system of linear equations is given by:

$$\begin{cases} x_i = 0, & \det(\mathbf{A}) \neq 0 \\ \text{dependent,} & \det(\mathbf{A}) = 0 \end{cases}$$

3 Linear Subspace (線性子空間)

I Linear subspace

A linear subspace of a vector space V over a field $\mathbb{K}$ is a nonempty subset W of V such that,

$$\forall w_1, w_2 \in W, a, b \in \mathbb{K} : aw_1 + bw_2 \in W.$$

II Four fundamental subspaces of a matrix (矩陣的四大子空間)

The four fundamental subspaces of a matrix are the row space, column space, null space, and left null space.

III Column space (zh-hant: 行/zh-hans: 列空間)

The column space of $\mathbf{A}$, denoted as $\text{Col}(\mathbf{A})$ or $\mathbf{C}(\mathbf{A})$, is defined as the set of all linear combinations of the column vectors of $\mathbf{A}$.

IV Row space (zh-hant: 列/zh-hans: 行空間)

The row space of $\mathbf{A}$ is defined as the set of all linear combinations of the row vectors of $\mathbf{A}$, that is, $\text{Col}(\mathbf{A}^\top)$.

V Null space (零空間), kernel (核), or kernel space (核空間)

The kernel of a linear map $L : V \rightarrow W$, denoted as $\ker(L)$ or $\text{Null}(L)$, is defined as:

$$\ker(L) = \{v \in V : L(v) = 0\}.$$

$\dim(\ker(\mathbf{A}))$ is called nullity (零化度).

VI Left null space (左零空間), left kernel (左核), or left kernel space (左核空間)

The left null space of a linear map $L : V \rightarrow W$ is defined as $\ker(L^{-1})$.

VII Image space (像空間) or range (值域)

The image space of a linear map $T : V \rightarrow W$, denoted as $\text{im}(T)$

$$\text{im}(T) = \{T(v) : v \in V\}.$$

VIII Cokernel (餘核)

The cokernel of a linear map $T : V \rightarrow W$, denoted as $\text{coker}(T)$

$$\text{coker}(T) = W \setminus \text{im}(T).$$

IX Constant rank theorem (常秩定理)

For any matrix $\mathbf{A}$:

$$\dim(\text{Col}(\mathbf{A})) = \dim(\text{Col}(\mathbf{A}^\top))$$

Proof.

Let $\mathbf{A}$ be an $m \times n$ matrix, $r = \dim(\text{Col}(\mathbf{A}))$, $c_1, c_2, \dots, c_r$ be a basis of $\text{Col}(\mathbf{A})$, matrix $\mathbf{C} = [c_1, c_2, \dots, c_r]$, and $r \times n$ matrix $\mathbf{R}$ such that $\mathbf{A} = \mathbf{C}\mathbf{R}$.

Since:

$$\begin{aligned}\text{Col}(\mathbf{A}^\top) &\subseteq \text{Col}(\mathbf{R}^\top), \\ \dim(\text{Col}(\mathbf{A}^\top)) &\leq \dim(\text{Col}(\mathbf{R}^\top))\end{aligned}$$

Because there are only r rows in $\mathbf{R}$,

$$\dim(\text{Col}(\mathbf{R}^\top)) \leq \dim(\text{Col}(\mathbf{A}))$$

Similarly, we can obtain that

$$\dim(\text{Col}(\mathbf{A})) \leq \dim(\text{Col}(\mathbf{R}^\top)) \leq \dim(\text{Col}(\mathbf{A}^\top)).$$

Therefore,

$$\dim(\text{Col}(\mathbf{A})) = \dim(\text{Col}(\mathbf{R}^\top)) = \dim(\text{Col}(\mathbf{A}^\top)).$$

□

X Rank (秩)

If $\mathbf{A}$ is an $m \times n$ matrix, then the rank of $\mathbf{A}$, denoted as $\text{rank}(\mathbf{A})$ or $\text{rk}(\mathbf{A})$, is the maximum number of linearly independent columns (or rows) in $\mathbf{A}$, namely:

$$\text{rank}(\mathbf{A}) = \dim(\text{Col}(\mathbf{A}))$$

XI Rank-nullity theorem (秩—零化度定理)

For any $n \times n$ matrix $\mathbf{A}$:

$$\text{rank}(\mathbf{A}) + \dim(\ker(\mathbf{A})) = n$$

Proof.

Let $\mathbf{A}$ be an $m \times n$ matrix with r linearly independent columns (i.e., $\text{rank}(\mathbf{A}) = r$). We will show that there exists a set of $n - r$ linearly independent solutions to the homogeneous system $\mathbf{A}\mathbf{x} = \mathbf{0}$.

To do this, we will produce an $n \times (n - r)$ matrix $\mathbf{X}$ whose columns form a basis of the null space of $\mathbf{A}$.

Without loss of generality, assume that the first r columns of $\mathbf{A}$ are linearly independent. So, we can write

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 \end{pmatrix},$$

where

- $\mathbf{A}_1$ is an $m \times r$ matrix with r linearly independent column vectors, and
- $\mathbf{A}_2$ is an $m \times (n - r)$ matrix such that each of its $n - r$ columns is a linear combination of the columns of $\mathbf{A}_1$.

This means that $\mathbf{A}_2 = \mathbf{A}_1 \mathbf{B}$ for some $r \times (n - r)$ matrix $\mathbf{B}$, and hence,

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1 \mathbf{B} \end{pmatrix}.$$

Let

$$\mathbf{X} = \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix},$$

where $\mathbf{I}_{n-r}$ is the $(n - r) \times (n - r)$ identity matrix. So, $\mathbf{X}$ is an $n \times (n - r)$ matrix such that

$$\mathbf{A}\mathbf{X} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1 \mathbf{B} \end{pmatrix} \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} = -\mathbf{A}_1 \mathbf{B} + \mathbf{A}_1 \mathbf{B} = \mathbf{0}_{m \times (n-r)}.$$

Therefore, each of the $n - r$ columns of $\mathbf{X}$ are particular solutions of $\mathbf{A}\mathbf{x} = \mathbf{0}_{\mathbb{F}^m}$.

Furthermore, the $n - r$ columns of $\mathbf{X}$ are linearly independent because $\mathbf{X}\mathbf{u} = \mathbf{0}_{\mathbb{F}^n}$ will imply $\mathbf{u} = \mathbf{0}_{\mathbb{F}^{n-r}}$ for $\mathbf{u} \in \mathbb{F}^{n-r}$:

$$\mathbf{X}\mathbf{u} = \mathbf{0}_{\mathbb{F}^n} \implies \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} \mathbf{u} = \mathbf{0}_{\mathbb{F}^n} \implies \begin{pmatrix} -\mathbf{B}\mathbf{u} \\ \mathbf{u} \end{pmatrix} = \begin{pmatrix} \mathbf{0}_{\mathbb{F}^r} \\ \mathbf{0}_{\mathbb{F}^{n-r}} \end{pmatrix} \implies \mathbf{u} = \mathbf{0}_{\mathbb{F}^{n-r}}.$$

Therefore, the column vectors of $\mathbf{X}$ constitute a set of $n - r$ linearly independent solutions for $\mathbf{A}\mathbf{x} = \mathbf{0}_{\mathbb{F}^m}$.

We next prove that any solution of $\mathbf{A}\mathbf{x} = \mathbf{0}_{\mathbb{F}^m}$ must be a linear combination of the columns of $\mathbf{X}$.

Let

$$\mathbf{u} = \begin{pmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{pmatrix} \in \mathbb{F}^n$$

be any vector such that $\mathbf{A}\mathbf{u} = \mathbf{0}_{\mathbb{F}^m}$. Since the columns of $\mathbf{A}_1$ are linearly independent, $\mathbf{A}_1 \mathbf{x} = \mathbf{0}_{\mathbb{F}^m}$ implies $\mathbf{x} = \mathbf{0}_{\mathbb{F}^r}$.

Therefore,

$$\begin{aligned} \mathbf{A}\mathbf{u} &= \mathbf{0}_{\mathbb{F}^m} \\ \implies \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1 \mathbf{B} \end{pmatrix} \begin{pmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{pmatrix} &= \mathbf{A}_1 \mathbf{u}_1 + \mathbf{A}_1 \mathbf{B} \mathbf{u}_2 = \mathbf{A}_1 (\mathbf{u}_1 + \mathbf{B} \mathbf{u}_2) = \mathbf{0}_{\mathbb{F}^m} \\ \implies \mathbf{u}_1 + \mathbf{B} \mathbf{u}_2 &= \mathbf{0}_{\mathbb{F}^r} \\ \implies \mathbf{u}_1 &= -\mathbf{B} \mathbf{u}_2 \\ \implies \mathbf{u} &= \begin{pmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{pmatrix} = \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} \mathbf{u}_2 = \mathbf{X} \mathbf{u}_2. \end{aligned}$$

This proves that any vector $\mathbf{u}$ that is a solution of $\mathbf{A}\mathbf{x} = \mathbf{0}$ must be a linear combination of the $n - r$ special solutions given by the columns of $\mathbf{X}$. And we have already seen that the columns of $\mathbf{X}$ are linearly independent. Hence, the columns of $\mathbf{X}$ constitute a basis for the null space of $\mathbf{A}$. Therefore, the nullity of $\mathbf{A}$ is $n - r$. Since r equals the rank of $\mathbf{A}$, it follows that $\text{rank}(\mathbf{A}) + \text{ker}(\mathbf{A}) = n$. This concludes our proof. $\square$

XII Full rank (滿秩) and rank deficiency (秩虧)

For an $m \times n$ matrix $\mathbf{A}$, if

$$\text{rank}(\mathbf{A}) = \min(m, n)$$

$\mathbf{A}$ is called to have full rank; otherwise it is called to be rank-deficient (欠秩或秩虧的).

The rank deficiency of a $m \times n$ matrix $\mathbf{A}$ is defined as

$$\min(m, n) - \text{rank}(\mathbf{A})$$

XIII Property of rank

- For any $m \times n$ matrix $\mathbf{A}$:

$$\text{rank}(\mathbf{A}) \leq \min(m, n)$$

- A square matrix is invertible if and only if it has full rank.
- A matrix is a zero matrix if and only if its rank is zero.

4 Matrix Analysis (矩陣分析)

I Eigenvector or characteristic vector (特徵向量) and characteristic polynomial (特徵多項式)

i Eigenvector or characteristic vector (特徵向量) and eigenvalue (本徵值、固有值、特徵值或特徵根)

Given a linear transformation $L: V \rightarrow V$, an eigenvector of it is defined as the vector $v \in V$ such that:

$$\mathbf{A}v = \lambda v,$$

where λ is a scalar, which is called the eigenvalue of $\mathbf{A}$ or the eigenvalue of v .

ii Characteristic polynomial

For an $n \times n$ matrix $\mathbf{A}$, the characteristic polynomial of it is defined as

$$p_{\mathbf{A}}(\lambda) := \det(\lambda \mathbf{I}_n - \mathbf{A}),$$

where λ is a scalar variable.

A scalar λ is an eigenvalue of $\mathbf{A}$ if and only if

$$p_{\mathbf{A}}(\lambda) = 0.$$

iii Property

- The number of distinct eigenvalues of a square matrix is less than or equal to the number of linearly independent eigenvectors.
- The number of distinct eigenvalues of a $n \times n$ matrix is less than or equal to n .
- The algebraic multiplicity of a root λ of the characteristic polynomial of a square matrix is greater than or equal to the number of linearly independent eigenvectors of it whose eigenvalue is λ .

iv Cayley–Hamilton theorem (凱萊–哈密頓定理)

For a square matrix $\mathbf{A}$ and its characteristic polynomial $p_{\mathbf{A}}$, the Cayley–Hamilton theorem states that:

$$p_{\mathbf{A}}(\mathbf{A}) = 0$$

v Diagonalization (對角化)

The diagonalization of an $n \times n$ matrix $\mathbf{A}$ is the construction of an $n \times n$ matrix $\mathbf{P}$ and an $n \times n$ diagonal matrix $\mathbf{D}$ such that:

$$\mathbf{A} = \mathbf{P}^{-1} \mathbf{D} \mathbf{P}$$

An $n \times n$ matrix can be diagonalized if it has n distinct eigenvalues.

The steps of the diagonalization of an $n \times n$ matrix $\mathbf{A}$ is as follows:

1. For each eigenvalue λ of the matrix $\mathbf{A}$, solve the eigenvector $\mathbf{v}$:

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = 0.$$

2. Compose matrix $\mathbf{P}$ with each eigenvector of $\mathbf{A}$ as a row.

Common application:

$$\mathbf{A}^n = (\mathbf{P}^{-1} \mathbf{D} \mathbf{P})^n = \mathbf{P}^{-1} \mathbf{D}^n \mathbf{P}$$

vi Definiteness (正定性)

Let $A \in \mathbb{R}^{n \times n}$ and $\mathbf{x} \in \mathbb{R}^n$. A is called

- Positive definite: if

$$\mathbf{x}^T \mathbf{A} \mathbf{x} > 0, \quad \forall \mathbf{x} \neq 0.$$

- Negative definite: if

$$\mathbf{x}^T \mathbf{A} \mathbf{x} < 0, \quad \forall \mathbf{x} \neq 0.$$

- Positive semidefinite: if

$$\mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0, \quad \forall \mathbf{x},$$

$$\mathbf{x}^T \mathbf{A} \mathbf{x} > 0, \quad \exists \mathbf{x}.$$

- Negative semidefinite: if

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} \leq 0, \quad \forall \mathbf{x},$$

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} < 0, \quad \exists \mathbf{x}.$$

- Neutral/Zero semidefinite: if

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} = 0, \quad \forall \mathbf{x}.$$

- Indefinite: otherwise.

vii Eigenvalues Test of Definiteness for Symmetric Hessians of Scalar Fields

For a symmetric matrix A ,

- A is positive definite if all eigenvalues of it are positive.
- A is negative definite if all eigenvalues of it are negative.
- A is positive semidefinite if all eigenvalues of it are nonnegative and at least one eigenvalue of it is positive.
- A is negative semidefinite if all eigenvalues of it are nonpositive and at least one eigenvalue of it is negative.
- A is neutral semidefinite if all eigenvalues of it are zero.
- A is indefinite otherwise.

II Generalized Definition of Inverse

i One-sided inverse (單邊逆)

- Right inverse (右逆): For any $n \times m$ matrix A with rank $\text{rank}(A) = n$, there exists an $m \times n$ matrix A_R^{-1} , called the right inverse of $\mathbf{A}$, such that

$$A A_R^{-1} = I_n,$$

where I_n is the $n \times n$ identity matrix.

- Left inverse (左逆): For any $n \times m$ matrix A with rank $\text{rank}(A) = m$, there exists an $m \times n$ matrix A_L^{-1} , called the left inverse of $\mathbf{A}$, such that

$$A_L^{-1} A = I_m,$$

where I_m is the $m \times m$ identity matrix.

Properties:

- A matrix doesn't necessarily have a right inverse and doesn't necessarily have a left inverse.
- A square matrix has a right inverse if and only if it has left inverse if and only if it is invertible.

- For any invertible matrix, there exists only one right inverse and one left inverse, which are its multiplicative inverse.

ii Generalized inverse or g-inverse (廣義逆)

A matrix $A^g \in \mathbb{C}^{n \times m}$ is a generalized inverse of a matrix $A \in \mathbb{C}^{m \times n}$ if:

$$AA^gA = A.$$

Properties:

- For all matrices, there exists at least one generalized inverse, but it's not necessarily unique.
- For any invertible matrix, there exists only one generalized inverse, which is its multiplicative inverse.

iii Reflexive generalized inverse (自反廣義逆)

A reflexive generalized inverse of a matrix A is defined as a matrix A^g such that:

$$AA^gA = A$$

$$A^gAA^g = A^g$$

Properties:

- For all matrices, there exists at least one reflexive generalized inverse, but it's not necessarily unique.
- For any invertible matrix, there exists only one reflexive generalized inverse, which is its multiplicative inverse.

iv Moore–Penrose inverse (摩爾—彭若斯廣義逆) or pseudoinverse (偽逆或擬反)

A Moore–Penrose inverse of a matrix A is defined as a matrix A^+ such that:

$$AA^+A = A$$

$$A^+AA^+ = A^+$$

$$(AA^+)^* = AA^+$$

$$(A^+A)^* = A^+A$$

Properties:

- For all matrices, there exists only one Moore–Penrose inverse.
- For any invertible matrix, there exists only one Moore–Penrose inverse, which is its multiplicative inverse.

5 Transformation Matrices

I Groups

i General linear group

The general linear group of degree n over some field F is the set of $n \times n$ invertible matrices over F , together with the operation of matrix multiplication, denoted as $GL(n, F)$ or $GL_n(F)$, or simply $GL(n)$ if the field is understood. It is a topological group and a Lie group under operation $+$.

The Lie algebra of a general linear group is denoted as the symbol of the general linear group with GL replaced with $\mathfrak{gl}$.

ii Orthogonal group or general orthogonal group

The orthogonal group or general orthogonal group of degree n over some field F is the set of $n \times n$ invertible matrices over F such that the inverse equals the transpose, together with the operation of matrix multiplication, denoted as $O(n, F)$ or $O_n(F)$, or simply $O(n)$ if the field is understood. An element of an orthogonal group is called an orthogonal matrix. It is a topological group and a Lie group under operation $+$.

iii Special orthogonal group or rotation group

The special orthogonal group or rotation group of degree n over some field F is a subgroup of the orthogonal group of degree n over some field F such that for any matrix in it, the determinant is 1, that is, it preserves the orientation, denoted as $SO(n, F)$ or $SO_n(F)$, or simply $SO(n)$ if the field is understood. An element of a special orthogonal group is called an rotational matrix. It is a topological group and a Lie group under operation $+$.

The Lie algebra of a special orthogonal group is denoted as the symbol of the special orthogonal group with SO replaced with $\mathfrak{so}$.

II Transformation matrices in two-dimensional space

i Degenerate matrices

Degenerate to 1 dimension ($(a \vee b) \wedge (d \vee e) \neq 0$):

$$\begin{bmatrix} a & b \\ ca & cb \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ax + by \\ c(ax + by) \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ d & e \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ dx + ey \end{bmatrix}$$

Degenerate to 0 dimension:

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

ii Stretching (伸縮/縮放/拉伸)

A stretch is a linear transformation which enlarges all distances in a particular direction by a constant factor k but does not affect distances in the perpendicular direction. (Note that if $k > 1$, then this really is a "stretch"; if $k < 1$, it is technically a "compression", but we still call it a stretch. Also, if $k = 1$, then the transformation is an identity, i.e. it has no effect)

The matrix associated with a stretch by a factor h along the x -axis and a factor k along the y -axis is given by:

$$\begin{bmatrix} h & 0 \\ 0 & k \end{bmatrix}$$

iii Shearing (推移)

A shearing in the x -direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$$

A shearing in the y -direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$$

iv Rotation (旋轉)

A rotation by an angle θ counterclockwise about the origin is given by:

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

v Reflection (鏡射/反射)

A reflection about a line that passes the origin and with $(a, b) = \sqrt{a^2 + b^2}(\cos \theta, \sin \theta)$ being a vector in the direction of it is given by:

$$\begin{bmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{bmatrix} = \frac{1}{a^2 + b^2} \begin{bmatrix} a^2 - b^2 & 2ab \\ 2ab & b^2 - a^2 \end{bmatrix}$$

vi Orthogonal projection (正射影)

An orthogonal projection about a line that passes the origin and with $(a, b) = \sqrt{a^2 + b^2}(\cos \theta, \sin \theta)$ being a vector in the direction of it is given by:

$$\begin{bmatrix} \cos^2(\theta) & \cos(\theta)\sin(\theta) \\ \cos(\theta)\sin(\theta) & \sin^2(\theta) \end{bmatrix} = \frac{1}{a^2 + b^2} \begin{bmatrix} a^2 & ab \\ ab & b^2 \end{bmatrix}$$

vii Exmample 1

Question: The cricle $C : x^2 + y^2 = 1$ is transformed by $\mathbf{A} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ to obtain a new curve C' . Find the equation of C' and its area.

Answer:

$$\begin{aligned} \begin{bmatrix} x' \\ y' \end{bmatrix} &= \mathbf{A} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ 3y \end{bmatrix} \\ \begin{cases} x' = 2x \\ y' = 3y \end{cases} &\Rightarrow \begin{cases} x = \frac{x'}{2} \\ y = \frac{y'}{3} \end{cases} \\ \left(\frac{x'}{2}\right)^2 + \left(\frac{y'}{3}\right)^2 = 1 &\Rightarrow \frac{x'^2}{4} + \frac{y'^2}{9} = 1 \\ C' : \frac{x'^2}{4} + \frac{y'^2}{9} &= 1 \end{aligned}$$

The area of C' is $\pi \cdot 2 \cdot 3 = 6\pi$. Since $\det(\mathbf{A}) = 6$, and the area of C is π , the area of C' equals $\det(\mathbf{A})$ multiplied by the area of C .

viii Exmample 2

Question: The line $\mathbf{L} : 4x - 3y = 5$ is transformed in the x -direction dependent on the y coordination by twice and then in the y -direction dependent on the x coordination by negative threee times to obtain a new line $\mathbf{L}'$. Find the equation of $\mathbf{L}'$.

Answer:

$$\begin{aligned} \mathbf{A} &= \begin{bmatrix} 1 & 2 \\ -3 & 1 \end{bmatrix} \\ \mathbf{A}^{-1} &= \frac{1}{7} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix} \\ \begin{bmatrix} x \\ y \end{bmatrix} &= \mathbf{A}^{-1} \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \frac{x' - 2y'}{7} \\ \frac{3x' + y'}{7} \end{bmatrix} \\ \mathbf{L}' : 4\frac{x - 2y}{7} + 3\frac{3x + y}{7} &= 5 \\ \mathbf{L}' : 5x + 11y + 35 &= 0 \end{aligned}$$

ix Example 3

Question: Reflect the point $P(5, 2)$ over the line $\mathbf{M} : x - y + 1 = 0$ to obtain a new point Q . Find Q .

Answer:

Shift one unit to the right: $\mathbf{M}$ becomes $\mathbf{M}' : x - y = 0$ and C becomes $C'(6, 2)$.

The reflection matrix is:

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 1 & 0 \end{bmatrix}$$

$$D' = \mathbf{AC}' = (2, 6)$$

Shift one unit to the left: $D(1, 6)$

x Example 4

Question: The line $x + y - 2 = 0$ is transformed by $\mathbf{A} = \begin{bmatrix} a & b \\ 1 & 2 \end{bmatrix} \in \mathbb{R}^{2 \times 2}$ to obtain a new line $2x + 3y - 4 = 0$. Find a and b .

Answer:

$$\mathbf{A} \begin{bmatrix} x \\ -x + 2 \end{bmatrix} = \begin{bmatrix} x' \\ \frac{-2x' + 4}{3} \end{bmatrix}$$

$$x - 2x + 4 = \frac{-2x' + 4}{3}$$

$$x' = \frac{3}{2}x - 4$$

$$ax - bx - 2b = x' = \frac{3}{2}x - 4$$

$$(a, b) = \left(-\frac{1}{2}, -2\right)$$

III Transformation matrices in three-dimensional space

i Stretching (伸縮/縮放/拉伸)

The matrix associated with a stretch by a factor h along the x -axis, a factor k along the y -axis, and a factor w along the z -axis is given by:

$$\begin{bmatrix} h & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & w \end{bmatrix}$$

ii Shearing (推移)

A shearing in the x -direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & k & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the x -direction dependent on the z coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & k \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the y -direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ k & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the y -direction dependent on the z coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & k \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the z -direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ k & 0 & 1 \end{bmatrix}$$

A shearing in the z -direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & k & 1 \end{bmatrix}$$

iii Rotation (旋轉)

A rotation by an angle θ counterclockwise about a line that passes the origin and with (a, b, c) being a vector in the direction of it is given by:

$$\frac{1}{a^2 + b^2 + c^2} \begin{bmatrix} a^2(1 - \cos \theta) + \cos \theta & ab(1 - \cos \theta) - c \sin \theta & ac(1 - \cos \theta) + b \sin \theta \\ bc(1 - \cos \theta) + c \sin \theta & b^2(1 - \cos \theta) + \cos \theta & bc(1 - \cos \theta) - a \sin \theta \\ ac(1 - \cos \theta) - b \sin \theta & bc(1 - \cos \theta) + a \sin \theta & c^2(1 - \cos \theta) + \cos \theta \end{bmatrix}$$

iv Rodrigues' rotation formula

If $\mathbf{v}, \mathbf{k} \in \mathbb{R}^3$ and $\mathbf{k}$ is a unit vector describing an axis of rotation about which $\mathbf{v}$ rotates by an angle θ according to the right hand rule, the Rodrigues formula for the rotated vector $\mathbf{v}_r$ is

$$\mathbf{v}_r = \mathbf{v} \cos \theta + (\mathbf{k} \times \mathbf{v}) \sin \theta + \mathbf{k}(\mathbf{k} \cdot \mathbf{v})(1 - \cos \theta).$$

The skew-symmetric matrix $\mathbf{K}$ in Lie algebra $\mathfrak{so}(3)$ that generates the one-parameter subgroup of $SO(3)$:

$$\mathbf{R}(\theta) = \exp(\theta \mathbf{K}),$$

such that

$$\mathbf{v}_r = \mathbf{R}(\theta) \mathbf{v},$$

is given by

$$\mathbf{K} \mathbf{v} = \mathbf{k} \times \mathbf{v}, \quad \forall \mathbf{v} \in \mathbb{R}^3,$$

and

$$\mathbf{R}(\theta) = \mathbf{I} + \sin \theta \mathbf{K} + (1 - \cos \theta) \mathbf{K}^2,$$

where $\mathbf{I}$ is the 3×3 identity matrix.

v Reflection (鏡射/反射)

A reflection about a plane $ax + by + cz = 0$ is given by:

$$\mathbf{I} - \frac{1}{a^2 + b^2 + c^2} \begin{bmatrix} 2a^2 & 2ab & 2ac \\ 2ab & 2b^2 & 2bc \\ 2ac & 2bc & 2c^2 \end{bmatrix},$$

where $\mathbf{I}$ is the 3×3 identity matrix.

vi Orthogona projection (正射影)

An orthogonal projection about a plane $ax + by + cz = 0$ is given by:

$$\mathbf{I} - \frac{1}{a^2 + b^2 + c^2} \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$$

where $\mathbf{I}$ is the 3×3 identity matrix.