Linear Algebra and Functional Analysis

沈威宇

October 15, 2026

Contents

Chapter 1 Linear Algebra (線性代數) and Functional Analysis (泛函分析) 1

1 Vector Space and Functional 1

I General 1

i Vector space (向量空間) or Linear space (線性空間). 1

ii Linear combination (線性組合) 2

iii Isomorphism (同構) 2

iv Division 2

v Affine space (仿射空間) 2

vi Linear map (線性映射) or linear operator 3

vii Linear transformation (線性變換) 3

viii Invertible map (可逆映射) 3

ix Linear subspace, vector subspace, or subspace 3

x Linear span, linear hull, or span 3

xi In the span iff new span unchanged 3

xii Linear independence (線性獨立) 4

xiii Linearly dependent set condition 4

xiv Linear independent set as minimum span generating set by inclusion of span 4

xv Spanning set element exchange theorem 4

xvi Steinitz exchange lemma 5

xvii Basis (基) 5

xviii Orientation 5

xix Dimension (維度) 5

xx Basis extension theorem 5

xxi Standard (unit/basis) vector 6

xxii Bilinear form (雙線性形式) 6
xxiii Symmetric bilinear form (對稱雙線性形式) 6
xxiv Topological vector space (TVS) (拓樸向量空間) 6
xxv Functional (泛函) 6
xxvi Seminorm (半範數) 6
xxvii Norm (範數/模長) 7
xxviii Sublinear function (亞線性函數) or quasi-seminorm (準半範數) 7
xxix Operator norm or induced norm 7
xxx Absolutely continuous function (絕對連續函數) 7
xxxi Bounded linear operator (有界線性運算子) 8
xxxii Uniform norm, sup norm, supremum norm, Chebyshev norm, infinity norm, or max norm 8
xxxiii Hahn–Banach theorem (哈恩-巴拿赫定理) 8
xxxiv p-norm (p-範數) 8
xxxv Locally convex topological vector space (LCTVS) or locally convex space . 9
xxxvi Banach space (巴拿赫空間) 9
xxxvii Dot product (點積) 9
xxxviii Inner product space (內積空間) 9
xxxix Inner product is bilinear form 10
xl Orthonormal basis (正交基) 10
xli Balanced set (平衡集), circled set, or disk set 10
xlii Well-defined (定義良好)/Unambiguous (不模糊的) 10
xliii Affine map (仿射映射) 10
xliv Affine transformation (仿射變換) 10
xlv Von Neumann bounded or bounded 10
xlvi Convex set (凸集) 10
xlvii (Algebraic) dual (對偶) (vector) (space) 11
xlviii Continuous dual (space) 11
xlix Transpose, dual map, or adjoint 11
l Strong dual (space) 11
li Orthogonal complement (正交補) 11
lii Affine combination 11

liii Convex combination 12
liv Coordinate system (座標/坐標系統) 12
lv Projection (投影) 12
lvi Orthographic projection (正射影) 12
lvii Hilbert space (希爾伯特空間) 12
lviii Angle (夾角) between two vectors 12
lix Principal angles (主夾角) between two subspaces 12
lx Flat, hyperplane, or affine subspace (仿射子空間) 13
lxi Hypersurface 13
lxii Parallel (平行) 13
lxiii Coplanar 13
lxiv Elementwise multiplication 13
lxv Unit vector (單位向量) or normalized vector 13
lxvi Affine independence 13
lxvii Simplex 13
lxviii Gram matrix, Gramian matrix, or Gramian 14
lxix Support of functional 14
lxx Euclidean vector space 14
lxxi Euclidean affine space (歐幾里德仿射空間) or Euclidean space (歐幾里得空間) 14
lxxii Affine hull (仿射包) 14
lxxiii Convex hull (凸包) 14
lxxiv Cartesian coordinate system (笛卡爾座標系統) or rectangular coordinate system (直角座標系統) 15
lxxv Lattice points (格子點) 15
lxxvi Curvilinear coordinate system 15
lxxvii Standard, unit, or probability simplex 15
lxxviii Rectangle 15
II Tensor 15
i Tensor 15
ii Upper and lower indices 16
III Two dimensional Euclidean vector space 16
i Quadrant (象限) 16

IV Three dimensional Euclidean vector space . . . 17
i Right-hand rule (右手定則) . . . 17
ii Octant (卦限) . . . 17
iii Cross product (叉積), external product (外積), or vector product (向量積) . . . 17
iv Basic properties of cross product . . . 17
v Triple product identity (三重積恆等式)/Jacobi identity (雅可比恆等式) . . . 17
vi Scalar triple product . . . 18
2 Matrix (矩陣) . . . 18
I Matrix . . . 18
II Basic operations . . . 19
i Submatrix . . . 19
ii Matrix equality . . . 19
iii Matrix addition . . . 19
iv Scalar multiplication . . . 19
v Matrix multiplication (矩陣乘法) . . . 19
vi Outer product . . . 19
vii Positive integer exponent . . . 19
viii Zero matrix (零矩陣) . . . 19
ix Identity matrix (單位矩陣) . . . 19
x Hadamard product (哈達瑪乘積), Schur product (舒爾乘積), elementwise product, or entrywise product (逐項乘積) . . . 20
xi Kronecker product (克羅內克積) . . . 20
xii Merging matrices . . . 20
xiii Property . . . 20
xiv Zero exponent . . . 22
xv Trace (跡) . . . 22
III Matrix row and column operations . . . 23
i Elementary row operations (zh-hant: 列/zh-hans: 行運算) and elementary column operations (zh-hant: 行/zh-hans: 列運算) . . . 23
ii Row operation as matrix multiplication and elementary matrix . . . 23
iii Zero row, nonzero row, and leading entry . . . 23
iv Transpose (轉置) . . . 24

v Symmetric matrix (對稱矩陣) . . . 24
vi Skew-symmetric matrix/antisymmetric matrix (反對稱矩陣) . . . 24
vii Transpose identities. . . . 24
IV Complex matrix . . . 25
i (Complex）conjugate. . . . 25
ii Conjugate transpose (共軛轉置)/Hermitian conjugate (埃爾米特共軛)/Hermitian transpose (埃爾米特轉置) . . . 25
iii Conjugate transpose identities . . . 26
V Cofactor matrix, adjugate matrix, determinant, and inverse . . . 26
i Determinant (行列式或判別式) . . . 26
ii Determinant identities . . . 26
iii Cofactor matrix (餘因子矩陣) . . . 27
iv Adjugate matrix (伴隨矩陣)/classical adjoint (經典伴隨矩陣) . . . 27
v Inverse (反方陣、逆方陣、反矩陣或逆矩陣), multiplicative inverse (乘法反方陣), or reciprocal. . . . 27
vi Negative integer exponent. . . . 27
vii Invertible matrix (可逆矩陣), orthogonal matrix (正交矩陣), orthonormal matrix (正則矩陣), non-singular matrix (非奇異矩陣), or non-degenerate matrix (非退化矩陣) . . . 27
viii Non-invertible matrix (不可逆矩陣), non-orthogonal matrix (非正交矩陣), non-orthonormal matrix (非正則矩陣), singular matrix (奇異矩陣), or degenerate matrix (退化矩陣). . . . 27
ix Inverse of a two-by-two matrix . . . 27
x Unitary matrix (么正矩陣或酉矩陣) . . . 27
xi Determinant property . . . 28
xii Inverse identities . . . 28
xiii Inverse of elementary matrix . . . 28
xiv Vandermonde matrix (范德蒙矩陣) . . . 29
VI Diagonal matrix . . . 29
i Diagonal matrix (對角矩陣) . . . 29
ii Diagonal matrix identities . . . 29
VII System of Linear Equations (線性方程組) . . . 29
i System of equations (方程組) . . . 29
ii Linear Equation (線性方程) . . . 30

iii System of Linear Equations (線性方程組) or linear system 30
iv Independent system of linear equations has exactly one solution 30
v Dependent system general solution 30
vi Coefficient matrix (係數矩陣) 30
vii Constant term vector 30
viii Augmented matrix (增廣矩陣) 30
ix Row equivalent means same solutions 31
x Cramer Rule or Cramer formula (克拉瑪公式) 31
VIII Row echelon form 32
i (General) row echelon form (REF) (（廣義）zh-hant: 列/zh-hans: 行階梯形式) . 32
ii Reduced row echelon form (RREF) 32
iii Pivot position 32
iv Gaussian Elimination (高斯消去法), row reduction, Gauss-Jordan Elimination, and reduced row echelon form test for consistency and obtaining general solution 33
IX Subspaces of matrices and linear maps 34
i Four fundamental subspaces of a matrix (矩陣的四大子空間) 34
ii Column space (zh-hant: 行/zh-hans: 列空間) 34
iii Row space (zh-hant: 列/zh-hans: 行空間) 34
iv Null space (零空間), kernel (核), or kernel space (核空間) 34
v Left null space (左零空間), left kernel (左核), or left kernel space (左核空間) . 34
vi Image space (像空間) or range (值域) 34
vii Cokernel (餘核) 34
X Rank and nullity 34
i Rank (秩) 34
ii Constant rank theorem (常秩定理) 35
iii Rank-nullity theorem (秩—零化度定理) 35
iv Rank-left nullity theorem 36
v Full rank (滿秩) and rank deficiency (秩虧) 36
vi Zero rank 37
vii Rank preservation 37
XI Properties 37
i Column correspondence property 37

ii Elementary row operation preservations . . . 37
iii Row-equivalence is an equivalence relation . . . 38
iv Rank and number of pivot columns . . . 38
v Nullity and number of non-pivot columns . . . 39
vi Left nullity and number of non-pivot rows . . . 39
vii Existence and uniqueness of reduced row echelon form . . . 39
viii Column space test for consistency of linear system . . . 39
ix Invertible coefficient matrix . . . 40
x Solution space is an affine space with kernel as associated vector space . . . 40
xi Zero nullity . . . 40
xii Pivot column set is column space basis . . . 41
xiii Invertible iff full rank . . . 41
xiv Invertible matrix reduced row echelon form is identity matrix . . . 41
xv Invertible matrix must be product of elementary matrix . . . 41
xvi Algorithm for computing inverse with elementary row operations . . . 41
xvii Algorithm for computing inverse right multiplied by another matrix with elementary row operations . . . 41
XII Partitioning . . . 42
i Partitioning . . . 42
ii Block multiplication . . . 42
iii Matrix multiplication by rows . . . 42
iv Matrix multiplication by outer product . . . 42
XIII Lower-upper (LU) decomposition or factorization . . . 42
i Lower triangular matrix . . . 42
ii Upper triangular matrix . . . 42
iii Lower triangular square matrices product . . . 42
iv Upper triangular square matrices product . . . 43
v Lower-upper (LU) decomposition or factorization with partial pivoting . . . 43
vi Lower-upper (LU) decomposition or factorization with partial pivoting . . . 44
vii Solving a system of linear equations using an LU decomposition . . . 44
viii Solving a system of linear equations using an LU decomposition with partial pivot . . . 44
XIV Flop count of various procedure . . . 45

3 Matrix Analysis (矩陣分析) . . . 45
I Eigenvector or characteristic vector (特徵向量) and characteristic polynomial (特徵多項式) . . . 45
i Eigenvector or characteristic vector (特徵向量) and eigenvalue (本徵值、固有值、特徵值或特徵根) . . . 45
ii Characteristic polynomial . . . 45
iii Multiplicity of eigenvalues . . . 45
iv Number of eigenvalues . . . 46
v Invertible iff has nonzero eigenvalue . . . 46
vi Cayley–Hamilton theorem (凱萊–哈密頓定理) . . . 46
vii Diagonalization (對角化) . . . 46
viii Definiteness (正定性) . . . 46
ix Eigenvalues Test of Definiteness for Symmetric Hessians of Scalar Fields . . . 47
II Generalized Definition of Inverse . . . 47
i One-sided inverse (單邊逆) . . . 47
ii Generalized inverse or g-inverse (廣義逆) . . . 48
iii Reflexive generalized inverse (自反廣義逆) . . . 48
iv Moore–Penrose inverse (摩爾—彭若斯廣義逆) or pseudoinverse (偽逆或擬反) . . . 48
4 Transformation Matrices . . . 49
I Groups . . . 49
i General linear group . . . 49
ii Orthogonal group or general orthogonal group . . . 49
iii Special orthogonal group or rotation group . . . 49
II Transformation matrices in two-dimensional space . . . 49
i Degenerate matrices . . . 49
ii Stretching (伸縮/縮放/拉伸) . . . 50
iii Shearing (推移) . . . 50
iv Rotation (旋轉) . . . 50
v Reflection (鏡射/反射) . . . 50
vi Orthogona projection (正射影) . . . 50
vii Exmaple 1 . . . 51
viii Exmaple 2 . . . 51

ix Example 3 51
x Example 4 52
III Transformation matrices in three-dimensional space 52
i Stretching (伸縮/縮放/拉伸) 52
ii Shearing (推移) 52
iii Rotation (旋轉) 53
iv Rodrigues' rotation formula 53
v Reflection (鏡射/反射) 54
vi Orthogona projection (正射影) 54

Chapter 1 Linear Algebra (線性代數) and Functional Analysis (泛函分析)

1 Vector Space and Functional

I General

i Vector space (向量空間) or Linear space (線性空間)

A vector space over a field F is a non-empty set V together with a binary operation and a binary function that satisfy the eight axioms listed below. In this context, the elements of V are commonly called vectors, and the elements of F are called scalars.

The binary operation, called vector addition or simply addition assigns to any two vectors v and w in V a third vector in V which is commonly written as $v + w$, and called the sum of these two vectors.

The binary function, called scalar multiplication, assigns to any scalar a in F and any vector v in V another vector in V, which is denoted av.

To have a vector space, the eight following axioms must be satisfied for every u, v and w in V, and a and b in F.

1. Associativity of vector addition: $u + (v + w) = (u + v) + w$.
2. Commutativity of vector addition: $u + v = v + u$.
3. Identity element of vector addition: There exists an element $0 \in V$, called the zero vector, such that $\forall v \in V : v + 0 = v$.
4. Inverse elements of vector addition: $\forall v \in V : \exists$ an element $-v \in V$, called the additive inverse or inverse of v, such that $v + (-v) = 0$.
5. Compatibility of scalar multiplication with field multiplication: $a(bv) = (ab)v$.
6. Identity element of scalar multiplication: $1v = v$, where 1 denotes the multiplicative identity in F.
7. Distributivity of scalar multiplication with respect to vector addition: $a(u + v) = au + av$.
8. Distributivity of scalar multiplication with respect to field addition: $(a + b)v = av + bv$.

When the scalar field is the real numbers, the vector space is called a real vector space, and when the scalar field is the complex numbers, the vector space is called a complex vector space. These two cases are the most common ones, but vector spaces with scalars in an arbitrary field F are also commonly considered. Such a vector space is called an F-vector space or a vector space over F.

When a vector is an n-scalar tuple, we call the scalars components and denote the vector as

$$(x_1, x_2, \dots, x_n)$$

or

$$\langle x_1, x_2, \dots, x_n \rangle$$

ii Linear combination (線性組合)

A linear combination of a series of vectors $(v_1, \dots, v_n)$ in a vector space V over a field F is any vector of the form

$$\sum_{i=1}^{n} c_i v_i$$

where $(c_1, \dots, c_n)$ is a series of scalars in F called coefficients.

iii Isomorphism (同構)

Vector spaces X and Y are called isomorphic if there exists a mapping $f : X \to Y$ between them such that f is bijective and continuous and f^{-1} is continuous, written as $X \cong Y$, and f is called a isomorphism between them.

iv Division

For nonzero vector a and vector b in a vector space such that for some scalar s, $sa = b$, $\frac{b}{a} \coloneqq s$.

v Affine space (仿射空間)

An affine space is a point set A together with a vector space $\vec{A}$, and a transitive and free action of the additive group of $\vec{A}$ on the set A. The vector space $\vec{A}$ is said to be associated to the affine space, and its elements are called vectors, translations, or sometimes free vectors.

Explicitly, the definition above means that the action is a mapping, generally denoted as an addition,

$$A \times \vec{A} \to A$$
$$(a, v) \mapsto a + v$$

that has the following properties.

1. Right identity:
$$\forall a \in A : \ a + 0 = a, \quad \text{where } 0 \text{ is the zero vector in } \vec{A}$$
2. Associativity:
$$\forall v, w \in \vec{A}, \forall a \in A, \ (a + v) + w = a + (v + w)$$

3. Free and transitive action:

$$\forall a \in A : \text{ the mapping } \vec{A} \to A : \ v \mapsto a + v \text{ is a bijection.}$$

4. Existence of one-to-one translations

$$\forall v \in \vec{A} : \text{ the mapping } \vec{A} \to A : \ v \mapsto a + v \text{ is a bijection.}$$

vi Linear map (線性映射) or linear operator

For vector spaces V, W, a map $T : \ V \to W$ between vector spaces V and W over field F is linear if

$$\forall a, b \in F, v, w \in V : \ T(av + bw) = aT(v) + bT(w),$$

also called linear operator.

The space of linear operators from V to W is a vector space, denoted as $L(V, W)$.

vii Linear transformation (線性變換)

A linear transformation is a bijection from an vector space onto itself that is an linear map.

viii Invertible map (可逆映射)

A map $T : \ V \to W$ is Invertible if and only if it is bijective. The unique inverse of T, called T^{-1}, is defined to be:

$$T^{-1} : \ W \to V; \forall v \in V : \ T(v) \mapsto v = T^{-1}(T(v)).$$

ix Linear subspace, vector subspace, or subspace

A linear subspace of a vector space is a subset of it that is closed under vector addition and scalar multiplication, and thus a vector space.

x Linear span, linear hull, or span

A span of a subset S, denoted as $\operatorname{span} S$ or $\langle S \rangle$, of a vector space is the smallest linear subspace containing S, and it is called that S is a generating/spanning set of $\operatorname{span} S$ and S generates/spans $\operatorname{span} S$.

xi In the span iff new span unchanged

Let S be a set of vectors and v be a vector. $v \in \operatorname{span} S$ iff $\operatorname{span} S = \operatorname{span}(S \cup \{v\})$.

Proof. Only if: Suppose $v \in \operatorname{span} S$. We can write v as a linear combination of vectors in S. For any $w \in \operatorname{span}(S \cup \{v\})$, we can write w as a linear combination of vectors in $S \cup \{S\}$. By replacing v with the linear combination of vectors in S, we can write w as a linear combination of vectors in S.

If: Suppose $v \notin \operatorname{span} S$. Since $v = 1v$, $v \in \operatorname{span}(S \cup \{v\}) \setminus S$, thus $\operatorname{span} S \neq \operatorname{span}(S \cup \{v\})$. □

xii Linear independence (線性獨立)

A set of nonzero vectors $\{v_1, \dots, v_n\}$ is linearly independent iff the equation:

$$\sum_{i=1}^{n} c_i v_i = 0$$

is true only if all coefficients $c_1, c_2, \dots, c_n$ are zero, then the set of vectors is linearly independent; otherwise the set of vectors is linearly dependent.

xiii Linearly dependent set condition

A set S of vectors is linearly dependent iff either $\mathbf{0} \in S$ or there exists $v \in S$ such that $v \in \text{span}(S \setminus \{v\})$.

Proof. If $\mathbf{0} \in S$, its coefficient can be any. Otherwise, pick a nonzero c_k, the corresponding vector can be written as a linear combination of others. □

xiv Linear independent set as minimum span generating set by inclusion of span

A subset X of a vector space is linearly dependent iff there exists a proper subset of X whose span is the span of X.

Proof. If: Let $X = \{x_1, \dots, x_n\}$. There exist not all zero c_i such that

$$\sum_{i=1}^{n} c_i x_i = 0$$

Take any k such that $c_k \neq 0$, we can write x_k in terms of other x_i, thus x_k is in the span of $X \setminus \{x_k\}$, thus $X \setminus \{x_k\}$ is a proper subset of X whose span is the span of X.

Only if: Suppose Y is proper subset of X such that the span of Y is the span of X. Choose $x \in X \setminus Y$. By the definition, there exists $y_1, \dots, y_m \in Y$ such that x is a linear combination of them. □

xv Spanning set element exchange theorem

Let S be a set of vectors and v a nonzero vector.

$$v \in \text{span}\, S \implies \exists w \in S : \ w \in \text{span}((S \setminus \{w\} \cup \{v\}))$$

Proof. Write v as a linear combination of elements of S. Choose one of them with nonzero coefficient, w, which must exist since v is nonzero, w can be written as a linear combination of elements in $S \setminus \{w\} \cup \{v\}$. □

Let S be a set of linearly independent vectors and v a nonzero vector.

$$v \in \text{span}\, S \iff \exists w \in S : \ w \in \text{span}((S \setminus \{w\} \cup \{v\}))$$

Proof. Write w as a linear combination of elements of elements in $(S \setminus \{w\} \cup \{v\}$. The v term must have nonzero coefficient since otherwise $w \in \text{span}(S \setminus \{w\})$ and S is not linearly independent. Rearrange the linear combination gives v as a linear combination of elements in S. □

xvi Steinitz exchange lemma

Let U be a set of $m \in \mathbb{N}$ linearly independent vectors and W be a set of $n \in \mathbb{N}$ vectors such that $U \subseteq \operatorname{span} W$. Then, $m \leq n$, and, there exists a subset W' of W such that $\left|W'\right| \leq n - m$ and $\operatorname{span}\left(U \cup W'\right) = \operatorname{span} W$.

Proof. First statement: Write both U and W as column vectors of arbitrary order. By $U \subseteq \operatorname{span} W$, there exists $A \in \mathbb{F}^{m \times n}$ such that $U = AW$. Because U is linearly independent, A must have row rank m. Since $\operatorname{rank}(A) \leq \min(m, n)$, $m \leq n$.

Second statement: Let $W_0 = W$. For each $u_i \in U$, write u_i as a linear combination of elements of W_{i-1}. Choose one of them with nonzero coefficient, w_i, which must exist since U is linearly independent, w_i can be written as a linear combination of elements in $W_i \coloneqq W_{i-1} \setminus \{w_i\} \cup \{u_i\}$. □

xvii Basis (基)

The basis of a vector space X over a field F is a set of linearly independent vectors $\mathbf{v} = \{v_1, v_2, \dots\} \subseteq X$ such that for any vector $\mathbf{P} \in X$, there exists a unique coefficient vector $(c_1, c_2, \dots) \subseteq F$ such that:

$$\sum_{i=1} c_i v_i = \mathbf{P}.$$

xviii Orientation

The orientation of a vector space is the arbitrary choice of which ordered bases are ”positively” oriented and which are ”negatively” oriented.

xix Dimension (維度)

The dimension of a vector space V, denoted as $\dim V$, is defined as the cardinality of a basis of it, which is unique (for infinite dimension, if axiom of choice is assumed).

The dimension of an affine space V, denoted as $\dim V$, is defined as the dimension of its associated vector space.

xx Basis extension theorem

For a vector space V of dimension $n \in \mathbb{N}$, for any linearly independent set of finitely many vectors $\{v_1, \dots, v_m\}$ in V, if $m = n$, it is a basis; otherwise, there exists a set of $n - m$ vectors $\{v_{m+1}, \dots, v_n\}$ in V such that $\{v_1, \dots, v_n\}$ is linearly independent, and any such $v_1, \dots, v_n$ is a basis of V.

Proof. For $m = n$, $\operatorname{span}\{v_1, \dots, v_n\} = V$, thus $\{v_1, \dots, v_n\}$ is a basis. For $m < n$, since $S \coloneqq \operatorname{span}\{v_1, \dots, v_m\}$ is of dimension $m < n$, choose any $v_{m+1} \in V \setminus S$. Assume for contradiction that $\{v_1, \dots, v_{m+1}\}$ is linearly dependent, we can write v_{m+1} as a linear combination of $v_1, \dots, v_m$, which means $v_{m+1} \in S$. Repeat until n. □

xxi Standard (unit/basis) vector

For a vector space, the standard (unit/basis) vectors are the unit vectors with one component being the multiplicative identity 1_F and others being the additive identity 0_F.

xxii Bilinear form (雙線性形式)

Let V be a vector space of dimension n over a field $\mathbb{K}$. A map $B : V \times V \to \mathbb{K}$ is a bilinear form on the space if:

1. $\forall u, v, w \in V : B(u + v, w) = B(u, w) + B(v, w)$.
2. $\forall u, v \in V, \lambda \in \mathbb{K} : B(\lambda u, v) = \lambda B(u, v)$.
3. $\forall u, v, w \in V : B(u, v + w) = B(u, v) + B(u, w)$.
4. $\forall u, v \in V, \lambda \in \mathbb{K} : B(u, \lambda v) = \lambda B(u, v)$.

xxiii Symmetric bilinear form (對稱雙線性形式)

Let V be a vector space of dimension n over a field $\mathbb{K}$. A map $B : V \times V \to \mathbb{K}$ is a symmetric bilinear form on the space if:

1. $\forall u, v \in V : B(u, v) = B(v, u)$.
2. $\forall u, v, w \in V : B(u + v, w) = B(u, w) + B(v, w)$.
3. $\forall u, v \in V, \lambda \in \mathbb{K} : B(\lambda u, v) = \lambda B(u, v)$.

xxiv Topological vector space (TVS) (拓樸向量空間)

A topological vector space X is a vector space over a topological field $\mathbb{K}$, such that vector addition $X \times X \to X$ and scalar multiplication $\mathbb{K} \times X \to X$ are continuous functions.

xxv Functional (泛函)

A functional is a function from a vector space into the field of real or complex numbers. For a vector space X, $x \in X$ and a functional on f on X, $f(x)$ is also denoted as $\langle f, x \rangle$ or fx.

xxvi Seminorm (半範數)

Given a vector space X over a ordered field F, a seminorm on X is a real-valued function $p : X \to \mathbb{R}$ with the following conditions, where $|s|$ denotes the usual absolute value of a scalar s:

1. Subadditivity/Triangle inequality:

$$\forall x, y \in X : p(x + y) \leq p(x) + p(y).$$

2. Absolute homogeneity:

$$\forall s \in \mathbb{R}, x \in X : p(sx) = |s|p(x).$$

These conditions implies that:

1. Non-negativity: $\forall x \in X :\ p(x) \geq 0$.
2. $p(0) = 0$.

xxvii Norm (範數/模長)

A norm on X is a seminorm $p :\ X \to \mathbb{R}$ with the following properties:

- Positive definiteness/Positiveness/Point-separating:

$$\forall x \in X :\ p(x) = 0 \implies x = 0$$

A vector x such that $p(x) = 1$ is called an unit vector.

xxviii Sublinear function (亞線性函數) or quasi-seminorm (準半範數)

Let $p :\ X \to \mathbb{R}$ be a function on a vector space X over the field $\mathbb{K}$, which is either $\mathbb{R}$ or $\mathbb{C}$. p is called a sublinear function or quasi-seminorm if it satisfies the following conditions:

1. Subadditivity/Triangle inequality:

$$\forall x, y \in X :\ p(x + y) \leq p(x) + p(y).$$

2. Nonnegative homogeneity:

$$\forall x \in X, r \geq 0 :\ p(rx) = rp(x).$$

xxix Operator norm or induced norm

For normed vector spaces V, W, the vector space of linear operators $L(V, W)$ is equipped with a norm $\|\cdot\|_L$ called operator norm or induced norm defined as, for $A \in L(V, W)$,

$$\|A\|_L = \sup_{\|v\|_V \leq 1} \|A(v)\|_W.$$

xxx Absolutely continuous function (絕對連續函數)

Function $f :\ I \subseteq \mathbb{R} \to \mathbb{R}$ is absolutely continuous on I if for every positive number ε, there is a positive number δ such that, for every finite sequence of pairwise disjoint sub-intervals (x_k, y_k) of I with $x_k < y_k \in I$ and cardinality N,

$$\sum_{k=1}^{N} (y_k - x_k) < \delta \implies \sum_{k=1}^{N} |f(y_k) - f(x_k)| < \varepsilon.$$

The collection of all absolutely continuous functions on I is denoted $\mathrm{AC}(I)$.

xxxi Bounded linear operator (有界線性運算子)

A bounded linear operator is a linear transformation $L : X \to Y$ between topological vector spaces X and Y that maps bounded subsets of X to bounded subsets of Y. If X and Y are normed vector spaces, then L is bounded if and only if there exists some $M > 0$ such that:

$$\forall x \in X : \|Lx\|_Y \leq M\|x\|_X.$$

The smallest such M is called the operator norm of L and denoted by $\|L\|$. A bounded operator between normed spaces is continuous and vice versa.

xxxii Uniform norm, sup norm, supremum norm, Chebyshev norm, infinity norm, or max norm

Uniform norms are defined on the function space of bounded functions valued in a normed space. The uniform norm of bounded function $f : X \to (Y, \|\|_Y)$ is defined by

$$\|f\|_\infty = \sup_{x \in X} \|f(x)\|_Y.$$

xxxiii Hahn–Banach theorem (哈恩-巴拿赫定理)

Let $p : X \to \mathbb{R}$ be a sublinear functional on a vector space X over the field $\mathbb{K}$, which is either $\mathbb{R}$ or $\mathbb{C}$. If $f : M \to \mathbb{K}$ is a linear functional on a vector subspace M such that

$$\forall m \in M : f(m) \leq p(m),$$

then there exists a linear functional $F : X \to \mathbb{K}$ such that

$$\forall m \in M : F(m) = f(m),$$

$$\forall x \in X : F(x) \leq p(x).$$

xxxiv p-norm (p-範數)

The p-norm of a n-dimensional vector $x = ((x_i)_{i=1}^n)$ is defined to be

$$\|x\|_p = \left(\sum_{i=1}^{n} |x_i|^p \right)^{\frac{1}{p}}, \quad p \in \mathbb{N}_\infty,$$

where the ∞-norm is

$$\|x\|_\infty = \max_{i=1}^{n} |x_i|.$$

The 1-norm is also called Manhattan norm (曼哈頓範數); the 2-norm is also called Euclidean norm (歐幾里德範數).

xxxv Locally convex topological vector space (LCTVS) or locally convex space

A locally convex topological vector space (LCTVS) $(X, \mathcal{T})$ is a TVS whose topology is generated by a family of seminorms P on it such that:

$$\bigcap_{p \in P} \{x : p(x) = 0\} = \{0\}$$

That is, $\mathcal{T}$ is generated by a basis $\{U_{\epsilon,x}\}_{\epsilon \in \mathbb{R}_{\geq 0}, x \in X}$ of neighborhoods defined to be:

$$U_{\epsilon,0} = \{y \in X : \forall p \in P : p(y) < \epsilon\}.$$

$$U_{\epsilon,x} = \{y \in X : y - x \in U_{\epsilon,0}\}.$$

This definition implies that a LCTVS is necessarily a Hausdorff space (T2 space).

xxxvi Banach space (巴拿赫空間)

A Banach space is a normed vector space $(X, \|\cdot\|)$ with induced metric d

$$d(x, y) = \|x - y\|, \quad \forall x, y \in X$$

that is a Complete metric space (aka Cauchy space).

xxxvii Dot product (點積)

The dot product of two vectors $\mathbf{a} = (a_1, a_2, \ldots, a_n)$ and $\mathbf{b} = (b_1, b_2, \ldots, b_n)$ is defined as:

$$\mathbf{a} \cdot \mathbf{b} = \sum_{i=1}^{n} a_i b_i.$$

Two nonzero vector are called orthogonal or perpendicular iff their dot product is zero.

xxxviii Inner product space (內積空間)

A bar over a scalar denotes the complex conjugate of it. An inner product space is a vector space V over the real or complex field F together with an inner product, that is, a map

$$\langle \cdot, \cdot \rangle : V \times V \to F$$

that satisfies the following three properties for all vectors $x, y, z \in V$ and all scalars $a, b \in F$.

- Conjugate symmetry:

  $$\langle x, y \rangle = \overline{\langle y, x \rangle}.$$

  This implies that $\langle x, x \rangle$ is a real number.

- Linearity in the first argument:

  $$\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle.$$

- Positive-definiteness: if x is not zero, then $\langle x, x \rangle > 0$.

Every inner product space has an induced norm called canonical norm and is often assumed to be equipped defined by

$$\|x\| = \sqrt{\langle x, x \rangle}$$

The inner product space over $\mathbb{C}$ is also called unitary space.

xxxix Inner product is bilinear form

An inner product is a bililinear form. TODO

Proof. TODO □

xl Orthonormal basis (正交基)

An orthonormal basis for an inner product space V with finite dimension is a basis for V such that for e, f in it:

$$\langle e, f \rangle = \begin{cases} 1, & \text{if } e = f, \\ 0, & \text{if } e \neq f. \end{cases}$$

xli Balanced set (平衡集), circled set, or disk set

Let X be a vector space over a field $\mathbb{K}$ with an absolute value function $|\cdot|$. A subset S of X is called a balanced set or balanced if:

$$\forall a \in \mathbb{K} \text{ s.t. } |a| \leq 1 : \ aS \subseteq S.$$

xlii Well-defined (定義良好)/Unambiguous (不模糊的)

A well-defined expression or unambiguous expression is an expression whose definition assigns it a unique interpretation or value. Otherwise, the expression is said to be not well defined, ill defined or ambiguous.

xliii Affine map (仿射映射)

Given two affine spaces A and B whose associated vector spaces are $\mathscr{A}$ and $\mathscr{B}$, an affine map from A to B is a map $f : A \to B$ such that

$$\begin{aligned} f &: \mathscr{A} \to \mathscr{B} \\ b - a &\mapsto f(b) - f(a) \end{aligned}$$

is a well-defined linear map.

xliv Affine transformation (仿射變換)

An affine transformation is a bijection from an affine space onto itself that is an affine map.

xlv Von Neumann bounded or bounded

Suppose X is a topological vector space over a field $\mathbb{K}$. A subset B of X is called von Neumann bounded (or bounded) in X if for every neighborhood V of the origin, there exists a real $r > 0$ such that $B \subseteq sV$ for all scalars s satisfying $|s| \geq r$.

xlvi Convex set (凸集)

Let S be a vector space or an affine space over some ordered field. A subset C of S is convex if, for all x and y in C, the line segment connecting x and y is included in C.

xlvii (Algebraic) dual (對偶) (vector) (space)

Given any vector space V over a field F, the dual space V^* is defined as the set of all linear maps $\varphi : V \to F$. The dual space V^* itself thus becomes a vector space over F when equipped with an addition and scalar multiplication satisfying that for any $\varphi, \psi \in V^*$, for any $x \in V$, for any $a \in F$,

$$(\varphi + \psi)(x) = \varphi(x) + \psi(x),$$

$$(a\varphi)(x) = a\,(\varphi(x))\,.$$

Element in dual is called covector.

xlviii Continuous dual (space)

Given any vector space V over a field F, the continuous dual space V' is defined as the set of all continuous linear maps $\varphi : V \to F$. The continuous dual space V' itself thus becomes a vector space over F when equipped with an addition and scalar multiplication satisfying that for any $\varphi, \psi \in V'$, for any $x \in V$, for any $a \in F$,

$$(\varphi + \psi)(x) = \varphi(x) + \psi(x),$$

$$(a\varphi)(x) = a\,(\varphi(x))\,.$$

xlix Transpose, dual map, or adjoint

For two vector spaces X, Y and a continuous linear map $A : X \to Y$, the transpose, dual map, or adjoint of A is defined as the unique map ${}^tA : Y' \to X'$ such that for all $x \in X$, for all $y \in Y'$,

$$\langle y, A(x)\rangle = \langle {}^tA(y), x\rangle.$$

l Strong dual (space)

A strong dual space is a continuous dual space of a TVS equipped with the strong (dual) topology, also called topology of uniform convergence on bounded subsets. Suppose X is a TVS over field $\mathbb{F}$ and $\mathcal{B}$ is a family of bounded subsets of X such that every bounded subset of X is a subset of some $B \in \mathcal{B}$. Define the family T of seminorms $\|\cdot\|_B$ on X' for all $x' \in X'$ as

$$T = \{\|x'\|_B \coloneqq \sup_{x \in B} \left|x'(x)\right| \mid B \in \mathcal{B}\}.$$

The strong dual topology is defined as the topology generated by T, and thus B' is a LCTVS.

li Orthogonal complement (正交補)

An orthogonal complement of a subspace W of a vector space V equipped with a bilinear form B is the set $W^\perp$ of all vectors in V that are orthogonal to every vector in W, of which $v \in V$ is orthogonal to $w \in V$ if and only if $B(v, w) = 0$.

lii Affine combination

An affine combination is a linear combination of points where all coefficients sum up to 1.

liii Convex combination

A convex combination is a linear combination of points where all coefficients are non-negative and sum up to 1.

liv Coordinate system (座標/坐標系統)

A coordinate system is a system that uses one or more numbers, or coordinates, to uniquely determine the points on a manifold.

lv Projection (投影)

A (vector) projection onto a subset U of a vector space V is a linear operator $P : V \to U$ such that $P^2 = P$, denoted as $\text{proj}_U P$. The property $P^2 = P$ is called idempotency (冪等性). Its norm is called scalar projection denoted as $\text{comp}_U P$.

lvi Orthographic projection (正射影)

An orthographic on a subset U of a vector space V is a projection $P : V \to U$ such that

$$\forall v \in V : v - P(v) \in U^*$$

where U^* is the orthogonal complement of U.

lvii Hilbert space (希爾伯特空間)

A Hilbert space is a real or complex inner product space that is also a complete metric space with respect to the distance function induced by the inner product.

lviii Angle (夾角) between two vectors

Given two vectors u, v in an inner product space, the angle θ between than is defined as

$$\theta = \arccos(u \cdot v).$$

lix Principal angles (主夾角) between two subspaces

Given two subspaces U, W of an inner product space with $\dim(U) = k \leq \dim(W) = m$, there exists then a sequence of k angles $0 \leq \theta_1 \leq \theta_2 \leq \ldots \leq \theta_k \leq \frac{\pi}{2}$ called the principal angles between U and W, the first one defined as

$$\theta_1 := \min \left\{ \arccos \left(\frac{|\langle u, w \rangle|}{\|u\| \|w\|} \right) \middle| u \in U \wedge w \in W \right\} = \angle(u_1 \in U, w_1 \in W),$$

The vectors u_1 and w_1 are the corresponding principal vectors.

The other principal angles and vectors are then defined recursively via

$$\theta_i := \min \left\{ \arccos \left(\frac{|\langle u, w \rangle|}{\|u\| \|w\|} \right) \middle| u \in U \wedge w \in W \wedge \forall j \in \mathbb{N} \wedge j < i : \langle u_i, u_j \rangle = \langle w_i, w_j \rangle = 0 \right\} = \angle(u_i \in U, w_i \in W$$

lx Flat, hyperplane, or affine subspace (仿射子空間)

A flat or an affine subspace of an affine space is a subspace of it that is an affine space.

lxi Hypersurface

A hypersurface is a $(n-1)$-dimensional manifold that is a subspace of an n-dimensional manifold.

lxii Parallel (平行)

Two nonzero vectors u, v in a vector space over field F are called to be parallel if there exists a scalar $x \in F$ such that $xu = v$, denoted as $u \parallel v$.

Two affine subspaces are called to be parallel if the direction spaces of them D, E satisfies:

$$D \subseteq E \vee E \subseteq D,$$

denoted as $D \parallel E$.

lxiii Coplanar

Multiple elements of a vector or affine space are called coplanar if they are in the same 2D affine subspace.

lxiv Elementwise multiplication

For two vectors consists of scalar components $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$, their elementwise multiplication, denoted as $\mathbf{u} \odot \mathbf{v}$, is defined as

$$\mathbf{u} \odot \mathbf{v} = (u_1 v_1, \dots, u_n v_n).$$

lxv Unit vector (單位向量) or normalized vector

A unit vector or a normalized vector in a normed vector space is a vector of norm 1, denoted by a lowercase letter with a circumflex or hat, as in $\hat{\mathbf{v}}$ (pronounced "v-hat"). The unit vector or normalized vector $\hat{\mathbf{u}}$ of a nonzero vector $\mathbf{u}$ in a normed vector space equipped with a norm $\|\cdot\|$ is defined as:

$$\hat{\mathbf{u}} = \frac{\mathbf{u}}{\|\mathbf{u}\|}.$$

lxvi Affine independence

For an affine space U associated with vector space V, $u_0, u_1, \dots, u_n \in U$ are affinely independent iff $(u_1 - u_0), (u_2 - u_0), \dots, (u_n - u_0)$ are linearly independent in V.

lxvii Simplex

A k-simplex or k-dimensional simplex in an affine space U is a set

$$\{\sum_{i=0}^{k} \theta_i u_i \mid \sum_{i=0}^{k} \theta_i = 1 \wedge \theta_0, \theta_1, \dots, \theta_k \in \mathbb{R}_{\geq 0}\},$$

where

$$u_0, u_1, \dots, u_k \in U$$

are affine independent and called vertices of the simplex.

lxviii Gram matrix, Gramian matrix, or Gramian

For n vectors $v_1, v_2, \dots, v_n$ in an inner product space, the Gram matrix, Gramian matrix, or Gramian G is defined as an $n \times n$ matrix where

$$G_{ij} = \langle v_i, v_j \rangle.$$

The Gram determinant or Gramian is the determinant of the Gram matrix.

lxix Support of functional

For a function $f : U \subseteq \mathbb{R}^n \to \mathbb{R}$ (or $\mathbb{C}$), the support of it, denoted as $\operatorname{supp}(f)$, is defined as

$$\operatorname{supp}(f) = \complement(\{x \in U \mid f(x) \neq 0\}).$$

If $\operatorname{supp}(f)$ is a compact set, f is called with compact support. If $\operatorname{supp}(f)$ is a compact subset of a set $V \subseteq U$, f is called with compact support on V.

lxx Euclidean vector space

A Euclidean vector space is a finite-dimensional inner product space over the real numbers, denoted as $\mathbb{R}^n$ for dimension n. The norm-induced metric $d(v, w) = \|v - w\|$ is called the Euclidean distance

lxxi Euclidean affine space (歐幾里德仿射空間) or Euclidean space (歐幾里得空間)

A Euclidean space, also known as Euclidean affine space, is an affine space over $\mathbb{R}$ such that the associated vector space is a Euclidean vector space, and is equipped with the same norm and metric as the associated Euclidean vector space.

lxxii Affine hull (仿射包)

The affine hull $\operatorname{aff}(S)$ of a set of points S in a Euclidean space is the smallest affine space that contains all the points in S, namely, the set of all affine combinations of the points in S, i.e.:

$$\operatorname{aff}(S) = \left\{ \sum_{i=1}^{k} \alpha_i x_i : k > 0, x_i \in S, \alpha_i \in \mathbb{R}, \sum_{i=1}^{k} \alpha_i = 1 \right\}$$

lxxiii Convex hull (凸包)

The convex hull $\operatorname{conv}(S)$ of a set of points S in a Euclidean space is the smallest convex set that contains all the points in S, namely, the set of all convex combinations of points in S, i.e.:

$$\operatorname{conv}(S) = \left\{ \sum_{i=1}^{k} \alpha_i x_i : k > 0, x_i \in S, \alpha_i \in [0, 1], \sum_{i=1}^{k} \alpha_i = 1 \right\}$$

lxxiv Cartesian coordinate system (笛卡爾座標系統) or rectangular coordinate system (直角座標系統)

The Cartesian or rectangular coordinate system is a coordinate system in a Euclidean vector space E determined by an orthogonal basis of unit vectors such that the inner product of two vectors in E equals to the dot product of them, and that the norm of E is 2-norm.

lxxv Lattice points (格子點)

In a Cartesian coordinate system, vectors whose components are all integers are called lattice points.

lxxvi Curvilinear coordinate system

A curvilinear coordinate system is a coordinate system for Euclidean space such that there exists an invertible map from a Cartesian coordinate system to it.

lxxvii Standard, unit, or probability simplex

The standard, unit, or probability k-simplex is the simplex in $\mathbb{R}^{k+1}$ whose vertices are the unit vectors of each axis, that is, the set

$$\{(p_0, p_1, \dots, p_k) \mid \sum_{i=0}^{k} p_i = 1 \wedge p_0, p_1, \dots, p_k \in \mathbb{R}_{\geq 0}\}.$$

lxxviii Rectangle

An open rectangle in $\mathbb{R}^n$ is the product of n open intervals.

An closed rectangle in $\mathbb{R}^n$ is the product of n closed intervals.

Rectangle may refer to either, or product of n intervals, or product of n intervals satisfying certain constraints, depending on context, and often the case they make no non-trivial difference if not specified.

II Tensor

i Tensor

Let V be a vector space and V^* be its dual. A tensor of type (r, s) is a multilinear map

$$T : (V^*)^r \times V^s \to \mathbb{R},$$

where r is the number of contravariant slots, which take covectors in V^*, and s is the number of covariant slots, which take vectors in V. The order or rank of it is $r + s$.

This map

$$T : (V^*)^r \times V^s \to \mathbb{R}$$

is also a map

$$T : (V^*)^q \times V^t \to \{S : (S : (V^*)^{r-q} \times V^{s-t} \to \mathbb{R})\}$$

for any $q \in \mathbb{Z} \cap [0, r]$ and $t \in \mathbb{Z} \cap [0, s]$.

A scalar is a tensor of type $(0, 0)$. A vector is a tensor of type $(1, 0)$, which maps a covector to a real number, thus also called a contravariant tensor. A covector is a tensor of type $(0, 1)$, which maps a vector to a real number, thus also called a covariant tensor.. A matrix as a linear map $V \to V$ is a tensor of type $(1, 1)$. A matrix as a bilinear form $V \times V \to \mathbb{R}$ is a tensor of type $(0, 2)$. A matrix as $V^* \times V^* \to \mathbb{R}$ is a tensor of type $\mathbb{R}$.

It's conventional that a vector is a column vector and a covector is a row vector. Tensor convention follows.

A tensor can be represented as an array whose dimension is its rank when a basis is chosen.

ii Upper and lower indices

Dummy indices such as μ and ν are used to indicate contravariant and covariant vectors and tensors.

We use a superscript of r dummy indices and a subscript of s dummy indices to indicate a tensor of type (r, s), i.e., $(V^*)^r \times V^s \to \mathbb{R}$ for vector space V, such as $T^{\mu_1 \cdots \mu_r}_{\nu_1 \cdots \nu_s}$. Each ith component of each upper index μ is denoted by replacing μ with i. Each ith component of each lower index μ is denoted by replacing μ with i. For vectors, that is, $A^\mu = (A^0, A^1, A^2, A^3)$. For covectors, that is, $A_\mu = (A_0, A_1, A_2, A_3)$.

Each upper index μ is fixed to a summation $\sum_{\mu=0}^{\dim(V)}$ when it takes an input of the same upper index μ, i.e.

$$T^{\mu \cdots 1}_{\cdots 2} U^{\mu \cdots 3}_{\cdots 4} = \sum_{\mu=0}^{\dim(V)} T^{\mu \cdots 1}_{\cdots 2} U^{\mu \cdots 3}_{\cdots 4}.$$

Each lower index μ is fixed to a summation $\sum_{\mu=0}^{\dim(V)}$ when it takes an input of the same lower index μ, i.e.

$$T^{\cdots 1}_{\mu \cdots 2} U^{\cdots 3}_{\mu \cdots 4} = \sum_{\mu=0}^{\dim(V)} T^{\cdots 1}_{\mu \cdots 2} U^{\cdots 3}_{\mu \cdots 4}.$$

The indexing replaces the order of them in a formula and makes the multiplication notation of tensors work without extra notations.

III Two dimensional Euclidean vector space

i Quadrant (象限)

I(+, +), II(-, +), III(-, -), IV(+, -).

IV Three dimensional Euclidean vector space

i Right-hand rule (右手定則)

Right-hand rule is a common used orientation for a three-dimensional space. In a right-handed Cartesian coordinate system, if you point the thumb of your right hand in the positive x-axis direction and your index finger in the positive y-axis direction, then your middle finger (extended perpendicularly from the palm) will point in the positive z-axis direction. If not specified otherwise, right-handed Cartesian coordinate system is usually used.

ii Octant (卦限)

I(+, +, +), II(-, +, +), III(-, -, +), IV(+, -, +), V(+, +, -), VI(-, +, -), VII(-, -, -), VIII(+, -, -).

iii Cross product (叉積), external product (外積), or vector product (向量積)

The cross product of vector $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$ is defined as:

$$\mathbf{a} \times \mathbf{b} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{pmatrix},$$

where $\mathbf{i} = (1, 0, 0)$, $\mathbf{j} = (0, 1, 0)$, $\mathbf{k} = (0, 0, 1)$.

iv Basic properties of cross product

- Self cross product is the zero vector:

$$\mathbf{a} \times \mathbf{a} = \mathbf{0}.$$

- Anticommutative law (反交換律):

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}.$$

- Left and right distributive over addition and subtraction:

$$\mathbf{a} \times (\mathbf{b} \pm \mathbf{c}) = \mathbf{a} \times \mathbf{b} \pm \mathbf{a} \times \mathbf{c}.$$

$$(\mathbf{b} \pm \mathbf{c}) \times \mathbf{a} = \mathbf{b} \times \mathbf{a} \pm \mathbf{c} \times \mathbf{a}.$$

- Commutative and associative over scalar multiplication:

$$(r\mathbf{a}) \times \mathbf{b} = a \times (r\mathbf{b}) = r(\mathbf{a} \times \mathbf{b}).$$

- Two vector in $\mathbb{R}^3$ are parallel iff their cross product is the zero vector.

v Triple product identity (三重積恆等式)/Jacobi identity (雅可比恆等式)

$$\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) + \mathbf{B} \times (\mathbf{C} \times \mathbf{A}) + \mathbf{C} \times (\mathbf{A} \times \mathbf{B}) = 0.$$

$$\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \mathbf{B} \cdot (\mathbf{C} \times \mathbf{A}) = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B}) = \begin{vmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{vmatrix}$$

$$(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = (\mathbf{A} \cdot \mathbf{C})\mathbf{B} - (\mathbf{B} \cdot \mathbf{C})\mathbf{A}.$$

vi Scalar triple product

$$[\mathbf{a},\mathbf{b},\mathbf{c}] = \det\big(\mathbf{a},\mathbf{b},\mathbf{c}\big) = \mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})$$

2 Matrix (矩陣)

I Matrix

A matrix (plural: matrices) is a rectangular array of numbers (or other mathematical objects), called the entries (元或元素) of the matrix. Most commonly, a matrix over a field F is a rectangular array of elements of F.

The size of a matrix is defined by the number of rows (zh-hant: 列/zh-hans: 行) and columns (zh-hant: 行/zh-hans: 列) it contains. A matrix with m rows and n columns is called an $m \times n$ matrix, or m-by-n matrix, where m and n are called its dimensions (階). A matrix with the same number of rows and columns n is called a n square matrix (方陣). A matrix with an infinite number of rows or/and columns is called an infinite matrix. A matrix with exactly one row is called a row vector. A matrix with exactly one column is called a column vector.

The entry in the i-th row and j-th column, called the i,jth entry (第 i,j 元) or ijth entry (第 ij 元), is commonly denoted as $a_{i,j}$ or a_{ij}.

An $m \times n$ matrix $\mathbf{A}$ is commonly denoted as:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix},$$

$$\mathbf{A} = [a_{ij}]_{1\le i\le m, 1\le j\le n},$$

$$\mathbf{A} = (a_{ij})_{1\le i\le m, 1\le j\le n},$$

$$\mathbf{A} = [a_{ij}]_{m\times n},$$

or

$$\mathbf{A} = (a_{ij})_{m\times n}.$$

The i,ith entries are called diagonal entries (對角線元), and other entries are called off-diagonal or non-diagonal entry. The diagonal line of diagonal entries is called main diagonal.

A matrix with only one row is called a row matrix or row vector, and a matrix with only one column is called a column matrix or column vector.

The ith row of the matrix $\mathbf{A}$ is commonly denoted as $\mathbf{A}_{i,:}$ or $\mathbf{A}_i$, and the jth column is commonly denoted as $\mathbf{A}_{:,j}$.

The set of all $m \times n$ matrices over $\mathbb{F}$ is denoted as $\mathbb{F}^{m \times n}$.

II Basic operations

i Submatrix

A submatrix of a matrix M is obtained by deleting from M entire rows, entire columns, or both.

ii Matrix equality

We say that two matrices $\mathbf{A} = [a_{ij}]_{m \times n}$ and $\mathbf{B} = [b_{ij}]_{m \times n}$ are equal if

$$\forall 1 \leq i \leq m, 1 \leq j \leq n : \ a_{ij} = b_{ij}.$$

iii Matrix addition

The sum of two $m \times n$ matrices **A** and **B** is calculated entrywise.

iv Scalar multiplication

The product $c\mathbf{A}$ of a scalar c is calculated by multiplying every entry of **A** by c.

v Matrix multiplication (矩陣乘法)

Multiplication of two matrices is defined only if the number of columns of the left matrix is the same as the number of rows of the right matrix. The matrix product **AB** of an $m \times n$ matrix **A** and $n \times p$ matrix **B** is the $m \times p$ matrix whose ijth entry is the dot product of the ith row of **A** and the jth column of **B**.

vi Outer product

For two column vectors u, v, the outer product uv^{T}, sometimes denoted as $u \otimes v$, is the matrix product uv^{T}.

vii Positive integer exponent

Given a square matrix **A** and a positive integer n, $\mathbf{A}^n$ is defined as $\prod_{i=1}^{n} \mathbf{A}$.

viii Zero matrix (零矩陣)

A zero matrix over a field F is a matrix of which all entries are the additive identity 0_F of F. Zero matrix is often denoted as **O** or O. $n \times n$ zero matrix is often denoted as $\mathbf{O}_n$ or O_n.

ix Identity matrix (單位矩陣)

An identity matrix over a field F is a square matrix of which all diagonal entries are the multiplicative identity 1_F of F and all off-diagonal entries are the multiplicative absorbing element 0_F of F. Identity matrix is often denoted as **I** or I. $n \times n$ identity matrix is often denoted as $\mathbf{I}_n$ or I_n.

x Hadamard product (哈達瑪乘積), Schur product (舒爾乘積), elementwise product, or entrywise product (逐項乘積)

For two $m \times n$ matrices **A** and **B**, the Hadamard product $\mathbf{A} \odot \mathbf{B}$ or $\mathbf{A} \circ \mathbf{B}$ is an $m \times n$ matrix defined as:

$$(\mathbf{A} \odot \mathbf{B})_{ij} = (\mathbf{A})_{ij}(\mathbf{B})_{ij}.$$

xi Kronecker product (克羅內克積)

If **A** is an $m \times n$ matrix and **B** is a $p \times q$ matrix, then the Kronecker product $\mathbf{A} \otimes \mathbf{B}$ is the $pm \times qn$ block matrix:

$$(\mathbf{A} \otimes \mathbf{B})_{i,j} = a_{\left\lceil \frac{i}{p} \right\rceil, \left\lceil \frac{j}{q} \right\rceil} b_{((i-1) \mod p)+1,((j-1) \mod q)+1}$$

xii Merging matrices

The matrix formed by merging matrices with the same number of rows left and right is denoted by writing them in columns of a row vector.

The matrix formed by merging matrices with the same number of columns up and down is denoted by writing them in rows of a column vector.

xiii Property

For any scalars r and s, $m \times n$ matrices A, B, C, D, zero matrix O, and identity matrix I, the following statements are true,

$$A + B = B + A$$

$$A + (B + C) = (A + B) + C$$

$$r(A + B) = rA + rB$$

$$rA + sA = (r + s)A$$

$$(rs)A = r(sA) = s(rA)$$

$$(AB)C = A(BC)$$

$$r(AB) = (rA)B = A(rB)$$

$$A(B + C) = AB + AC$$

$$(A + B)C = AC + BC$$

$$(A + B)(C + D) = AC + AD + BC + BD$$

$$A \odot B = B \odot A$$

$$A \odot (B \odot C) = (A \odot B) \odot C$$

$$A \odot (B + C) = A \odot B + A \odot C$$

$$IA = A$$

$$AI = A$$

$$A + O = A$$

$$AO = O$$

$$OA = O$$

the following statements are not necessarily true,

$$AB = BA$$

Proof. Counterexample:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$AB = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$BA = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

□

$$(AB = AC \wedge A \neq O) \Rightarrow B = C$$

Proof. Counterexample:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$C = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$$

$$AB = AC = A$$

□

$$(ACB = ADB \wedge A \neq O \wedge B \neq O) \Rightarrow C = D$$

Proof. Counterexample:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$B = C = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$D = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$$

$$ACB = AC = ADB = AD = A$$

□

$$A^2 = O \Rightarrow A = O$$

Proof. Counterexample:

$$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

□

$$AB = O \Rightarrow (A = O \vee B = O)$$

Proof. Counterexample:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

□

$$AB = B \Rightarrow A = I$$

Proof. Counterexample:

$$A = B = AB = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

□

xiv Zero exponent

Given an $n \times n$ matrix **A** and $n \times n$ identity matrix **I**, $\mathbf{A}^0$ is defined as **I**.

xv Trace (跡)

The trace of a matrix **A** is the sum of all diagonal entries of **A**, denoted as tr(**A**).

III Matrix row and column operations

i Elementary row operations (zh-hant: 列/zh-hans: 行運算) and elementary column operations (zh-hant: 行/zh-hans: 列運算)

Elementary row operations include:

- Row addition (zh-hant: 列/zh-hans: 行相加): Adding a row multiplied by a nonzero constant to another.
- Row multiplication/scaling (zh-hant: 列/zh-hans: 行加倍): Multiplying a row by a nonzero constant.
- Row interchange/swap/switch (zh-hant: 列/zh-hans: 行交換): Interchanging two rows of a matrix.

Elementary column operations are defined the same with row replaced with column.

If one matrix can be transformed to another through a sequence of elementary row operations, the two matrices are called row-equivalent.

If one matrix can be transformed to another through a sequence of elementary column operations, the two matrices are called column-equivalent.

ii Row operation as matrix multiplication and elementary matrix

Every elementary row operation on a $m \times n$ matrix A can be performed by left-multiplying an appropriate $m \times m$ matrix:

- Row addition: Adding row i multiplied by $k \neq 0$ to row j can be performed by left-multiplying I_m with the j, ith entry replaced with k.
- Row multiplication: Multiplying row i by a constant $k \neq 0$ can be performed by left-multiplying I_m with the i, ith entry replaced with k.
- Row interchange: Interchanging row i and row j can be performed by left-multiplying I_m with row i and row j interchanged.

A square matrix is called elementary matrix if it can be obtained from the identity matrix of its size with a single elementary row operations.

Let A be a $m \times n$ matrix and E be an $m \times m$ elementary matrix that can be obtained by performing an elementary row operation on I_m. Then, EA is the matrix obtained by performing the same operation on A.

iii Zero row, nonzero row, and leading entry

A row whose all entries are zero is called a zero row, otherwise it is called a nonzero row. The left-most nonzero entry of a nonzero row is called leading entry.

iv Transpose (轉置)

The transpose of an $m \times n$ matrix $\mathbf{A}$ is the $n \times m$ matrix $\mathbf{A}^\top$:

$$(\mathbf{A}^\top)_{i,j} = a_{j,i}$$

Inverse map as linear map is transpose as matrix.

v Symmetric matrix (對稱矩陣)

$\mathbf{A}$ is called a symmetric matrix iff $\mathbf{A} = \mathbf{A}^\top$.

vi Skew-symmetric matrix/antisymmetric matrix (反對稱矩陣)

$\mathbf{A}$ is called a skew-symmetric matrix or an antisymmetric matrix if $\mathbf{A} = -\mathbf{A}^\top$.

vii Transpose identities

For an index set of matrices $\mathbf{A} : [1, n] \cap \mathbb{N} \to \mathbb{F}^{p \times q};\ i \mapsto \mathbf{A}(i)$ for some fixed p, q,

$$\left(\sum_{i=1}^{n} \mathbf{A}(i)\right)^\top = \sum_{i=1}^{n} (\mathbf{A}(i))^\top$$

Proof.

$$\left(\sum_{i=1}^{n} \mathbf{A}(i)\right)_{jk} = \sum_{i=1}^{n} \mathbf{A}(i)_{jk}$$

$$\left(\left(\sum_{i=1}^{n} \mathbf{A}(i)\right)^\top\right)_{jk} = \sum_{i=1}^{n} \mathbf{A}(i)_{kj}$$

$$= \sum_{i=1}^{n} \left((\mathbf{A}(i))^\top\right)_{jk}$$

$$= \left(\sum_{i=1}^{n} (\mathbf{A}(i))^\top\right)_{jk}$$

□

For two matrices $X \in \mathbb{F}^{p \times q}, Y \in \mathbb{F}^{q \times r}$,

$$(XY)^\top = Y^\top X^\top$$

Proof.

$$(XY)_{jk} = \sum_{i=1}^{q} X_{ji} Y_{ik}$$

$$((XY)^\top)_{jk} = \sum_{i=1}^{q} X_{ki} Y_{ij}$$

$$= \sum_{i=1}^{q} (X^\top)_{ik} (Y^\top)_{ji}$$

$$= (Y^\top X^\top)_{jk}$$

□

For an index set of matrices $\mathbf{A}: [1,n] \cap \mathbb{N} \to \mathbb{F}^{\mathbb{N}\times\mathbb{N}};\ i \mapsto \mathbf{A}(i)$ such that $\mathbf{A}(i)\mathbf{A}(i+1)$ is defined for all $i \in [1,n) \cap \mathbb{N}$,

$$\left(\prod_{i=1}^{n} \mathbf{A}(i)\right)^{\mathsf{T}} = \prod_{i=1}^{n} (\mathbf{A}(n-i+1))^{\mathsf{T}}$$

Proof. Prove by mathematical induction on n.

Base case $n = 1$:

$$\left(\prod_{i=1}^{1} \mathbf{A}(i)\right)^{\mathsf{T}} = \prod_{i=1}^{1} (\mathbf{A}(1-i+1))^{\mathsf{T}},$$

which is true.

Assume the statement holds for $n = k$:

$$\left(\prod_{i=1}^{k} \mathbf{A}(i)\right)^{\mathsf{T}} = \prod_{i=1}^{k} (\mathbf{A}(k-i+1))^{\mathsf{T}}$$

For $n = k+1$, we have

$$\begin{aligned}
\left(\prod_{i=1}^{k+1} \mathbf{A}(i)\right)^{\mathsf{T}} &= \left(\prod_{i=1}^{k} \mathbf{A}(i) A(k+1)\right)^{\mathsf{T}} \\
&= (A(k+1))^{\mathsf{T}} \left(\prod_{i=1}^{k} \mathbf{A}(i)\right)^{\mathsf{T}} \\
&= (A(k+1))^{\mathsf{T}} \prod_{i=1}^{k} (\mathbf{A}(k-i+1))^{\mathsf{T}} \\
&= \prod_{i=1}^{k+1} (\mathbf{A}(k-i+1))^{\mathsf{T}}
\end{aligned}$$

□

IV Complex matrix

i (Complex) conjugate

For a complex matrix A, its complex conjugate $\bar{A}$ is defined by taking complex conjugate for all its entries.

ii Conjugate transpose (共軛轉置)/Hermitian conjugate (埃爾米特共軛)/Hermitian transpose (埃爾米特轉置)

For a complex matrix A, its conjugate transpose A^* is defined as

$$A^* = \bar{A}^{\mathsf{T}}$$

iii Conjugate transpose identities

For an index set of matrices $\mathbf{A}: [1,n]\cap\mathbb{N}\to\mathbb{C}^{p\times q};\ i\mapsto\mathbf{A}(i)$ for some fixed p,q,

$$\left(\sum_{i=1}^{n}\mathbf{A}(i)\right)^{*}=\sum_{i=1}^{n}(\mathbf{A}(i))^{*}$$

For two matrices $X\in\mathbb{C}^{p\times q},Y\in\mathbb{C}^{q\times r}$,

$$(XY)^{*}=Y^{*}X^{*}$$

For an index set of matrices $\mathbf{A}: [1,n]\cap\mathbb{N}\to\mathbb{C}^{\mathbb{N}\times\mathbb{N}};\ i\mapsto\mathbf{A}(i)$ such that $\mathbf{A}(i)\mathbf{A}(i+1)$ is defined for all $i\in[1,n)\cap\mathbb{N}$,

$$\left(\prod_{i=1}^{n}\mathbf{A}(i)\right)^{*}=\prod_{i=1}^{n}(\mathbf{A}(n-i+1))^{*}$$

V Cofactor matrix, adjugate matrix, determinant, and inverse

Let $\mathbf{M}_{ij}$ of $\mathbf{A}$ be the determinant of the submatrix of $\mathbf{A}$ without the ith row and jth column.

i Determinant (行列式或判別式)

The determinant of a 1×1 matrix is the only element of it; the determinant of an $n\times n$ matrix $\mathbf{A}$, denoted $\det(\mathbf{A})$ or

$$\begin{vmatrix}a_{11}&a_{12}&\dots&a_{1n}\\a_{21}&a_{22}&\dots&a_{2n}\\\vdots&\vdots&\ddots&\vdots\\a_{n1}&a_{n2}&\dots&a_{nn}\end{vmatrix},$$

is

$$\det(\mathbf{A})=\sum_{i=1}^{n}a_{ij}(-1)^{i+j}\mathbf{M}_{ij}\quad j\le n\in\mathbb{N}$$

$$\det(\mathbf{A})=\sum_{j=1}^{n}a_{ij}(-1)^{i+j}\mathbf{M}_{ij}\quad i\le n\in\mathbb{N}$$

ii Determinant identities

Let $\mathbf{A}$ and $\mathbf{A}_i$ be square matrices.

$$\det\left(\sum_{i=1}^{n}\mathbf{A}_i\right)=\sum_{i=1}^{n}\det\left(\mathbf{A}_i\right)$$

$$\det\left(\prod_{i=1}^{n}\mathbf{A}_i\right)=\prod_{i=1}^{n}\det\left(\mathbf{A}_i\right)$$

$$\det(\mathbf{A})\neq0\implies\det\left(\mathbf{A}^{n}\right)\neq0$$

iii Cofactor matrix (餘因子矩陣)

The cofactor matrix of an $n \times n$ matrix $\mathbf{A}$ is the $n \times n$ matrix $\operatorname{cof}(\mathbf{A})$:

$$[\operatorname{cof}(\mathbf{A})]_{ij} = \left((-1)^{i+j}\mathbf{M}_{ij}\right)$$

iv Adjugate matrix (伴隨矩陣)/classical adjoint (經典伴隨矩陣)

The adjugate matrix or classical adjoint matrix of an $n \times n$ matrix $\mathbf{A}$ is the $n \times n$ matrix $\operatorname{adj}(\mathbf{A})$:

$$\operatorname{adj}(\mathbf{A}) = \operatorname{cof}(\mathbf{A})^{\mathsf{T}}$$

v Inverse (反方陣、逆方陣、反矩陣或逆矩陣), multiplicative inverse (乘法反方陣), or reciprocal

The inverse of an $n \times n$ matrix $\mathbf{A}$ is an $n \times n$ matrix $\mathbf{A}^{-1}$ such that:

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$$

If such $\mathbf{A}^{-1}$ exists, $\mathbf{A}$ is called invertible.

vi Negative integer exponent

Given an invertible matrix $\mathbf{A}$ and a negative integer n, $\mathbf{A}^n$ is defined as $(\mathbf{A}^{-1})^n$.

vii Invertible matrix (可逆矩陣), orthogonal matrix (正交矩陣), orthonormal matrix (正則矩陣), non-singular matrix (非奇異矩陣), or non-degenerate matrix (非退化矩陣)

An invertible matrix, orthogonal matrix, orthonormal matrix, non-singular matrix, or non-degenerate matrix is a matrix such that its inverse exists.

viii Non-invertible matrix (不可逆矩陣), non-orthogonal matrix (非正交矩陣), non-orthonormal matrix (非正則矩陣), singular matrix (奇異矩陣), or degenerate matrix (退化矩陣)

An non-invertible matrix, non-orthogonal matrix, non-orthonormal matrix, singular matrix, or degenerate matrix is a matrix that is not an invertible matrix.

ix Inverse of a two-by-two matrix

For an invertible 2×2 matrix $\mathbf{A}$:

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \implies \mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

x Unitary matrix (么正矩陣或酉矩陣)

An unitary matrix is a square matrix whose inverse equals to its conjugate transpose.

xi Determinant property

- When a row (or column) is added to another, the determinant remains unchanged.
- When all entries of a row (or column) is multiplied by a real number n, the determinant is multiplied by n.
- When two rows (or columns) are swapped, the determinant is multiplied by -1.
- Given matrix $\mathbf{A}$ and its inverse $\mathbf{A}^{-1}$, $\det\left(\mathbf{A}^{-1}\right) = \dfrac{1}{\det(\mathbf{A})}$.

xii Inverse identities

A matrix is invertible iff its transpose is invertible. Product of invertible matrices is invertible.

Let $\mathbf{A}$ and $\mathbf{A}_i$ be invertible matrices, $\mathbf{B}$ be a matrix of the same size of $\mathbf{A}$.

$$\mathbf{A}^{-1} = \frac{\operatorname{adj}(\mathbf{A})}{\det(\mathbf{A})}$$

$$\left(\mathbf{A}^{\top}\right)^{-1} = \left(\mathbf{A}^{-1}\right)^{\top}$$

Proof.

$$\mathbf{A}^{\top}\left(\mathbf{A}^{-1}\right)^{\top} = \left(\mathbf{A}^{-1}\mathbf{A}\right)^{\top} = \mathbf{I}^{\top} = \mathbf{I}$$

□

$$\left(\prod_{i=1}^{n}\mathbf{A}_i\right)^{-1} = \prod_{i=1}^{n}\left(\mathbf{A}_{n-i+1}\right)^{-1}$$

$$(\mathbf{A}^{-1}\mathbf{B}\mathbf{A})^n = \mathbf{A}^{-1}\mathbf{B}^n\mathbf{A}, \quad n \in \mathbb{N}$$

Let A, B be square matrix of same size,

$$AB = I \Rightarrow (\exists A^{-1} \wedge A^{-1} = B)$$

xiii Inverse of elementary matrix

Every elementary matrix is invertible. Moreover, the inverse of an elementary is an elementary matrix.

- Row addition: The inverse of I_m with the j,ith entry replaced with k is I_m with the j,ith entry replaced with $-k$.
- Row multiplication: The inverse of I_m with the i,ith entry replaced with k is I_m with the i,ith entry replaced with $\dfrac{1}{k}$.
- Row interchange: The inverse of I_m with row i and row j interchanged is itself.

Thus, for any two row-equivalent $m \times n$ matrices A, B, there exists an $m \times m$ invertible matrix C, which is the product of a sequence of $m \times m$ elementary matrices, such that $B = CA$.

xiv Vandermonde matrix (范德蒙矩陣)

A Vandermonde matrix **V** is:

$$V_{i,j} = \alpha_i^{\,j-1}$$

An $n \times n$ Vandermonde matrix **V** satisfies:

$$\det(V) = \prod_{1 \le i < j \le n} (\alpha_j - \alpha_i)$$

VI Diagonal matrix

i Diagonal matrix (對角矩陣)

A diagonal matrix is a square matrix whose all off-diagonal entries are the additive identity of 0_F.

ii Diagonal matrix identities

If:

$$\mathbf{D} = \begin{bmatrix} d_1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_n \end{bmatrix}$$

where all $d_i \neq 0$, then:

$$\mathbf{D}^{-1} = \begin{bmatrix} \frac{1}{d_1} & 0 & \cdots & 0 \\ 0 & \frac{1}{d_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{1}{d_n} \end{bmatrix}$$

Let **E**, **F** be two $n \times n$ diagonal matrices:

$$\mathbf{EF} = [e_{ij} f_{ij}]_{1 \le i,j \le n}$$

VII System of Linear Equations (線性方程組)

i System of equations (方程組)

A solution to a system of equations is a choice of variables such that all equations are satisfied. Two systems of equations are called equivalent if they have exactly the same set of solutions.

- **Consistent system (相容方程組)**: A system of equations is consistent iff it has at least one solution.
- **Inconsistent/contradictory system (相容方程組)**: A system of equations is inconsistent or contradictory iff it has no solution.
- **Dependent system (相依方程組)**: A consistent system of equations is dependent iff it has infinitely many solutions.

- **Independent system (獨立方程組)**: A system of equations is independent iff it has finitely many solutions.

‘

ii Linear Equation (線性方程)

A linear equation is an equation that may be put in the from:

$$\sum_{k=1}^{n} a_k x_k = c,$$

where a_k are called coefficients and c is called constant term.

If $c = 0$, the linear equation is called a homogeneous linear equation; otherwise, it is called a heterogeneous or non-homogeneous linear equation.

iii System of Linear Equations (線性方程組) or linear system

A system of linear equations is a collection of two or more linear equations involving the same tuple of variables.

iv Independent system of linear equations has exactly one solution

An independent system of linear equations must have exactly one solution.

Proof. If more than one solution exist, their linear combinations form infinitely many solutions. □

v Dependent system general solution

For a dependent system, we can solve some of the variables, called basic variables, in terms of some other variables, called free variables, such that the free variables are independent from each other and that a choice of free variables uniquely determines basic variables, providing a general solution of the system.

vi Coefficient matrix (係數矩陣)

The coefficients of a system of linear equations written as a matrix, with one row for each equation.

vii Constant term vector

The column vector of the constant terms on the right side of the equal signs of a system of linear equations.

viii Augmented matrix (增廣矩陣)

Coefficient matrix with the constant term vector augmented to the last column of the matrix. For coefficient matrix A and constant term vector b, we write the augmented matrix as $[A|b]$.

ix Row equivalent means same solutions

Two systems of linear equations with same size of augmented matrix have the same solutions if their augmented matrices are row-equivalent.

Two consistent systems of linear equations with same size of augmented matrix have the same solutions iff their augmented matrices are row-equivalent.

Proof. Let $M = [A|b]$ and $N = [C|d]$ both be with m rows and $n+1$ columns.

If: There exists elementary matrices $E_1, \dots, E_k$ such that

$$N = E_k \dots E_1 M$$

We prove that multiplying augmented matrix E by an elementary matrix doesn't change its solution set.

$$EM = [EA|Eb]$$

If $b \notin \text{row}(A)$, since elementary row operation doesn't alter row space, $Eb \notin \text{row}(EA)$. Otherwise, since E is invertible with inverse being elementary matrix, we have

$$Ax = b \implies EAx = EAb$$

$$EAx = Eb \implies E^{-1}EAx = E^{-1}EAb \implies Ax = b$$

Only if: By the existence and uniqueness of reduced row echelon form, if they have the same non-empty solution set, they have the same reduced row echelon form R, and we can construct sequences of elementary matrices $E_1, \dots, E_k$ and $F_1, \dots, F_k$ such that

$$R = E_k \dots E_1 M = F_k \dots F_1 N$$

Since an elementary matrix is invertible with inverse being elementary matrix, we have

$$M = E_1^{-1} \dots E_k^{-1} F_k \dots F_1 N,$$

and that M, N are row-equivalent. □

x Cramer Rule or Cramer formula (克拉瑪公式)

Given a system of linear equations that can be written in the form:

$$\mathbf{AX} = \mathbf{C}$$

where **A** is an $n \times n$ coefficient matrix, **X** is a variable matrix:

$$\mathbf{X} = \left(x_1, x_2, x_3, \dots, x_n\right)^\top,$$

and **C** is a coefficient matrix:

$$\mathbf{C} = \left(c_1, c_2, c_3, \dots, c_n\right)^\top.$$

Let $\mathbf{A}_i$ be $\mathbf{A}$ with the ith column replaced by $\mathbf{C}$. The solution to the system of linear equations is given by:

$$\begin{cases} x_i = \dfrac{\det\left(\mathbf{A}_i\right)}{\det\left(\mathbf{A}\right)}, & \det\left(\mathbf{A}\right) \neq 0 \\ \text{inconsistent}, & \det\left(\mathbf{A}\right) = 0 \wedge \bigvee_{i=1}^{n} \det\left(\mathbf{A}_i\right) \neq 0 \\ \text{dependent}, & \det\left(\mathbf{A}\right) = 0 \wedge \bigvee_{i=1}^{n} \det\left(\mathbf{A}_i\right) = 0 \end{cases}$$

where $x_i = \dfrac{\det\left(\mathbf{A}_i\right)}{\det\left(\mathbf{A}\right)}$ can be geometrically viewed as the volume of the parallelogram formed by the column vectors of $\mathbf{A}_i$ divided by the volume of the parallelogram formed by the column vectors of $\mathbf{A}$.

Particularly, when $\mathbf{C}$ is a zero matrix, that is,

$$\mathbf{AX} = \mathbf{0},$$

the solution to the system of linear equations is given by:

$$\begin{cases} x_i = 0, & \det\left(\mathbf{A}\right) \neq 0 \\ \text{dependent}, & \det\left(\mathbf{A}\right) = 0 \end{cases}$$

VIII Row echelon form

i (General) row echelon form (REF) (（廣義）zh-hant: 列/zh-hans: 行階梯形式)

A matrix is said to be in row echelon form iff:

1. Each nonzero lies above every zero row.
2. The leading entry of every nonzero row lies in a column to the right of the column containing the leading entry of any row above.

Thus, for a leading entry of any row, all entries below it in the same column are zero.

ii Reduced row echelon form (RREF)

A matrix is said to be in reduced row echelon form iff:

1. It is in row echelon form.
2. Each leading entry is the only nonzero entry in the column.
3. The leading entries of all nonzero rows are 1.

iii Pivot position

For a matrix A and its (unique, which will be proven later) reduced row echelon form R, the position containing a leading entry of a nonzero row is called pivot position, a column containing a pivot position is called pivot column, and a row containing a pivot position is called pivot row.

iv Gaussian Elimination (高斯消去法), row reduction, Gauss-Jordan Elimination, and reduced row echelon form test for consistency and obtaining general solution

Gaussian elimination or row reduction refers to either algorithm, or the forward pass only with Gaussian-Jordan elimination referring to either the entire algorithm or the backward pass only.

To find the row echelon form of a matrix, do the followings, called forward pass:

1. Determine the leftmost nonzero column. This is a pivot column, and the topmost position in this column is a pivot position.
2. In the pivot column, choose any nonzero entry, and perform the appropriate row interchange to bring this entry into the pivot position.
3. Add an appropriate multiple of the row containing the pivot position to each lower row in order to change each entry below the pivot position into 0.
4. Ignore the row containing the pivot position. If there is a nonzero row below it, repeat steps 1–4 on the submatrix that remains.

To find the reduced row echelon form of a matrix from a row echelon form of it, from the topmost to bottommost nonzero row, do the followings, called backward pass:

1. Perform the appropriate row scaling to make the leading entry 1.
2. Add an appropriate multiple of the row to each upper row in order to change each entry above the leading entry into 0.

A system of linear equations is consistent iff the reduced row echelon form of its augmented matrix has no row with leading entry in the rightmost column.

Proof. If the reduced row echelon form of its augmented matrix has row with leading entry in the rightmost column, it is obviously inconsistent. If the reduced row echelon form of its augmented matrix has no row with leading entry in the rightmost column, we can construct a general solution. □

Given the reduced row echelon form of the augmented matrix of a linear system, if it has no row with leading entry in the rightmost column, we can obtain the general solution by

1. Assign the variables of non-pivot columns as free variables and variables of pivot columns as basic variables.
2. For each nonzero row, write an equation of the variable of the leading entry as a linear combination of the variables (constant term for the rightmost column and free variables for others) of other nonzero entries.

IX Subspaces of matrices and linear maps

i Four fundamental subspaces of a matrix (矩陣的四大子空間)

The four fundamental subspaces of a matrix are the row space, column space, null space, and left null space.

ii Column space (zh-hant: 行/zh-hans: 列空間)

The column space of $\mathbf{A}$, denoted as $\mathrm{Col}(\mathbf{A})$ or $\mathbf{C}(\mathbf{A})$, is defined as the span of the set of column vectors of $\mathbf{A}$.

iii Row space (zh-hant: 列/zh-hans: 行空間)

The row space of $\mathbf{A}$, denoted as $\mathrm{Row}(\mathbf{A})$, is defined as the span of the set of the row vectors of $\mathbf{A}$, that is, $\mathrm{Col}(\mathbf{A}^\top)$.

iv Null space (零空間), kernel (核), or kernel space (核空間)

The kernel of a linear map $L\colon V \to W$, denoted as $\ker(L)$ or $\mathrm{Null}(L)$, is defined as:

$$\ker(L) = \{v \in V \colon L(v) = 0\}.$$

$\dim\big(\ker(\mathbf{A})\big)$ is called nullity (零化度).

v Left null space (左零空間), left kernel (左核), or left kernel space (左核空間)

The left null space of a linear map $L\colon V \to W$ is defined as $\ker\big(L^{-1}\big)$.

$\dim\big(\ker\big(L^{-1}\big)\big)$ is called left nullity.

vi Image space (像空間) or range (值域)

The image space of a linear map $T\colon V \to W$, denoted as $\mathrm{im}(T)$, is defined as,

$$\mathrm{im}(T) = \{T(v) \colon v \in V\}.$$

vii Cokernel (餘核)

The cokernel of a linear map $T\colon V \to W$, denoted as $\mathrm{coker}(T)$, is defined as,

$$\mathrm{coker}(T) = W \setminus \mathrm{im}(T).$$

X Rank and nullity

i Rank (秩)

The rank of a matrix $\mathbf{A}$, denoted as $\mathrm{rank}(\mathbf{A})$ or $\mathrm{rk}(\mathbf{A})$, is defined as the dimension of its column space (or row space, which are the same by constant rank theorem), that is,

$$\mathrm{rank}(\mathbf{A}) = \dim\big(\mathrm{Col}(\mathbf{A})\big)$$

ii Constant rank theorem (常秩定理)

For any matrix $\mathbf{A}$:

$$\dim\big(\text{Col}(\mathbf{A})\big) = \dim\big(\text{Col}(\mathbf{A}^\top)\big)$$

Proof. Let $\mathbf{A}$ be an $m \times n$ matrix, $r = \dim\big(\text{Col}(\mathbf{A})\big)$, $c_1, c_2, \dots, c_r$ be a basis of $\text{Col}(\mathbf{A})$, matrix $\mathbf{C} = [c_1, c_2, \dots, c_r]$, and $r \times n$ matrix $\mathbf{R}$ such that $\mathbf{A} = \mathbf{CR}$.

Since:

$$\text{Col}(\mathbf{A}^\top) \subseteq \text{Col}(\mathbf{R}^\top),$$

$$\dim\big(\text{Col}(\mathbf{A}^\top)\big) \leq \dim\big(\text{Col}(\mathbf{R}^\top)\big)$$

Because there are only r rows in $\mathbf{R}$,

$$\dim\big(\text{Col}(\mathbf{R}^\top)\big) \leq \dim\big(\text{Col}(\mathbf{A})\big)$$

Similarly, we can obtain that

$$\dim\big(\text{Col}(\mathbf{A})\big) \leq \dim\big(\text{Col}(\mathbf{R}^\top)\big) \leq \dim\big(\text{Col}(\mathbf{A}^\top)\big).$$

Therefore,

$$\dim\big(\text{Col}(\mathbf{A})\big) = \dim\big(\text{Col}(\mathbf{R}^\top)\big) = \dim\big(\text{Col}(\mathbf{A}^\top)\big).$$

□

iii Rank-nullity theorem (秩—零化度定理)

For any $m \times n$ matrix $\mathbf{A}$:

$$\text{rank}(\mathbf{A}) + \dim\big(\ker(\mathbf{A})\big) = n$$

Proof. Let $\mathbf{A}$ be an $m \times n$ matrix with rank r. We will show that there exists a set of $n - r$ linearly independent solutions to the homogeneous system $\mathbf{Ax} = \mathbf{0}_m$ where $\mathbf{x}$ is an n-dimensional column vector.

To do this, we will produce an $n \times (n - r)$ matrix $\mathbf{X}$ whose columns form a basis of the null space of $\mathbf{A}$.

Without loss of generality, assume that the first r columns of $\mathbf{A}$ are linearly independent. So, we can write

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 \end{pmatrix},$$

where

- $\mathbf{A}_1$ is an $m \times r$ matrix with r linearly independent column vectors, and
- $\mathbf{A}_2$ is an $m \times (n - r)$ matrix such that each of its $n - r$ columns is a linear combination of the columns of $\mathbf{A}_1$.

This means that $\mathbf{A}_2 = \mathbf{A}_1\mathbf{B}$ for some $r \times (n - r)$ matrix $\mathbf{B}$, and hence,

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1\mathbf{B} \end{pmatrix}.$$

Let

$$\mathbf{X} = \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix},$$

Then,

$$\mathbf{AX} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1\mathbf{B} \end{pmatrix} \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} = -\mathbf{A}_1\mathbf{B} + \mathbf{A}_1\mathbf{B} = \mathbf{0}_{m\times(n-r)}.$$

Therefore, each of the $n-r$ columns of $\mathbf{X}$ is a solution of $\mathbf{Ax} = \mathbf{0}_m$.

Furthermore, we claim that the $n-r$ columns of $\mathbf{X}$ are linearly independent. We show that $\mathbf{0}_{n-r}$ is the only $(n-r)$-dimensional column vector $\mathbf{u}$ such that

$$\mathbf{Xu} = \mathbf{0}_n$$

$$\begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} \mathbf{u} = \begin{pmatrix} -\mathbf{Bu} \\ \mathbf{u} \end{pmatrix} = \begin{pmatrix} \mathbf{0}_r \\ \mathbf{0}_{n-r} \end{pmatrix}$$

We next prove that any solution of $\mathbf{Ax} = \mathbf{0}_m$ must be in the column space of $\mathbf{X}$. Let the submatrix of the top r rows of $\mathbf{x}$ be $\mathbf{x}_1$ and the submatrix of the bottom $n-r$ rows of $\mathbf{x}$ be $\mathbf{x}_2$.

$$\mathbf{Ax} = \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_1\mathbf{B} \end{pmatrix} \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{pmatrix} = \mathbf{A}_1\mathbf{x}_1 + \mathbf{A}_1\mathbf{B}\mathbf{x}_2 = \mathbf{A}_1(\mathbf{x}_1 + \mathbf{B}\mathbf{x}_2) = \mathbf{0}_m$$

Since the columns of $\mathbf{A}_1$ are linearly independent, $\mathbf{A}_1\mathbf{u} = \mathbf{0}_r$ implies $\mathbf{u} = \mathbf{0}_r$, thus

$$\mathbf{u}_1 + \mathbf{B}\mathbf{u}_2 = \mathbf{0}_r$$

$$\mathbf{u} = \begin{pmatrix} -\mathbf{B} \\ \mathbf{I}_{n-r} \end{pmatrix} \mathbf{u}_2 = \mathbf{X}\mathbf{u}_2$$

This proves that any vector $\mathbf{u}$ that is a solution of $\mathbf{Ax} = \mathbf{0}$ must be a linear combination of the $n-r$ special solutions given by the columns of $\mathbf{X}$. And we have already seen that the columns of $\mathbf{X}$ are linearly independent.

Hence, the column space of $\mathbf{X}$ is the null space of $\mathbf{A}$. Therefore, the nullity of $\mathbf{A}$ is $n-r$. Since r is the rank of $\mathbf{A}$, it follows that $\text{rank}(\mathbf{A}) + \ker(\mathbf{A}) = n$. □

iv Rank-left nullity theorem

For any $m \times n$ matrix $\mathbf{A}$:

$$\text{rank}(\mathbf{A}) + \dim\left(\ker\left(\mathbf{A}^\top\right)\right) = m$$

Proof. Apply rank-nullity theorem on $A^\top$. □

v Full rank (滿秩) and rank deficiency (秩虧)

For any $m \times n$ matrix $\mathbf{A}$:

$$\text{rank}(\mathbf{A}) \leq \min(m, n)$$

Iff

$$\text{rank}(\mathbf{A}) = \min(m, n),$$

A is called to have full rank; otherwise it is called to be rank-deficient (欠秩或秩虧的).

The rank deficiency of a $m \times n$ matrix **A** is defined as

$$\min(m, n) - \operatorname{rank}(\mathbf{A})$$

vi Zero rank

A matrix is a zero matrix iff its rank is zero.

vii Rank preservation

For real matrix,

$$\operatorname{rank}(A) = \operatorname{rank}\left(A^{\mathsf{T}}\right) = \operatorname{rank}\left(AA^{\mathsf{T}}\right) = \operatorname{rank}\left(A^{\mathsf{T}}A\right)$$

For complex matrix,

$$\operatorname{rank}(A) = \operatorname{rank}\left(A^{\mathsf{T}}\right) = \operatorname{rank}\left(\bar{A}\right) = \operatorname{rank}\left(A^{*}\right) = \operatorname{rank}\left(AA^{*}\right) = \operatorname{rank}\left(A^{*}A\right)$$

XI Properties

i Column correspondence property

Let A, B be row-equivalent matrices. If a column j of A is a linear combination of other columns of A, then column j of B is a linear combination of the corresponding columns of B using the same coefficients.

Proof. There exists an invertible matrix E, which is the product of a sequence of $m \times m$ elementary matrices, such that $B = EA$. Let a matrix with subscript denote its submatrix of that column. For every column j

$$B_j = EA_j$$

Suppose

$$A_j = \sum_{i \neq j} c_i A_i$$

$$B_j = EA_j = E \sum_{i \neq j} c_i A_i = \sum_{i \neq j} c_i B_i$$

□

ii Elementary row operation preservations

Suppose E is an $m \times m$ elementary matrix and A is an $m \times n$ matrix. Then A and $B \coloneqq EA$ have the same

- row space (and thus also rank)
- kernel (and thus also nullity)

Proof. Since $B = EA$, each row of B is a linear combination of rows of A, thus $\text{Row}(B) \subseteq \text{Row}(A)$. Since $A = E^{-1}B$, each row of A is a linear combination of rows of B, thus $\text{Row}(A) \subseteq \text{Row}(B)$. Thus $\text{Row}(A) = \text{Row}(B)$.

$Bx = 0 \iff EAx = 0 \iff Ax = E^{-1}0 = 0 \iff EAx = Bx = E0 = 0$, thus $\ker(A) = \ker(B)$. □

The statement does not hold for column space and left kernel.

Proof. Counterexample for column space:

$$A = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

B is obtained by row swapping A, but $\text{Col}(A) = \text{span}\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$ and $\text{Col}(B) = \text{span}\left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ are different.

Counterexample for left kernel:

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$$

B is obtained by adding row 1 of A to row 2, but $\ker\left(A^{\mathsf{T}}\right) = \text{span}\left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\ker\left(B^{\mathsf{T}}\right) = \text{span}\left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ are different. □

iii Row-equivalence is an equivalence relation

Row-equivalence is an equivalence relation.

Proof. Reflexivity: Obviously

Symmetry: Since elementary matrices are invertible

Transitivity: If B is row-equivalent to A with $B = EA$ and C is row-equivalent to B with $C = FB$, then $C = EFA$, thus C is row-equivalent to A. □

iv Rank and number of pivot columns

For a matrix A and its reduced row echelon form R, the followings are equal:

- The rank of A (or R)
- The number of pivot positions of A (or R)
- The number of pivot columns of A (or R)

- The number of pivot rows of A (or R)
- The number of nonzero rows of R
- The number of basic variables in the general solution of any consistent system of linear equations whose coefficient matrix is A (or R)

v Nullity and number of non-pivot columns

For a matrix A and its reduced row echelon form R, the followings are equal:

- The nullity of A (or R)
- The number of non-pivot columns of A (or R)
- The number of free variables in the general solution of any consistent system of linear equations whose coefficient matrix is A (or R)

vi Left nullity and number of non-pivot rows

For a matrix A and its reduced row echelon form R, the followings are equal:

- The left nullity of A (or R)
- The number of non-pivot rows of A (or R)
- The number of zero rows of R
- The number of free variables in the general solution of any consistent system of linear equations whose coefficient matrix is $A^\top$ (or $R^\top$)

vii Existence and uniqueness of reduced row echelon form

Every matrix can be transformed into one and only one matrix in reduced row echelon form by means of a sequence of elementary row operations.

Proof. Existence: By Gauss-Jordan elimination.

Uniqueness:

Let A be a matrix and R be any reduced row-equivalent form of it. Since elementary row operation preserves rank, which is the dimension of column space, which is the number of pivot columns of both A and R, the pivot columns of R are uniquely determined.

Let a matrix with subscript denote its submatrix of that column. For every non-pivot column j, by column correspondence property, since column j of A is a linear combination of pivot columns of A, column j of R is a linear combination of the corresponding columns of A using the same coefficients, which is unique. □

viii Column space test for consistency of linear system

A system of linear equations is consistent iff its constant term vector is in the column space of its coefficient matrix.

Proof. By definition, obviously. □

Thus,

- A homogeneous linear system is always consistent.

  Proof. Zero vector is in all linear subspaces. □

- A linear system whose coefficient matrix's rank is equal to its the number of rows is always consistent.

  Proof. Such matrix's row space is the whole vector space $\mathbb{F}^m$, where m is the number of row. □

ix Invertible coefficient matrix

For a linear system $Ax = b$ with invertible coefficient matrix A and constant term vector b, then $x = A^{-1}b$ is the unique solution to it.

x Solution space is an affine space with kernel as associated vector space

Suppose y is a solution to a linear system $Ax = b$ with coefficient matrix A and constant term vector b, then the set of all solutions of it

$$\{y + k : \ k \in \ker(A)\}$$

Proof. Suppose $z - y \in \ker(A)$.

$$Az = Ay + A(z - y) = Ay = b$$

Suppose $z - y \notin \ker(A)$.

$$Az = Ay + A(z - y) = b + A(z - y) \neq b$$

□

Thus, the solution space is an affine space with kernel as the associated vector space.

xi Zero nullity

For a matrix $A \in \mathbb{R}^{m \times n}$, the followings are equivalent:

- A has zero nullity (i.e., $\operatorname{rank}(A) = n$).
- A linear system whose coefficient matrix is A has at most one solution.

  Proof. If consistent, since the associated vector space $\ker(A)$ of solution space is $\{\mathbf{0}\}$, the solution space has only one element. □

- The only solution to the homogeneous linear system whose coefficient matrix is A is $\mathbf{0}$.
- The columns of the reduced row echelon form of A are the n distinct standard vectors of $\mathbb{R}^n$.

xii Pivot column set is column space basis

The set of pivot columns of a matrix is a basis of its column space.

Proof. For reduced row echelon form, obviously. By column correspondence property and unique existence of reduced row echelon form, the statement holds. □

xiii Invertible iff full rank

A square matrix is invertible iff it has full rank.

Proof. Implies: If A is invertible $Ax = 0$ has the only solution $x = 0$, thus A has nullity 0. By rank-nullity thereom, A has rank n.

Implied by: Since A has full rank, elementary row operations preserve rank, and I_n is the only RREF with full rank, there exists an invertible matrix E, which is the product of a sequence of elementary matrices, such that $EA = I_n$, then $E = A^{-1}$. □

xiv Invertible matrix reduced row echelon form is identity matrix

Let A be an $n \times n$ matrix. Then A is invertible iff its reduced row echelon form is I_n.

Proof. Implies: Since A has full rank, elementary row operations preserve rank, and I_n is the only RREF with full rank, A must have reduced row echelon form I_n.

Implied by: Since elementary row operations preserve rank and I_n has full rank, A has full rank, thus A is invertible. □

xv Invertible matrix must be product of elementary matrix

Let A be an $n \times n$ matrix. Then A is invertible iff it's a product of $n \times n$ elementary matrices.

xvi Algorithm for computing inverse with elementary row operations

Let A be an $n \times n$ matrix. Use elementary row operations to transform $[AI_n]$ into the form $[RB]$ where R is the reduced row echelon form of A. Then either

- $R = I_n$, in which case A is invertible and $B = A^{-1}$; or
- $R \neq I_n$, in which case A is not invertible.

xvii Algorithm for computing inverse right multiplied by another matrix with elementary row operations

Let A be an $n \times n$ matrix and B be an $n \times p$ matrix. Use elementary row operations to transform $[AB]$ into the form $[RC]$ where R is the reduced row echelon form of A. Then either

- $R = I_n$, in which case A is invertible and $C = A^{-1}$; or
- $R \neq I_n$, in which case A is not invertible.

XII Partitioning

i Partitioning

We can break up any matrix by drawing horizontal and vertical lines within the matrix, which divides the matrix into a two-dimensional array of submatrices called blocks. The resulting array is called a partition of the matrix, and the process of forming these blocks is called partitioning.

ii Block multiplication

Suppose two matrices A and B are partitioned into blocks so that the number of blocks in each row of A is the same as the number of blocks in each column of B. Then the matrices can be multiplied according to the usual rules for matrix multiplication, treating the blocks as if they were scalars, provided that the individual products are defined.

iii Matrix multiplication by rows

Let A be an $m \times n$ matrix and B an $n \times p$ matrix. The ith row of AB can be obtained by multiplying the ith row of A by B.

iv Matrix multiplication by outer product

Let A be an $m \times n$ matrix and B an $n \times p$ matrix. The matrix AB is the sum of matrix products of each column of A with the corresponding row of B.

XIII Lower-upper (LU) decomposition or factorization

i Lower triangular matrix

A lower triangular matrix is a matrix whose all entries a_{ij} with $i \leq j$ are the additive identity of 0_F.

ii Upper triangular matrix

An upper triangular matrix is a matrix whose all entries a_{ij} with $i \geq j$ are the additive identity of 0_F.

iii Lower triangular square matrices product

Let A, B be two $n \times n$ lower triangular matrices. Then AB is an $n \times n$ lower triangular matrix.

Proof.

$$(AB)_{ij} = \sum_{k=1}^{n} a_{ik} b_{kj}$$

For $i < j$,

- If $k > i$, then $a_{ik} = 0$.

- If $k \leq i$, then $k < j$, so $b_{kj} = 0$

Thus $(AB)_{ij} = 0$. □

iv Upper triangular square matrices product

Let A, B be two $n \times n$ upper triangular matrices. Then AB is an $n \times n$ upper triangular matrix.

Proof.

$$(AB)_{ij} = \sum_{k=1}^{n} a_{ik} b_{kj}$$

For $i > j$,

- If $k < i$, then $a_{ik} = 0$.
- If $k \geq i$, then $k > j$, so $b_{kj} = 0$

Thus $(AB)_{ij} = 0$. □

v Lower-upper (LU) decomposition or factorization with partial pivoting

LU decomposition with partial pivoting expresses an $m \times n$ matrix A as the product a $m \times m$ lower triangular matrix L whose all diagonal entries being multiplicative identity 1_F and an $m \times n$ upper triangular matrix U:

$$A = LU$$

If Gaussian elimination forward pass on A doesn't need row interchange (i.e., only needs row additions). LU decomposition of A exists. To obtain it, let L_0 be I_m, and perform Gaussian elimination forward pass on A to obtain a row echelon form U of A with l row additions. For each kth operation of adding row i multiplied by c to row j in the process, assign L_{k-1} with the j, ith entry replaced with $-c$ to L_k. Let $L = L_l$. We obtain the LU decomposition of A:

$$A = LU$$

Proof. Observe that U is an $m \times n$ upper triangular matrix. Let the elementary matrices of the operations be $E_1, \ldots, E_l$.

$$U = E_l \ldots E_1 A$$

Since all E_i are invertible,

$$A = E_1^{-1} \ldots E_l^{-1} U$$

Since all E_i are lower triangular with all diagonal entries being multiplicative identity 1_F,

$$L = E_1^{-1} \ldots E_l^{-1}$$

is lower triangular with all diagonal entries being multiplicative identity 1_F.

In forward pass, no two same row additions are performed, thus

$$L_k = E_1^{-1} \ldots E_k^{-1}$$

□

vi Lower-upper (LU) decomposition or factorization with partial pivoting

LU decomposition with partial pivoting expresses an $m \times n$ matrix A multiplied by an $m \times m$ permutation matrix P whose columns and rows are standard basis vectors (i.e., each column and each row has exactly one multiplicative identity 1_F and $m - 1$ additive identity 0_F), as the product a $m \times m$ lower triangular matrix L whose all diagonal entries being multiplicative identity 1_F and an $m \times n$ upper triangular matrix U:

$$PA = LU$$

Let $L_0 = O_m$ and $P_0 = I_m$. Perform Gaussian elimination forward pass on A to obtain a row echelon form U of A with l row additions and interchanges. For each elementary row operation k in it,

- if it is row addition that adds row i multiplied by c to row j, assign L_{k-1} with the j, ith entry replaced with $-c$ to L_k and assign P_{k-1} to P_k;
- if it is row interchange that swaps row i and j, do the same operation on L_{k-1} and P_{k-1} to obtain L_k and P_k.

Let $L = I_m + L_l$ and $P = P_l$. We obtain the LU decomposition with partial pivoting of A:

$$PA = LU$$

Proof. Observe that LU is the LU decomposition of PA, and observe the relation of the forward pass of A and PA to construct the algorithm for sequence L_k. □

vii Solving a system of linear equations using an LU decomposition

The system $Ax = b$ can be rewritten as $LUx = b$. Set $y = Ux$.

First, solve $Ly = b$, called forward substitution. Let $y = (y_1, \dots, y_m)^\top$ and $b = (b_1, \dots, b_m)^\top$. If $m > n$, then

- if $b_{n+1}, \dots, b_m$ are all zero, then the system is consistent, and we can discard the bottom $m - n$ rows,
- otherwise, the system is inconsistent, and there is no solution.

Solve row by row from the top. The top row gives $y_1 = b_1$. For each following row i, substituting results of rows above gives y_i.

Second, solve $Ux = y$, called backward substitution. Let $x = (x_1, \dots, x_n)^\top$. If $m < n$, $x_{m+1}, \dots, x_n$ are free variables. Solve row by row from the bottom. The bottom row gives x_m as a function of the free variables. For each following row i, substituting results of rows below gives x_i as a function of the free variables.

viii Solving a system of linear equations using an LU decomposition with partial pivot

The system $Ax = b$ can be rewritten as $LUx = Pb$ since $P = P^{-1}$. Solve it using the LU decomposition LU.

XIV Flop count of various procedure

Each floating point operation (flop) is an arithmetic operation of floating numbers. Suppose we have a system $Ax = b$ of floating numbers with A being an $n \times n$ invertible matrix. The approximate flop counts are:

- computing the LU decomposition of A: $\frac{2}{3}n^3$.
- solving the system using Gaussian elimination: $\frac{2}{3}n^3$.
- solving the system given the LU decomposition of A: $2n^2$.
- calculating the inverse of A: $2n^3$

Proof. TODO □

3 Matrix Analysis (矩陣分析)

I Eigenvector or characteristic vector (特徵向量) and characteristic polynomial (特徵多項式)

i Eigenvector or characteristic vector (特徵向量) and eigenvalue (本徵值、固有值、特徵值或特徵根)

Given a linear transformation $L : V \to V$, an eigenvector of it is defined as any nonzero vector $v \in V$ such that:

$$\mathbf{A}v = \lambda v,$$

where λ is a scalar, which is called the eigenvalue of $\mathbf{A}$ or the eigenvalue of v.

Obviously, v is an eigenvector implies av is an eigenvector of the same eigenvalue for any nonzero scalar a.

ii Characteristic polynomial

For an $n \times n$ matrix $\mathbf{A}$, the characteristic polynomial of it is defined as

$$p_{\mathbf{A}}(\lambda) := \det\bigl(\lambda \mathbf{I}_n - \mathbf{A}\bigr),$$

where λ is a scalar variable.

A scalar λ is an eigenvalue of $\mathbf{A}$ if and only if

$$p_{\mathbf{A}}(\lambda) = 0.$$

iii Multiplicity of eigenvalues

The algebraic multiplicity of a root λ of the characteristic polynomial of a square matrix is greater than or equal to the number of linearly independent eigenvectors of it whose eigenvalue is λ.

Proof. TODO □

iv Number of eigenvalues

The number of distinct eigenvalues of a square matrix is less than or equal to the number of linearly independent eigenvectors, which less than or equal to n.

Proof. TODO □

v Invertible iff has nonzero eigenvalue

A square matrix is invertible iff it has nonzero eigenvalue.

Proof. TODO □

vi Cayley–Hamilton theorem (凱萊–哈密頓定理)

For a square matrix **A** and its characteristic polynomial $p_{\mathbf{A}}$, the Cayley–Hamilton theorem states that:

$$p_{\mathbf{A}}(\mathbf{A}) = 0$$

vii Diagonalization (對角化)

The diagonalization of an $n \times n$ matrix **A** is the construction of an $n \times n$ matrix **P** and an $n \times n$ diagonal matrix **D** such that:

$$\mathbf{A} = \mathbf{P}^{-1}\mathbf{D}\mathbf{P}$$

An $n \times n$ matrix can be diagonalized if it has n distinct eigenvalues.

The steps of the diagonalization of an $n \times n$ matrix **A** is as follows:

1. For each eigenvalue λ of the matrix **A**, solve an eigenvector v:

$$(\mathbf{A} - \lambda\mathbf{I})\mathbf{v} = 0.$$

2. Compose matrix **P** with each eigenvector of **A** as a column (or row).

Common application:

$$\mathbf{A}^n = (\mathbf{P}^{-1}\mathbf{D}\mathbf{P})^n = \mathbf{P}^{-1}\mathbf{D}^n\mathbf{P}$$

viii Definiteness (正定性)

Let $A \in \mathbb{R}^{n \times n}$. A is called

- Positive definite: iff

$$\forall \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \ \mathbf{x}^\top A \mathbf{x} > 0.$$

- Negative definite: iff

$$\forall \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \ \mathbf{x}^\top A \mathbf{x} < 0.$$

- Positive semi-definite: iff

$$\forall \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \ \mathbf{x}^\top A \mathbf{x} \geq 0.$$

- Negative semi-definite: iff
$$\forall \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \mathbf{x}^\top A \mathbf{x} \leq 0.$$
- Indefinite: otherwise, that is,
$$\exists \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \mathbf{x}^\top A \mathbf{x} > 0$$
and
$$\exists \mathbf{x} \in \mathbb{R}^n, \mathbf{x} \neq 0 : \mathbf{x}^\top A \mathbf{x} < 0.$$

ix Eigenvalues Test of Definiteness for Symmetric Hessians of Scalar Fields

For a symmetric matrix A,

- A is positive definte if all eigenvalues of it are positive.
- A is negative definte if all eigenvalues of it are negative.
- A is positive semidefinte if all eigenvalues of it are nonnegative and at least one eigenvalue of it is positive.
- A is negative semidefinte if all eigenvalues of it are nonpositive and at least one eigenvalue of it is negative.
- A is neutral semidefinte if all eigenvalues of it are zero.
- A is indefinte otherwise.

II Generalized Definition of Inverse

i One-sided inverse (單邊逆)

- Right inverse (右逆): For any $n \times m$ matrix A with rank $\operatorname{rank}(A) = n$, there exists an $m \times n$ matrix A_{R}^{-1}, called the right inverse of **A**, such that
$$A A_{\mathrm{R}}^{-1} = I_n,$$
where I_n is the $n \times n$ identity matrix.
- Left inverse (左逆): For any $n \times m$ matrix A with rank $\operatorname{rank}(A) = m$, there exists an $m \times n$ matrix A_{L}^{-1}, called the left inverse of **A**, such that
$$A_{\mathrm{L}}^{-1} A = I_m,$$
where I_m is the $m \times m$ identity matrix.

Properties:

- A matrix doesn't necessarily have a right inverse and doesn't necessarily have a left inverse.
- A square matrix has a right inverse if and only if it has left inverse if and only if it is invertible.
- For any invertible matrix, there exists only one right inverse and one left inverse, which are its multiplicative inverse.

ii Generalized inverse or g-inverse (廣義逆)

A matrix $A^{\mathrm{g}} \in \mathbb{C}^{n\times m}$ is a generalized inverse of a matrix $A \in \mathbb{C}^{m\times n}$ if:

$$AA^{\mathrm{g}}A = A.$$

Properties:

- For all matrices, there exists at least one generalized inverse, but it's not necessarily unique.
- For any invertible matrix, there exists only one generalized inverse, which is its multiplicative inverse.

iii Reflexive generalized inverse (自反廣義逆)

A reflexive generalized inverse of a matrix **A** is defined as a matrix $\mathbf{A}^g$ such that:

$$\mathbf{A}\mathbf{A}^g\mathbf{A} = \mathbf{A}$$

$$\mathbf{A}^g\mathbf{A}\mathbf{A}^g = \mathbf{A}^g$$

Properties:

- For all matrices, there exists at least one reflexive generalized inverse, but it's not necessarily unique.
- For any invertible matrix, there exists only one reflexive generalized inverse, which is its multiplicative inverse.

iv Moore–Penrose inverse (摩爾—彭若斯廣義逆) or pseudoinverse (偽逆或擬反)

A Moore–Penrose inverse of a matrix **A** is defined as a matrix $\mathbf{A}^+$ such that:

$$\mathbf{A}\mathbf{A}^+\mathbf{A} = \mathbf{A}$$

$$\mathbf{A}^+\mathbf{A}\mathbf{A}^+ = \mathbf{A}^+$$

$$\left(\mathbf{A}\mathbf{A}^+\right)^* = \mathbf{A}\mathbf{A}^+$$

$$\left(\mathbf{A}^+\mathbf{A}\right)^* = \mathbf{A}^+\mathbf{A}$$

Properties:

- For all matrices, there exists only one Moore–Penrose inverse.
- For any invertible matrix, there exists only one Moore–Penrose inverse, which is its multiplicative inverse.

4 Transformation Matrices

I Groups

i General linear group

The general linear group of degree n over some field F is the set of $n \times n$ invertible matrices over F, together with the operation of matrix multiplication, denoted as $\mathrm{GL}(n, F)$ or $\mathrm{GL}_n(F)$, or simply $\mathrm{GL}(n)$ if the field is understood. It is a topological group and a Lie group under operation $+$.

The Lie algebra of a general linear group is denoted as the symbol of the general linear group with GL replaced with $\mathfrak{gl}$.

ii Orthogonal group or general orthogonal group

The orthogonal group or general orthogonal group of degree n over some field F is the set of $n \times n$ invertible matrices over F such that the inverse equals the transpose, together with the operation of matrix multiplication, denoted as $\mathrm{O}(n, F)$ or $\mathrm{O}_n(F)$, or simply $\mathrm{O}(n)$ if the field is understood. An element of an orthogonal group is called an orthogonal matrix. It is a topological group and a Lie group under operation $+$.

iii Special orthogonal group or rotation group

The special orthogonal group or rotation group of degree n over some field F is a subgroup of the orthogonal group of degree n over some field F such that for any matrix in it, the determinant is 1, that is, it preserves the orientation, denoted as $\mathrm{SO}(n, F)$ or $\mathrm{SO}_n(F)$, or simply $\mathrm{SO}(n)$ if the field is understood. An element of a special orthogonal group is called an rotational matrix. It is a topological group and a Lie group under operation $+$.

The Lie algebra of a special orthogonal group is denoted as the symbol of the special orthogonal group with SO replaced with $\mathfrak{so}$.

II Transformation matrices in two-dimensional space

i Degenerate matrices

Degenerate to 1 dimension ($(a \vee b) \wedge (d \vee e) \neq 0$):

$$\begin{bmatrix} a & b \\ ca & cb \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ax + by \\ c(ax + by) \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ d & e \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ dx + ey \end{bmatrix}$$

Degenerate to 0 dimension:

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

ii Stretching (伸縮/縮放/拉伸)

A stretch is a linear transformation which enlarges all distances in a particular direction by a constant factor k but does not affect distances in the perpendicular direction. (Note that if $k > 1$, then this really is a ”stretch”; if $k < 1$, it is technically a ”compression”, but we still call it a stretch. Also, if $k = 1$, then the transformation is an identity, i.e. it has no effect)

The matrix associated with a stretch by a factor h along the x-axis and a factor k along the y-axis is given by:

$$\begin{bmatrix} h & 0 \\ 0 & k \end{bmatrix}$$

iii Shearing (推移)

A shearing in the x-direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$$

A shearing in the y-direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$$

iv Rotation (旋轉)

A rotation by an angle θ counterclockwise about the origin is given by:

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

v Reflection (鏡射/反射)

A reflection about a line that passes the original and with $(a, b) = \sqrt{a^2 + b^2}(\cos\theta, \sin\theta)$ being a vector in the direction of it is given by:

$$\begin{bmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{bmatrix} = \frac{1}{a^2 + b^2} \begin{bmatrix} a^2 - b^2 & 2ab \\ 2ab & b^2 - a^2 \end{bmatrix}$$

vi Orthogona projection (正射影)

An orthogonal projection about a line that passes the original and with $(a, b) = \sqrt{a^2 + b^2}(\cos\theta, \sin\theta)$ being a vector in the direction of it is given by:

$$\begin{bmatrix} \cos^2(\theta) & \cos(\theta)\sin(\theta) \\ \cos(\theta)\sin(\theta) & \sin^2(\theta) \end{bmatrix} = \frac{1}{a^2 + b^2} \begin{bmatrix} a^2 & ab \\ ab & b^2 \end{bmatrix}$$

vii Exmaple 1

Question: The cricle C : $x^2 + y^2 = 1$ is transformed by $\mathbf{A} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ to obtain a new curve C′. Find the equation of C′ and its area.

Answer:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{A} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ 3y \end{bmatrix}$$

$$\begin{cases} x' = 2x \\ y' = 3y \end{cases} \Rightarrow \begin{cases} x = \dfrac{x'}{2} \\ y = \dfrac{y'}{3} \end{cases}$$

$$\left(\frac{x'}{2}\right)^2 + \left(\frac{y'}{3}\right)^2 = 1 \Rightarrow \frac{x'^2}{4} + \frac{y'^2}{9} = 1$$

$$\mathrm{C}' : \frac{x'^2}{4} + \frac{y'^2}{9} = 1$$

The area of C′ is $\pi \cdot 2 \cdot 3 = 6\pi$. Since $\det(\mathbf{A}) = 6$, and the area of C is π, the area of C′ equals $\det(\mathbf{A})$ multiplied by the area of C.

viii Exmaple 2

Question: The line **L** : $4x - 3y = 5$ is transformed in the x-direction dependent on the y coordination by twice and then in the y-direction dependent on the x coordination by negative theee times to obtain a new line **L**′. Find the equation of **L**′.

Answer:

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ -3 & 1 \end{bmatrix}$$

$$\mathbf{A}^{-1} = \frac{1}{7} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \mathbf{A}^{-1} \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \dfrac{x' - 2y'}{7} \\ \dfrac{3x' + y'}{7} \end{bmatrix}$$

$$\mathbf{L}' : 4\frac{x - 2y}{7} + 3\frac{3x + y}{7} = 5$$

$$\mathbf{L}' : 5x + 11y + 35 = 0$$

ix Example 3

Question: Reflect the point P$(5, 2)$ over the line **M** : $x - y + 1 = 0$ to obtain a new point Q. Find Q.

Answer:

Shift one unit to the right: **M** becomes **M**′ : $x - y = 0$ and C becomes C′$(6, 2)$.

The reflection matrix is:

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\mathrm{D}' = \mathbf{A}\mathrm{C}' = (2, 6)$$

Shift one unit to the left: $\mathrm{D}(1, 6)$

x Example 4

Question: The line $x + y - 2 = 0$ is transformed by $\mathbf{A} = \begin{bmatrix} a & b \\ 1 & 2 \end{bmatrix} \in \mathbb{R}^{2\times 2}$ to obtain a new line $2x + 3y - 4 = 0$. Find a and b.

Answer:

$$\mathbf{A}\begin{bmatrix} x \\ -x + 2 \end{bmatrix} = \begin{bmatrix} x' \\ \dfrac{-2x' + 4}{3} \end{bmatrix}$$

$$x - 2x + 4 = \frac{-2x' + 4}{3}$$

$$x' = \frac{3}{2}x - 4$$

$$ax - bx - 2b = x' = \frac{3}{2}x - 4$$

$$(a, b) = (-\frac{1}{2}, -2)$$

III Transformation matrices in three-dimensional space

i Stretching (伸縮/縮放/拉伸)

The matrix associated with a stretch by a factor h along the x-axis, a factor k along the y-axis, and a factor w along the z-axis is given by:

$$\begin{bmatrix} h & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & w \end{bmatrix}$$

ii Shearing (推移)

A shearing in the x-direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & k & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the x-direction dependent on the z coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & k \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the y-direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ k & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the y-direction dependent on the z coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & k \\ 0 & 0 & 1 \end{bmatrix}$$

A shearing in the z-direction dependent on the x coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ k & 0 & 1 \end{bmatrix}$$

A shearing in the z-direction dependent on the y coordination by a factor k is given by:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & k & 1 \end{bmatrix}$$

iii Rotation (旋轉)

A rotation by an angle θ counterclockwise about a line that passes the original and with (a, b, c) being a vector in the direction of it is given by:

$$\frac{1}{a^2+b^2+c^2}\begin{bmatrix} a^2(1-\cos\theta)+\cos\theta & ab(1-\cos\theta)-c\sin\theta & ac(1-\cos\theta)+b\sin\theta \\ bc(1-\cos\theta)+c\sin\theta & b^2(1-\cos\theta)+\cos\theta & bc(1-\cos\theta)-a\sin\theta \\ ac(1-\cos\theta)-b\sin\theta & bc(1-\cos\theta)+a\sin\theta & c^2(1-\cos\theta)+\cos\theta \end{bmatrix}$$

iv Rodrigues' rotation formula

If $\mathbf{v}, \mathbf{k} \in \mathbb{R}^3$ and $\mathbf{k}$ is a unit vector describing an axis of rotation about which $\mathbf{v}$ rotates by an angle θ according to the right hand rule, the Rodrigues formula for the rotated vector $\mathbf{v}_r$ is

$$\mathbf{v}_r = \mathbf{v}\cos\theta + (\mathbf{k}\times\mathbf{v})\sin\theta + \mathbf{k}(\mathbf{k}\cdot\mathbf{v})(1-\cos\theta).$$

The skew-symmetric matrix $\mathbf{K}$ in Lie algebra $\mathfrak{so}(3)$ that generates the one-parameter subgroup of $\mathrm{SO}(3)$:

$$\mathbf{R}(\theta) = \exp(\theta\mathbf{K}),$$

such that

$$\mathbf{v}_r = \mathbf{R}(\theta)\mathbf{v},$$

is given by

$$\mathbf{K}\mathbf{v} = \mathbf{k}\times\mathbf{v}, \quad \forall\mathbf{v}\in\mathbb{R}^3,$$

and

$$\mathbf{R}(\theta) = \mathbf{I} + \sin\theta\mathbf{K} + (1-\cos\theta)\mathbf{K}^2,$$

where $\mathbf{I}$ is the 3×3 identity matrix.

v Reflection (鏡射/反射)

A reflection about a plane $ax + by + cz = 0$ is given by:

$$\mathbf{I} - \frac{1}{a^2 + b^2 + c^2} \begin{bmatrix} 2a^2 & 2ab & 2ac \\ 2ab & 2b^2 & 2bc \\ 2ac & 2bc & 2c^2 \end{bmatrix},$$

where $\mathbf{I}$ is the 3×3 identity matrix.

vi Orthogona projection (正射影)

An orthogonal projection about a plane $ax + by + cz = 0$ is given by:

$$\mathbf{I} - \frac{1}{a^2 + b^2 + c^2} \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$$

where $\mathbf{I}$ is the 3×3 identity matrix.