Topology

沈威宇

November 8, 2026

Contents

1 Topology (拓樸學). 1

Ⅰ Basics 1

i Topological space (拓樸空間). 1

ii Complement 1

iii Homeomorphism (同胚) or isomorphism (同構) 1

iv Open set (開集) 1

v Open neighborhood (開鄰域) 1

vi Interior (內部) of subset 2

vii Neighborhood (鄰域) 2

viii A set is open iff all points in it has neighborhood that is a subset of it 2

ix Subspace topology 2

x Topological embedding 2

xi Limit point (極限點), cluster point, or accumulation point (集積點) . 2

xii Closure (閉包). 2

xiii Open cover 3

xiv Having open cover iff open set 3

xv Closed set (閉集) 3

xvi Set of all closed sets 3

xvii A subset is a closed set iff its closure is itself 3

xviii Closure as smallest intersection of closed superset 4

xix Clopen set 4

xx Borel set (博雷爾集) 4

xxi Filter 5

xxii Base or basis (基) 5

xxiii Prefilter or filter base 5

xxiv Quotient space (商空間) 5

xxv Locally finite cover 5
xxvi Dense (稠密) subset 5
xxvii Comparison of topologies 5
xxviii Product topology or Tychonoff topology 6
II Type of topological spaces 6
i Distinct 6
ii Kolmogorov space (柯爾莫果洛夫空間) or T_0 space 6
iii Kolmogorov quotient (space) 6
iv Fréchet space (弗蘭歇空間), accessible space, or T_1 space 6
v Symmetric space or R_0 space 7
vi Hausdorff space (郝斯多夫空間), separated space (分離空間), or T_2 space 7
vii Urysohn space or $T_{2½}$ space 7
viii Completely Hausdorff space, functionally Hausdorff space, or completely T_2 space 7
ix Regular space (正則空間) 7
x Regular Hausdorff space or T_3 space 8
xi Completely regular space 8
xii Completely regular Hausdorff space, Tychonoff space, completely T_3 space, or $T_{3½}$ space 8
xiii Normal space (正規空間) 8
xiv Urysohn's lemma 8
xv Normal Hausdorff space or T_4 space 8
xvi Completely normal space or hereditarily normal space 9
xvii Completely normal Hausdorff space, T_5 space, or completely T_4 space 9
xviii Perfectly normal space 9
xix Perfectly normal Hausdorff space, T_6 space, or perfectly T_4 space 9
xx Connected space (連通空間) 9

xxi Connected space iff empty set and space are the only two clopen sets 9
xxii Path-connected space (路徑連通空間) 10
xxiii Simply connected space or 1-connected space (單連通空間) . . . 10
xxiv Compact (緊緻) (subset) 10
xxv Compact space 10
xxvi (Weakly) locally compact space 10
xxvii Strongly locally compact or locally relatively compact space . . 10
xxviii σ-compact space 10
xxix σ-locally compact space 10
xxx Second-countable space or completely separable space 10
III Metric space (度量空間 or 賦距空間) 11
i Metric space 11
ii Cauchy sequence (柯西序列) 11
iii Complete metric space (完備度量空間) or Cauchy space (柯西空間) . 11
iv Metric induced by norm 11
v Complete subset or Cauchy subset 11
vi (Open) ball (開球) 12
vii Closed ball (閉球) 12
viii Bounded (有界) subset 12
ix Totally bounded or precompact subset 12
x ε-net 12
xi A compact subset of a Hausdorff space is closed 12
xii A closed subset of a compact set is compact 13
xiii A compact subset of a metric space is bounded 13
xiv A complete subset of a complete metric space is closed 13
xv A closed subset of a complete metric space is complete 14
xvi A closed and bounded subset of an Euclidean space is compact . 14
xvii Heine–Borel Theorem (海涅-博雷爾定理) 14

xviii A complete and totally bounded subset of a complete metric space is compact 14
xix A totally bounded subset is bounded 15
xx A bounded subset of a finite-dimensional Banach space is totally bounded 16
IV Normed space 16
i Norm-induced topology 16
ii Norm-induced topology is Hausdorff 16
iii Equivalence of norms 17
iv Norm equivalence theorem 17
v Unit sphere is closed 18
vi All norms of real or complex finite-dimensional space are equivalent 18
vii Linear map from finite-dimensional space is continuous and bounded 19
V Net (網) 20
i Net (網) 20
ii Eventually or residually. 20
iii Limit of nets 20
iv Cofinal (共尾) or frequent subset 20
v Cofinally (共尾地) or frequently 21
vi Cluster point, or accumulation point (集積點) 21
vii Subnet or Willard-subnet 21
viii Ultranet or universal net 21
ix In Hausdorff spaces 21
x Willard-subnet limit and cluster point theorems 21
VI Manifold 22
i Euclidean neighborhood 22
ii Locally Euclidean space. 22
iii (Topological) manifold (流形) 22

iv (Coordinate) chart 22
v Atlas 22
vi Transition function or transition map 23
vii Differentiable and smooth 23
viii Diffeomorphism 24
ix Local diffeomorphism 24
VII Combination with abstract algebra 24
i Topological group (拓樸群) 24
ii Topological field (拓樸域) 25
VIII Discrete space 25
i Discrete topology 25
ii Discrete (topological) space 25
iii Discrete subspace 25

1 Topology (拓樸學)

I Basics

i Topological space (拓樸空間)

A topological space consists of a set X and a topology $\mathcal{T}$ on it, denoted as $(X, \mathcal{T})$. Where the set X is the set of points in the space, and the topology $\mathcal{T}$ is the set of subsets of X that satisfies:

- Empty set and space itself is open:

$$\emptyset, X \in \mathcal{T}$$

- Closed under arbitrary unions:

$$\forall \mathcal{S} \subseteq \mathcal{T} : \bigcup_{A \in \mathcal{S}} A \in \mathcal{T}$$

- Closed under finite intersection:

$$\forall \mathcal{S} \subseteq \mathcal{T}, |\mathcal{S}| \in \mathbb{N} : \bigcap_{A \in \mathcal{S}} A \in \mathcal{T}$$

ii Complement

For a subset A of a topological space $(X, \mathcal{T})$, the complement A^c of A is defined as $X \setminus A$.

iii Homeomorphism (同胚) or isomorphism (同構)

Topological spaces $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ are called homeomorphic if there exists a mapping $f : X \to Y$ between them such that f is bijective and continuous and f^{-1} is continuous, written as $X \cong Y$, and f is called a homeomorphism between them.

iv Open set (開集)

In a topological space $(X, \mathcal{T})$, an open subset $O \subseteq X$ is defined to be

$$O \in \mathcal{T}.$$

v Open neighborhood (開鄰域)

An open neighborhood of a point P in a topological space X is any open subset $O \subseteq X$ such that $P \in O$.

An open neighborhood of a subset V in a topological space X is any open subset $O \subseteq X$ such that $V \subseteq O$.

vi Interior (內部) of subset

The interior of a subset S of a topological space X, denoted by $\operatorname{int}_X S$, $\operatorname{int} S$, $\operatorname{int}_X(S)$, $\operatorname{int}(S)$, or S°, is defined as

$$\operatorname{int}(S) = \{x \in S \mid \exists \text{ open } U \subseteq X \text{ s.t. } x \in U \subseteq S\}.$$

vii Neighborhood (鄰域)

A subset U is a neighborhood of a point P if and only if there exists an open set O such that $P \in O \subseteq U$.

A subset U is a neighborhood of a subset V if and only if $V \subseteq \operatorname{int}(U)$.

viii A set is open iff all points in it has neighborhood that is a subset of it

For a topological space $(X, \mathcal{T})$, a subset A is in $\mathcal{T}$ iff for all $x \in A$, there exists neighborhood N of x such that $N \subseteq A$.

Proof. If: Since $\mathcal{T}$ is closed under arbitrary union, pick a such neighborhood for each $x \in A$.

Only if: A is such neighborhood. □

ix Subspace topology

The subspace topology of a topological space $(X, \mathcal{T})$ in a subset I of X is defined as the set

$$\mathcal{I} = \{U \cap I | U \in \mathcal{T}\}.$$

x Topological embedding

A topological embedding is a homeomorphism onto its image with subspace topology.

xi Limit point (極限點), cluster point, or accumulation point (集積點)

Let S be a subset of a topological space X. A point $x \in X$ is called a limit point, cluster point, or accumulation point of S iff for every neighborhood N of x, there exists $y \in N \setminus \{x\}$ such that $y \in S$.

xii Closure (閉包)

The closure of a subset S of points in a topological space consists of all points in S together with all limit points of S.

xiii Open cover

For a subset A of a topological space $(X, \mathcal{T})$, an open cover A is a cover C of A such that $C \subseteq \mathcal{T}$.

xiv Having open cover iff open set

For a subset A of a topological space $(X, \mathcal{T})$, A has open cover iff A is open.

Proof. If: $\{A\}$ is an open cover.

Only if: $\mathcal{T}$ is closed under arbitrary union. □

xv Closed set (閉集)

A subset A of a topological space $(X, \mathcal{T})$ is called closed iff its complement $X \setminus A \in \mathcal{T}$.

xvi Set of all closed sets

For an topological space $(X, \mathcal{T})$, suppose $\mathcal{U}$ is the set of all closed subsets $\mathcal{U}$, then

- Empty set and space itself is closed:

$$\varnothing, X \in \mathcal{U}$$

- Closed under finite unions:

$$\forall \mathcal{S} \subseteq \mathcal{U}, |\mathcal{S}| \in \mathbb{N} : \bigcup_{A \in \mathcal{S}} A \in \mathcal{U}$$

- Closed under arbitrary intersection:

$$\forall \mathcal{S} \subseteq \mathcal{U} : \bigcap_{A \in \mathcal{S}} A \in \mathcal{U}$$

Proof. Each corresponds to each condition of a topology. □

xvii A subset is a closed set iff its closure is itself

A subset A of a topological space $(X, \mathcal{T})$ is closed iff its closure is itself.

Proof. If: Assume for contradiction that a limit point x of A is in A^c. Then all neighborhoods N of x is such that

$$N \cap A \neq \varnothing$$

Since $x \in A^c$ and $A^c \in \mathcal{T}$, A^c is a neighborhood of x, but

$$A^c \cap A = \varnothing$$

Only if: For all $x \in A^c$, there exists a neighborhood N of x such that

$$N \cap A = \varnothing,$$

that is,

$$N \subseteq A^c$$

By that a set is open if all points in it has neighborhood that is a subset of it, A^c is open. □

xviii Closure as smallest intersection of closed superset

For a subset A of a topological space $(X, \mathcal{T})$ with set of all closed set $\mathcal{U}$, the closure of A is

$$\bigcap_{F \in \mathcal{U} \wedge A \subseteq F} F$$

Proof. Let

$$C \coloneqq \bigcap_{F \in \mathcal{U} \wedge A \subseteq F} F$$

and D the closure of A.

$C \subseteq D$: By that a closure is closed and the definition of C.

$D \subseteq C$: Assume for contradiction that there exists a limit point x of A such that $x \notin C$. Then, C^c is open and contains x, which contradicts that C is closed. □

xix Clopen set

A subset A of a topological space $(X, \mathcal{T})$ is called clopen iff it is both open and closed.

xx Borel set (博雷爾集)

A Borel set B is any set in a topological space $(X, \mathcal{T})$ that can be formed from open sets through the operations of countable union, countable intersection, and relative complement, that is,

1. Every element of $\mathcal{T}$ is a Borel set.
2. For any indexed family $(B_i)_{i=1}^{\infty}$ of Borel sets, $\bigcap_{i=1}^{\infty} B_i$ is a Borel set.
3. If B, C are Borel sets, then $B \setminus C$ is a Borel set.
4. Any other set is not a Borel set.

xxi Filter

A filter on a set X is a family $\mathcal{B}$ of subsets of X such that:

1. $X \in \mathcal{B}$.
2. $\varnothing \in \mathcal{B}$.
3. $A \in \mathcal{B} \wedge B \in \mathcal{B} \implies A \cap B \in \mathcal{B}$.
4. $A \subseteq B \subseteq X \wedge A \in \mathcal{A} \in \mathcal{B} \implies B \in \mathcal{B}$.

xxii Base or basis (基)

Given a topological space $(X, \mathcal{T})$, a base (or basis) for the topology $\mathcal{T}$ (also called a base for X if the topology is understood) is a family $\mathcal{B} \subseteq \mathcal{T}$ of open sets such that every open set of the topology can be represented as the union of some subfamily of $\mathcal{B}$.

The topology generated by a base $\mathcal{B}$, generally denoted by $\tau(\mathcal{B})$ can be defined to be follows: A subset $O \subseteq X$ is to be declared as open, if for all $x \in O$, there exists some $B \in \mathcal{B}$ such that $x \in B \subseteq O$.

xxiii Prefilter or filter base

$\mathcal{B}$ is called a prefilter if its upward closure $\uparrow \mathcal{B}$ is a filter.

xxiv Quotient space (商空間)

Let X be a topological space and $\cong$ be an equivalence relation on it. The quotient space Y of X by $\cong$ is the quotient set of X by $\cong$ equipped with the quotient topology, defined as $U \subseteq Y$ is open in the quotient topology on Y if and only if $\{x \in X \mid [x] \in U\}$ is open in X and $V \subseteq Y$ is closed in the quotient topology on Y if and only if $\{x \in X \mid [x] \in V\}$ is closed in X.

xxv Locally finite cover

Let X be a topological space, and $\{U_i \mid i \in I\}$ be a cover of X. We say that U is locally finite if for all $x \in X$, there exists $V \ni x$ that is open in X such that the set $\{i \in I \mid V \cap U_i \neq \varnothing\}$ is finite.

xxvi Dense (稠密) subset

A subset A of a topological space X is said to be a dense subset of X if for every $x \in X$ and every neighborhood U of x, $U \cap A \neq \varnothing$.

xxvii Comparison of topologies

Let τ_1 and τ_2 be two topologies on a set X.

τ_1 is said to be a coarser (weaker or smaller) topology than τ_2, and τ_2 is said to be a finer (stronger or larger) topology than τ_1, iff τ_1 is a subset of τ_2.

τ_1 is said to be a strictly coarser topology than τ_2, and τ_2 is said to be a strictly finer topology than τ_1, iff τ_1 is a proper subset of τ_2.

The binary relation $\subseteq$ defines a partial ordering relation on the set of all possible topologies on X.

xxviii Product topology or Tychonoff topology

The product topology or Tychonoff topology on a set $X = \prod_{i=1}^{n} X_i$ where X_i are topological spaces and $n > 1$ is defined as the coarsest topology on X such that all ith projection map are continuous. X with the product topology on X is called a product space. Unless otherwise specified, product topology is assumed on a Cartesian product of topological spaces.

II Type of topological spaces

i Distinct

Point x and point y in a topological space are distinct if and only if they are not the same point.

ii Kolmogorov space (柯爾莫果洛夫空間) or T_0 space

Two points in a topological space are topological distinguishable if and only if there exists an open set that contains one of them but not the other.

A topological space is called a Kolmogorov space or T_0 space if and only if any two distinct points in it are topological distinguishable. This condition is called the T_0 condition or the first separation theorem.

iii Kolmogorov quotient (space)

The Kolmogorov quotient (space) of a topological space is the quotient space of it under the equivalence relation topological indistinguishablity.

iv Fréchet space (弗蘭歇空間), accessible space, or T_1 space

Two points x and y in a topological space are separated if and only if there exists open sets U, V such that $x \in U \wedge y \notin U \wedge y \in V \wedge x \notin V$.

A topological space is called a Fréchet space, accessible space, or T_1 space if and only if any two distinct points in it are separated. This condition is called the T_1 condition

or the second separation theorem. A topological space is a T_1 space implies it is a T_0 space.

v Symmetric space or R_0 space

A topological space is called a symmetric space or R_0 space if and only if any two topological distinguishable points in it are separated.

vi Hausdorff space (郝斯多夫空間), separated space (分離空間), or T_2 space

Points x and y in a topological space are separated by neighborhoods if and only if there exists a neighborhood U of x and a neighborhood V of y such that U and V are disjointed.

A topological space is called a Hausdorff space, separated space, or T_2 space if and only if any two distinct points in it are separated by neighborhoods. This condition is called the T_2 condition or the third separation theorem. A topological space is a T_2 space implies it is a T_1 space.

vii Urysohn space or $T_{2½}$ space

Points x and y in a topological space are separated by closed neighborhoods if and only if there exists a closed neighborhood U of x and a closed neighborhood V of y such that U and V are disjointed.

A topological space is called a Urysohn space or $T_{2½}$ space if and only if any two distinct points in it are separated by closed neighborhoods. A topological space is a $T_{2½}$ space implies it is a T_2 space.

viii Completely Hausdorff space, functionally Hausdorff space, or completely T_2 space

Points x and y in a topological space X are separated by a (continuous) function if and only if there exists a continuous function $f : X \to [0, 1]$ such that $f(x) = 0$ and $f(y) = 1$.

A topological space is called a completely Hausdorff space, functionally Hausdorff space, or completely T_2 space if and only if any two distinct points in it are separated by a continuous function. A topological space is a completely T_2 space implies it is a $T_{2½}$ space.

ix Regular space (正則空間)

A topological space X is called a regular space if and only if for any closed set F and any point $x \notin F$, there exists a neighbourhood U of x and a neighbourhood V of F such that U and V are disjointed.

x Regular Hausdorff space or T_3 space

A topological space is called a regular Hausdorff space or T_3 space if and only if it is a regular space and a Hausdorff space.

xi Completely regular space

A closed set F and a point $x \notin F$ in a topological space X are separated by a (continuous) function if and only if there exists a continuous function $f : X \to [0, 1]$ such that $f(x) = 0$ and $F \subseteq f^{-1}(\{1\})$.

A topological space X is called a completely regular space if and only if any closed set F and any point $x \notin F$ in it are separated by a (continuous) function. A topological space is a completely regular space implies it is a regular space.

xii Completely regular Hausdorff space, Tychonoff space, completely T_3 space, or $T_{3½}$ space

A topological space X is called a completely regular Hausdorff space, Tychonoff space, completely T_3 space, or $T_{3½}$ space if and only if it is a completely regular space and a Hausdorff space. A topological space is a completely T_3 space implies it is a completely T_2 space.

xiii Normal space (正規空間)

Disjoint closed sets E and F in a topological space are separated by neighborhoods if and only if there exists a neighborhood U of E and a neighborhood V of F such that U and V are disjointed.

A topological space is called a normal space if and only if any two distinct closed sets in it are separated by neighborhoods. A topological space is a normal space implies it is a regular space.

xiv Urysohn's lemma

Two closed sets E and F in a topological space X are separated by a (continuous) function if and only if there exists a continuous function $f : X \to [0, 1]$ such that $\subseteq f^{-1}(\{0\})$ and $F \subseteq f^{-1}(\{1\})$.

The Urysohn's lemma states that, a topological space is a normal space if and only if any two disjoint closed sets in it are separated by a (continuous) function.

xv Normal Hausdorff space or T_4 space

A topological space is called a normal Hausdorff space or T_4 space if and only if it is a normal space and a T_1 space. A topological space is a normal Hausdorff space implies

it is a regular Hausdorff space.

xvi Completely normal space or hereditarily normal space

A topological space is called a completely normal space or hereditarily normal space if and only if any subspace of it is a normal space.

xvii Completely normal Hausdorff space, T_5 space, or completely T_4 space

A topological space is called a completely normal Hausdorff space, T_5 space, or completely T_4 space if and only if it is a completely normal space and a T_1 space. A topological space is a completely normal Hausdorff space implies it is a normal Hausdorff space and a completely regular Hausdorff space.

xviii Perfectly normal space

Two closed sets E and F in a topological space X are precisely separated by a (continuous) function if and only if there exists a continuous function $f : X \to [0, 1]$ such that $E = f^{-1}(\{0\})$ and $F = f^{-1}(\{1\})$.

A topological space is called a perfectly normal space if and only if any two disjoint closed sets in it are precisely separated by a (continuous) function. A topological space is a perfect normal space implies it is a completely normal space.

xix Perfectly normal Hausdorff space, T_6 space, or perfectly T_4 space

A topological space is called a perfectly normal Hausdorff space, T_6 space, or perfectly T_4 space if and only if it is a perfectly normal space and a T_1 space. A topological space is a perfectly normal Hausdorff space implies it is a completely normal Hausdorff space.

xx Connected space (連通空間)

A topological space is called disconnected if and only if it can be expressed as the union of two disjoint non-empty open sets; otherwise, it is called connected.

xxi Connected space iff empty set and space are the only two clopen sets

A topological space $(X, \mathcal{T})$ is connected iff $\varnothing$ and X are the only two clopen sets.

Proof. If: Assume for contradiction that a non-empty proper subset A of X is clopen. Then A^c is a non-empty clopen proper subset of X too. Then X can be expressed as the union of two disjointed non-empty open sets A and A^c.

Only if: Assume for contradiction that X is disconnected and can be expressed as the union of two disjointed non-empty open sets A and A^c. Then A and A^c are both clopen. □

xxii Path-connected space (路徑連通空間)

A topological space X is called path-connected if and only if for any two points $x, y \in X$, there existes a continuous function $f : [0,1] \to X$ such that $f(0) = x$ and $f(1) = y$. A topological space is path-connected implies it is connected.

xxiii Simply connected space or 1-connected space (單連通空間)

A topological space is called simply connected if and only if it is path-connected, and that for any two points $x, y \in X$ and any two continuous functions $f : [0,1] \to X$ and $g : [0,1] \to X$ such that $f(0) = g(0) = x$ and $f(1) = g(1) = y$, there exists a continuous function $F : [0,1] \times [0,1] \to X$ such that $F(x,0) = f(x)$ and $F(x,1) = g(x)$.

xxiv Compact (緊緻) (subset)

A subset K of a topological space X is called compact iff for every open cover C of K, there exists a finite subcover.

xxv Compact space

A topological space X is called compact iff for every open cover C of X, there exists a finite subcover.

xxvi (Weakly) locally compact space

A topological space X is called (weakly) locally compact iff every point in it has a compact neighbourhood.

xxvii Strongly locally compact or locally relatively compact space

A topological space X is called strongly locally compact or locally relatively compact iff every point in it has a closed compact neighbourhood.

A Hausdorff space is weakly locally compact iff it is strongly locally compact.

xxviii σ-compact space

A topological space X is called σ-compact iff its topology is the union of countably many compact spaces.

xxix σ-locally compact space

A topological space X is called σ-locally compact iff it is σ-compact and locally compact.

xxx Second-countable space or completely separable space

A topological space is second-countable iff it has a countable basis.

III Metric space (度量空間 or 賦距空間)

i Metric space

A metric space is an ordered pair (M, d) where M is a set and d is a metric on M, i.e., a function $d : M \times M \to \mathbb{R}$ satisfying the following axioms for all points $x, y, z \in M$:

1. The distance from a point to itself is zero: $d(x, x) = 0$.
2. (Positivity) The distance between two distinct points is always positive:$x \neq y \implies d(x, y) > 0$.
3. (Symmetry) The distance from x to y is always the same as the distance from y to x: $d(x, y) = d(y, x)$.
4. The triangle inequality: $d(x, z) \leq d(x, y) + d(y, z)$.

ii Cauchy sequence (柯西序列)

Given a metric space (x, d), a sequence of elements of X:

$$x_1, x_2, x_3, \ldots$$

is Cauchy if for every positive real number ε there is a positive integer N such that for all positive integers $m, n > N$:

$$d\left(x_m, x_n\right) < \varepsilon.$$

iii Complete metric space (完備度量空間) or Cauchy space (柯西空間)

A metric space (X, d) is complete if every Cauchy sequence of points in X has a limit that is also in X.

iv Metric induced by norm

For a normed space $(X, \|\cdot\|)$, the metric

$$d(x, y) = \|x - y\|, \quad \forall x, y \in X.$$

is said to be induced by the norm. A normed space is assumed to be equipped with such metric unless otherwise specified.

v Complete subset or Cauchy subset

A subset S of a metric space (X, d) is complete if every Cauchy sequence of points in S has a limit that is also in S.

vi (Open) ball (開球)

In a metric space (X, d), the open ball or ball of radius $r > 0$ and centered at the point $a \in X$, denoted as $B_{<r}(a)$, $B_r(a)$, or $B(a, r)$, is defined as:

$$B_r(a) \coloneqq \{p \in X : d(a, p) < r\}.$$

vii Closed ball (閉球)

In a metric space (X, d), the closed ball of radius $r \geq 0$ and centered at the point $a \in X$, denoted as $B_{\leq r}(a)$, is defined as:

$$B_{\leq r}(a) \coloneqq \{p \in X : d(a, p) \leq r\}.$$

viii Bounded (有界) subset

A subset S of a metric space (M, d) is called bounded if and only if there exists $r > 0$ such that for all $s, t \in S$, we have $d(s, t) < r$.

ix Totally bounded or precompact subset

A subset S of a metric space (M, d) is called totally bounded or precompact if and only if there eixsts $r > 0$ such that there exists a finite collection of open balls of radius r that is a cover of S.

x ε-net

A subset E of a metric space X is an ε-net for another subset A of X if

$$A \subseteq \bigcup_{x \in E} B(x, \varepsilon).$$

If a finite ε-net exists for every $\varepsilon > 0$, then A is totally bounded.

xi A compact subset of a Hausdorff space is closed

If a subset K of a Hausdorff space X is compact, then it must be closed.

Proof. We want to show that K is closed, that is, $X \setminus K$ is open.

For any $a \in X \setminus K$, for each $x \in K$, by the definition of Hausdorff space, there exist a disjoint open set U_x such that $x \in U_x$, $a \in V_x$, and $U_x \cap V_x = \varnothing$.

The collection of $C = \{U_x \mid x \in K\}$ is an open cover of K. Since K is compact, a finite subset D of C is a cover of K.

Corresponding to each $U_i \in D$, we have an open neighborhood V_i of a such that $U_i \cap V_i = \varnothing$. Let $V = \bigcap_{U_i \in D} V_i$. Then V is an open neighborhood of a and is disjoint from $\bigcap_{U_i \in D} U_i \supseteq K$, so

$$V \cap K = \varnothing.$$

For every $a \in X \setminus K$, we found an open neighborhood V of a such that

$$V \subseteq X \setminus K \wedge V \cap K = \varnothing,$$

so K is closed. □

xii A closed subset of a compact set is compact

If K is a compact subset of a topological space X and $F \subseteq K$ is closed, then F is compact.

Proof. Let C be an open cover of F. Then

$$C \cup \{X \setminus F\}$$

is an open cover of K.

Since K is compact, there is a finite subcollection $D = C' \cup \{X \setminus F\}$ of $C \cup \{X \setminus F\}$ where $C' \subseteq C$ that is a cover of K. Then C' is a finite subcollection of C that is a cover of F. Thus F is compact. □

xiii A compact subset of a metric space is bounded

If K is a compact subset of a metric space (X, d), then K is bounded.

Proof. Since K is compact, for any point $p \in X$, there exist a finite collection of open balls

$$\{B(p, n) \mid n \in \mathbb{N} \wedge n \leq m \in \mathbb{N}\}$$

that is a cover of K, then $K \subseteq B(p, m)$, so K is bounded. □

xiv A complete subset of a complete metric space is closed

If a subset A of a complete metric space is compact, them A is closed.

Proof. Since a metric space is necessarily a Hausdorff space, any convergent sequence must have a unique limit. Since A is complete, every Cauchy sequence in A converges to a point in A. Thus, A is closed. □

xv A closed subset of a complete metric space is complete

If a subset A of a complete metric space X is closed, them A is compact.

Proof. Let (x_n) be a Cauchy sequence in A. Since the space is complete, it converges to some $x \in X$. Since A is closed in X, any limit of a convergent sequence in A must be in A. Thus, A is complete. □

xvi A closed and bounded subset of an Euclidean space is compact

If a subset A of an Euclidean space $\mathbb{R}^n$ is closed and bounded, then A is compact.

Proof. Let (x_k) be any sequence in A. We want to show that it has a convergent subsequence with its limit in A.

Since A is bounded, there exists $M > 0$ such that

$$\|x_k\| \leq M, \quad \forall k.$$

So

$$x_k \in B(0, M), \quad \forall k.$$

By the Bolzano–Weierstrass theorem, there exists a subsequence (x_{k_j}) that coverges to some point $x \in \mathbb{R}^n$.

Since A is closed, it is complete. Thus $x \in A$. □

xvii Heine–Borel Theorem (海涅-博雷爾定理)

For a subset S of a complete metric space (X, d), the following two statements are equivalent:

- S is compact;
- S is complete and totally bounded.

xviii A complete and totally bounded subset of a complete metric space is compact

If a subset A of a complete metric space (x, d) is complete and totally bounded, then A it is compact.

Proof. Take any sequence (x_n) in A. We want to find a convergent subsequence whose limit lies in A.

For $\varepsilon = 1$, since A is totally bounded, there exists a finite cover of A by closed balls of radius 1:

$$A \subseteq \bigcup_{i=1}^{N_1} B_1\left(y_i\right).$$

Since (x_n) is infinite but the cover is finite, by the pigeonhole principle, one of these balls must contain infinitely many terms of (x_n). Denote this ball by $B_1\left(y_{i_1}\right)$ and pick the subsequence $\left(x_n^{(1)}\right)$ of (x_n) that is entirely in $B_1\left(y_{i_1}\right)$.

Repeat it for $\varepsilon = \frac{1}{2}, \frac{1}{3}, \ldots$, we can construct a nested sequence of subsequences

$$\left(x_n^{(1)}\right) \supseteq \left(x_n^{(2)}\right) \supseteq \ldots$$

where $\left(x_n^{(k)}\right)$ lies in a closed ball of radius $\frac{1}{k}$, $B_{\frac{1}{k}}\left(y_{i_k}\right)$.

Pick the diagonal sequence $y_k = x_k^{(k)}$. Then

$$y_k \in \left(x_n^{(k)}\right) \subseteq B_{\frac{1}{k}}\left(y_{i_k}\right),$$

and the sequence (y_k) is a Cauchy sequence in A because

$$\forall N \in \mathbb{N} : \; m, n \geq N \implies d(y_m, y_n) \leq \frac{2}{N}.$$

Since A is complete, (y_k) must converges to some point $y \in A$. □

xix A totally bounded subset is bounded

If a subset A of a metric space (X, d) is totally bounded, them A is bounded.

Proof. By total boundedness, there exist finitely many points $x_1, x_2, \ldots x_n \in X$ such that:

$$A \subseteq \bigcup_{i=1}^{n} B(x_i, 1).$$

Pick one of these centers, say x_1.

For any $a \in A$, there exists some $i \in \{1, 2, \ldots n\}$ such that

$$d(a, x_i) < 1.$$

By triangle inequality:

$$d(a, x_1) \leq d(a, x_i) + d(x_i, x_1) \leq 1 + d(x_i, x_1).$$

Let:

$$M = 1 + \max_{i \in \{1,2,\ldots n\}} d(x_i, x_1).$$

So, for all $a \in A$:

$$d(a, x_1) < M.$$

Thus:

$$A \subseteq B(x_1, M).$$

Therefore, A is bounded. □

xx A bounded subset of a finite-dimensional Banach space is totally bounded

If a subset A of a n-dimensional Banach space X with $n \in \mathbb{N}$ is bounded, then A is totally bounded.

Proof. Pick a linear homeomorphism

$$T : \mathbb{R}^n \to X.$$

Define a norm on $\mathbb{R}^n$ by

$$\|v\|_* := \|Tv\|_X, \quad \forall v \in \mathbb{R}^n.$$

By that all norms on a finite-dimensional vector space are equivalent, the Euclidean norm $\|\cdot\|_2$ and $\|\cdot\|_*$ are equivalent, that is, there exist constants $c, C > 0$ such that for every $v \in \mathbb{R}^n$,

$$c\|v\|_2 \le \|v\|_* \le C\|v\|_2. \qquad (1)$$

Let $A \subseteq X$ be bounded. Then $B := T^{-1}(A)$ is bounded with respect to the metric induced by $\|\cdot\|_*$ and, by (1), is bounded with respect to the metric induced by $\|\cdot\|_2$. So there is $R > 0$ with $B_{\|\cdot\|_2}(0, R) \supseteq B$. For any $\varepsilon > 0$ and $\delta := \dfrac{\varepsilon}{C}$, there is a finite collection of $B_{\|\cdot\|_2}(v_i, R)$ where $v_i \in \mathbb{R}^n$ that is a cover of $B_{\|\cdot\|_2}(0, R)$.

Take any $b \in B$. Then $b \in B_{\|\cdot\|_2}(v_i, R)$ for some v_i, so $\|b - v_i\|_2 < \delta$.

By (1), we get

$$\|T(b) - T(v_i)\|_X = \|b - v_i\|_* \le C\|b - v_i\|_2 < C\delta = \varepsilon.$$

Therefore the finite set of all $T(v_i)$ is an ε-net for A. Since $\varepsilon > 0$ was arbitrary, A is totally bounded. □

IV Normed space

i Norm-induced topology

For a vector space and a norm $\|\cdot\|$ for it. The induced topology consists of all unions of open balls of the space under the metric induced by $\|\cdot\|$.

ii Norm-induced topology is Hausdorff

The topology induced by any norm on a vector space is Hausdorff.

Proof. Let $(V, \|\cdot\|)$ be a normed space. We want to show that for any distinct $x, y \in V$, there exist disjoint open neighborhoods of x and y.

$$d = \|x - y\| > 0$$

Choose $r = \dfrac{d}{3}$. Consider $B_r(x)$ and $B_r(y)$. Suppose for contradiction that

$$B_r(x) \cap B_r(y) \neq \varnothing$$

Then there exists $z \in B_r(x) \cap B_r(y)$, so

$$\|x - z\| < r, \quad \|y - z\| < r$$

By triangle inequality,

$$\|x - y\| \leq \|x - z\| + \|y - z\| \leq 2r = \frac{2}{3}\|x - y\|,$$

which is impossible because $\|x - y\| \neq 0$. □

iii Equivalence of norms

Two norms $\|\cdot\|_a$ and $\|\cdot\|_b$ for a vector space V are called equivalent iff there exists positive numbers C and D such that

$$C\|x\|_a \leq \|x\|_b \leq D\|x\|_a$$

for all $x \in V$.

iv Norm equivalence theorem

Two norms $\|\cdot\|_a$ and $\|\cdot\|_b$ for a vector space V induce the same topology iff they are equivalent.

Proof. Let $B_a(x, r)$ and $B_b(x, r)$ denotes open balls under the two norms.

$\Leftarrow$: For any $x \in V$ and $r > 0$, we have

$$B_a\left(x, \frac{r}{D}\right) \subseteq B_b(x, r) \subseteq B_a\left(x, \frac{r}{C}\right)$$

If $\|y - x\|_a < \frac{r}{D}$, then

$$\|y - x\|_b \leq D\|y - x\|_a < r$$

If $\|y - x\|_b < r$, then

$$\|y - x\|_a \leq \frac{1}{C}\|y - x\|_b < \frac{r}{C}$$

Thus every open set under either norm is open under the other norm. Hence the induced topologies coincide.

$\Rightarrow$: Since $B_b(0, 1)$ is open in the topology induced by $\|\cdot\|_a$, there exists $r > 0$ such that

$$B_a(0, r) \subseteq B_b(0, 1).$$

We show that

$$\|x\|_b \leq \frac{2}{r}\|x\|_a$$

for every $x \in V$.

For $x \neq 0$, choose any $\varepsilon > 0$ and set

$$t = \frac{r}{2(\|x\|_a + \varepsilon)}$$

Then $\|tx\|_a < r$, so $\|tx\|_b < 1$. By absolute homogeneity,

$$\|x\|_b < \frac{1}{t} = \frac{2}{r}(\|x\|_a + \varepsilon)$$

Let $\varepsilon \to 0^+$,

$$\|x\|_b \leq \frac{2}{r}\|x\|_a$$

Similarly, there also exists $s > 0$ such that

$$\|x\|_a \leq \frac{2}{s}\|x\|_b$$

Take $C = \frac{s}{2}$ and $D = \frac{2}{r}$. □

v Unit sphere is closed

Let $M = (X, \|\cdot\|)$ be a normed vector space. The unit sphere $S \coloneqq \{x \in X : \|x\| = 1\}$ is closed in M.

Proof.

$$S = B_{\leq 1}(0) \setminus B_1(0)$$

$$S' = B_1(0) \cup (X \setminus B_{\leq 1}(0))$$

which is union of open set. □

vi All norms of real or complex finite-dimensional space are equivalent

For a finite-dimensional vector space V over the real or complex numbers, all norms for it are equivalent.

Proof. Choose a unit vector ordered basis $B = (v_1, \ldots, v_n)$. Let $x = \sum_{i=1}^{n} x_i v_i$. Pick a norm, say, 1-norm

$$\left\| \sum_{i=1}^{n} x_i v_i \right\|_1 = \sum_{i=1}^{n} |x_1|$$

We show that every norm for V is equivalent to $\|\cdot\|_1$, i.e., there exist $C, D > 0$ such that

$$C\|x\|_1 \geq \|x\| \geq D\|x\|_1$$

First inequality:

By triangle inequality and positive homogeneity,

$$\begin{aligned}\left\|\sum_{i=1}^{n} x_i v_i\right\| &\leq \sum_{i=1}^{n} |x_i| \left\|v_i\right\| \\ &\leq \sum_{i=1}^{n} \left\|v_i\right\| \sum_{i=1}^{n} |x_i| \\ &= \sum_{i=1}^{n} \left\|v_i\right\| \left\|\sum_{i=1}^{n} x_i v_i\right\|_1\end{aligned}$$

Take $C = \sum_{i=1}^{n} \left\|v_i\right\|$.

Second inequality:

Define the unit sphere $S \coloneqq \{x \in X : \|x\|_1 = 1\}$, which is closed and totally bounded, and thus compact by Heine-Borel theorem. Moreover, denote $\|\cdot\|$ restricted to S as f.

$$|f(x) - f(y)| \leq \|x - y\| \leq C \|x - y\|_1$$

Thus f is continuous with respect to $\|\cdot\|_1$. Therefore, by extreme value theorem, f attains its minimum on S. Let

$$D = \min_{x \in S} \|x\|$$

Since $0 \notin S$, $D > 0$.

If $x = 0$, then we have equality.

Suppose $x \neq 0$. Let $y = \dfrac{x}{\|x\|_1} \in S$. We have

$$\|y\| \geq D$$

. By absolute homogeneity,

$$\|x\| \geq D \|x\|_1$$

□

vii Linear map from finite-dimensional space is continuous and bounded

Let X be normed real or complex finite-dimensional vector space, Y be vector space, and $f : X \to Y$ be a linear map. Then, f is continuous, equivalently, f is bounded.

Proof. Choose an ordered unit vector basis $(e_1, \ldots, e_n)$ of X. Then, with coordinate notation $(x_1, \ldots, x_n)$ of x for any $x \in X$,

$$f(x) = \sum_{i=1}^{n} x_i f(e_i)$$

and by triangle inequality and absolute homogeneity:

$$\|f(x)\| \leq \sum_{i=1}^{n} \left|x_i\right| \left\|f(e_i)\right\|$$

Let

$$M = \sup_i \left\|f(e_i)\right\|$$

By all norms of X are equivalent, there exists $C > 0$ such that

$$\sum_{i=1}^{n} \left|x_i\right| \leq C \|x\|$$

Thus

$$\|f(x)\| \leq \sum_{i=1}^{n} \left|x_i\right| M \leq CM \|x\|$$

□

V Net (網)

i Net (網)

A net in X, denoted as $x_\bullet = \left(x_a\right)_{a \in A}$ or $\langle x_a \rangle_{a \in A}$, is a function of the form $x_\bullet : A \to X$ whose domain A is some directed set, and whose values are $x_\bullet(a) = x_a$ in a topological space X. Elements of a net's domain are called its indices.

ii Eventually or residually

A net $x_\bullet = \left(x_a\right)_{a \in A}$ in X is said to be eventually or residually in a set S if there exists some $a \in A$ such that for every $b \in A$ with $b \geq a$, the point $x_b \in S$.

iii Limit of nets

A point x in a topological space X is called a limit point or limit of a net $x_\bullet$ in X if, for every open neighborhood U of x, the net $x_\bullet$ is eventually in U, also called that the net converges to/towards x, and denoted as $x_\bullet \to x$, $x_a \to x$, $\lim x_\bullet \to x$, $\lim\limits_{a \in A} x_a \to x$, or $\lim\limits_{a} x_a \to x$.

If $\lim x_\bullet \to x$ and this limit is unique (i.e. $\lim x_\bullet \to y$ only if $x = y$), then one writes $\lim x_\bullet = x$, $\lim\limits_{a \in A} x_a = x$, or $\lim\limits_{a} x_a = x$.

iv Cofinal (共尾) or frequent subset

A subset B of a directed set A is a cofinal or frequent subset if for all $a \in A$, there exists $b \in B$ such that $b \geq a$.

v Cofinally (共尾地) or frequently

A net $x_\bullet = \left(x_a\right)_{a\in A}$ in X is said to be cofinally or frequently in a set S if for every $a \in A$ there exists $b \in A$ such that $b \geq a \wedge x_b \in S$.

vi Cluster point, or accumulation point (集積點)

A point x in a topological space X is called a cluster point or accumulation point of a net $x_\bullet$ in X if, for every open neighborhood U of x, the net $x_\bullet$ is cofinally in U, also called that the net clusters to/towards x.

A limit point of a net is necessary a cluster point of it.

vii Subnet or Willard-subnet

A net $s_\bullet = \left(s_i\right)_{i\in I}$ is called a subnet or Willard-subnet of another net $x_\bullet = \left(x_a\right)_{a\in A}$ if there exists an order-preserving map $h : I \to A$ such that $h(I)$ is a cofinal subset of A and

$$s_i = x_{h(i)} \quad \forall i \in I.$$

viii Ultranet or universal net

A net $x_\bullet$ in X is called an ultranet or a universal net if for every subset $S \subseteq X$, $x_\bullet$ is eventually in S or $X \setminus S$.

A point is a limit point of a ultranet if and only if it a cluster point of it.

ix In Hausdorff spaces

A net in a Hausdorff space has at most one limit; if it has one, the limit point of it is the only cluster point of it.

Proof. TODO □

x Willard-subnet limit and cluster point theorems

For any nonempty net $\langle a_n\rangle_{n\in A}$ and its any Willard-subnet $\langle b_n\rangle_{n\in B}$, if x is a limit point of a_n, then x is a limit point of b_n.

Proof. Let $h : B \to A$ be order-preserving and $h(B)$ be a cofinal subset of A. Let U be an open neighborhood of x. There exists $a \in A$ such that $n \geq a \implies a_n \in U$. Then, for any $b \in B$ such that $h(b) \geq a$, $m \geq b \implies b_m = a_{h(m)} \in U$. □

For any nonempty net $\langle a_n\rangle_{n\in A}$ and its any Willard-subnet $\langle b_n\rangle_{n\in B}$, if x is a cluster point of b_n, then x is a cluster point of a_n.

Proof. Let $h: B \to A$ be order-preserving and $h(B)$ be a cofinal subset of A. Let U be an open neighborhood of x. For any $b \in B$, there exists $m \geq b$ such that $a_{h(m)} = b_m \in U$. Also, for any $c \in A$, there exists $d \in B$ such that $h(d) \geq c$. Thus, for any $a \in A$, there exists $n \geq a$ such that $a_n \in U$. □

Its special case when $A = B = \mathbb{N}_0$ or $\mathbb{N}_1$, and $a_n \in \mathbb{R}$ is called the subsequence convergence theorem.

VI Manifold

i Euclidean neighborhood

An Euclidean neighborhood of a point in a topological space is a neighborhood of it that is homeomorphic to an open subset of $\mathbb{R}^n$ for some $n \in \mathbb{N}_0$. Such Euclidean neighborhood is called n-dimensional.

A Euclidean neighborhood homeomorphic to an open ball in $\mathbb{R}^n$ is called a Euclidean ball.

ii Locally Euclidean space

A topological space X is called locally Euclidean iff it can be covered by n-dimensional Euclidean neighborhoods for some $n \in \mathbb{N}_0$. Such locally Euclidean space is called n-dimensional or an n-locally Euclidean space.

Euclidean balls form a basis for the topology of a locally Euclidean space.

iii (Topological) manifold (流形)

A (topological) manifold is a locally Euclidean space that is a Hausdorff space. An n-dimensional (topological) manifold or n-manifold is an n-dimensional locally Euclidean space that is a Hausdorff space.

An 1D manifold is called curve. A 2D manifold is a called a surface.

iv (Coordinate) chart

For any Euclidean neighborhood U, a homeomorphism $\phi: U \to \phi(U) \subseteq \mathbb{R}^n$ is called a (coordinate) chart on U.

v Atlas

For a locally Euclidean space M, an atlas $\{(U_i, \phi_i) \mid i \in I\}$ on M is a family of ordered pairs of Euclidean neighborhood U_i and coordinate chart ϕ_i on U_i such that $\{U_i \mid i \in I\}$ covers M.

vi Transition function or transition map

For an atlas $\{(U_i, \phi_i) \mid i \in I\}$ on an n-dimensional locally Euclidean space M, the transition function $\phi_i \phi_j^{-1}$ for $i, j \in I$ is defined as the function

$$\phi_i \phi_j^{-1} : \phi_i(U_i \cap U_j) \to \phi_j(U_i \cap U_j);\ x \mapsto \phi_i(\phi_j^{-1}(x)).$$

Such a map is a homeomorphism between open subsets of $\mathbb{R}^n$.

vii Differentiable and smooth

An atlas $\{(U_i, \phi_i) \mid i \in I\}$ is called a differentiable atlas iff for all $i, j \in I$, the transition map $\phi_i \phi_j^{-1}$ is differentiable.

An atlas $\{(U_i, \phi_i) \mid i \in I\}$ is called a (n-times) continuously differentiable atlas iff for all $i, j \in I$, the transition map $\phi_i \phi_j^{-1}$ is (n-times) continuously differentiable.

An atlas $\{(U_i, \phi_i) \mid i \in I\}$ is called a C^k atlas iff for all $i, j \in I$, the transition map $\phi_i \phi_j^{-1}$ is a C^k function.

An atlas $\{(U_i, \phi_i) \mid i \in I\}$ is called a smooth (C^∞) atlas iff for all $i, j \in I$, the transition map $\phi_i \phi_j^{-1}$ is a smooth (C^∞) function.

A differentiable manifold is a manifold such that there exists a differentiable atlas on it.

A (n-times) continuously differentiable manifold is a manifold such that there exists a (n-times) continuously differentiable atlas on it.

A C^k manifold is a manifold such that there exists a C^k atlas on it.

A smooth (C^∞) manifold is a manifold such that there exists a smooth (C^∞) atlas on it.

Let M be an n-dimensional differentiable manifold with differentiable atlas $\{(U_i, \phi_i) \mid i \in I\}$. A map $f : M \to M$ is a differentiable map iff for any charts (U_i, ϕ_i) such that $p \in U_i \wedge i \in I$ and (U_j, ϕ_j) such that $f(p) \in U_j \wedge j \in I$,

$$\phi_j \circ f \circ \phi_i^{-1} : \phi_i(U_i \cap f^{-1}(U_j)) \subseteq \mathbb{R}^n \to \mathbb{R}^n$$

is differentiable.

Let M be an n-dimensional (n-times) continuously differentiable manifold with (n-times) continuously differentiable atlas $\{(U_i, \phi_i) \mid i \in I\}$. A map $f : M \to M$ is an (n-times) continuously differentiable map iff for any charts (U_i, ϕ_i) such that $p \in U_i \wedge i \in I$ and (U_j, ϕ_j) such that $f(p) \in U_j \wedge j \in I$,

$$\phi_j \circ f \circ \phi_i^{-1} : \phi_i(U_i \cap f^{-1}(U_j)) \subseteq \mathbb{R}^n \to \mathbb{R}^n$$

is (n-times) continuously differentiable.

Let M be an n-dimensional C^k manifold with C^k atlas $\{(U_i, \phi_i) \mid i \in I\}$. A map $f : M \to M$ is a C^k map iff for any charts (U_i, ϕ_i) such that $p \in U_i \wedge i \in I$ and (U_j, ϕ_j) such that $f(p) \in U_j \wedge j \in I$,

$$\phi_j \circ f \circ \phi_i^{-1} : \phi_i(U_i \cap f^{-1}(U_j)) \subseteq \mathbb{R}^n \to \mathbb{R}^n$$

is a C^k function.

Let M be an n-dimensional smooth (C^∞) manifold with smooth (C^∞) atlas $\{(U_i, \phi_i) \mid i \in I\}$. A map $f : M \to M$ is a smooth (C^∞) map iff for any charts (U_i, ϕ_i) such that $p \in U_i \wedge i \in I$ and (U_j, ϕ_j) such that $f(p) \in U_j \wedge j \in I$,

$$\phi_j \circ f \circ \phi_i^{-1} : \phi_i(U_i \cap f^{-1}(U_j)) \subseteq \mathbb{R}^n \to \mathbb{R}^n$$

is a smooth (C^∞) function.

viii Diffeomorphism

Given two differentiable manifolds X, Y, a continuously differentiable map $f : X \to Y$ is a diffeomorphism iff it is a bijection and its inverse is differentiable. If there exists a diffeomorphism between two differentiable manifolds, they are called diffeomorphic.

Given two C^k manifolds X, Y, a C^k map $f : X \to Y$ is a C^k-diffeomorphism iff it is a bijection and its inverse is C^k. If there exists a C^k-diffeomorphism between two C^k manifolds, they are called C^k-diffeomorphic.

ix Local diffeomorphism

Given two differentiable manifolds X, Y, a map $f : X \to Y$ is a local diffeomorphism iff for any point $x \in X$, there exists an open neighborhood U of X such that $f_U : U \to f(U)$ is a diffeomorphism. If there exists a local diffeomorphism between two differentiable manifolds, they are called locally diffeomorphic.

Given two C^k manifolds X, Y, a map $f : X \to Y$ is a local C^k-diffeomorphism iff for any point $x \in X$, there exists an open neighborhood U of X such that $f_U : U \to f(U)$ is a C^k-diffeomorphism. If there exists a local C^k-diffeomorphism between two differentiable manifolds, they are called locally C^k-diffeomorphic.

VII Combination with abstract algebra

i Topological group (拓樸群)

A topological group is a triple $(X, \tau, *)$ such that (X, τ) is a topological space, $(X, *)$ is a group, and the maps of $(x, y) \in X^2 \mapsto x * y$ and $x \mapsto x^{-1}$ are continuous mapa. Such topology τ is called a group topology.

ii Topological field (拓樸域)

A topological field is defined with a topological space (X, τ) and a field $(X, +, *)$ such that the maps of $a \mapsto -a$ and $a \mapsto a^{-1}$ are continuous maps.

VIII Discrete space

i Discrete topology

A discrete topology on X is a topology such that every subset of X is open.

ii Discrete (topological) space

A space equipped with discrete topology.

iii Discrete subspace

A topological subspace whose subspace topology is a discrete topology.