Trigonometric Functions

沈威宇

November 8, 2026

Contents

1 Trigonometric Functions (三角函數) 1

I Trigonometric Ratios (三角比) or Trigonometric Functions 1

i Right triangle definitions for acute angles 1

ii Unit-circle definitions and properties 1

iii Exponential definitions 2

iv Power notation 2

v Trigonometric functions of important angles 2

vi Sinusoid 3

vii Sinc function 3

II Phasor or complex amplitude 3

i Definition 3

ii Addition 4

iii Multiplication 4

iv Differentiation 4

v Integration 4

vi Phasor diagram 4

III Inverse trigonometric functions (反三角函數) 4

i Definition 4

ii atan2 函數. 5

iii Trigonometric functions of inverse trigonometric functions 5

IV Identities 5

i 三角函數基本關係 5

ii 平移關係 6
iii 奇變偶不變，正負看象限 7
iv 歐拉關係 7
v 正切萬能公式 7
vi 二倍角公式 8
vii 半角公式或平方化倍角公式 8
viii 三倍角公式 8
ix 和差角公式 8
x 平方化正切平方公式 9
xi 三角形內角正切公式 9
xii 正餘弦函數疊合 9
xiii 和差化積公式 9
xiv 積化和差公式 10
xv 連加公式 10
xvi 正餘切和等於正餘割積公式 10
xvii Inverse tangent addition formula 10
xviii 正餘弦四次方和公式 10
xix 正餘弦四次方差公式 10
xx 正餘弦六次方和公式 10
V 正弦連乘、餘切連加、餘割平方級數與餘切平方級數公式 11

1 Trigonometric Functions (三角函數)

I Trigonometric Ratios (三角比) or Trigonometric Functions

i Right triangle definitions for acute angles

Given a right triangle with an acute angle being theta:

- Sine (sin) (正弦):

$$\sin\theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

- Cosine (cos) (餘弦):

$$\cos\theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

- Tangent (tan) (正切):

$$\tan\theta = \frac{\text{hypotenuse}}{\text{adjacent}}$$

- Cotangent (cot) (餘切):

$$\cot\theta = \frac{1}{\tan\theta}$$

- Secant (sec) (正割):

$$\sec\theta = \frac{1}{\cos\theta}$$

- Cosecant (csc) (餘割):

$$\csc\theta = \frac{1}{\sin\theta}$$

ii Unit-circle definitions and properties

Let Γ be the ray obtained by rotating by an angle θ the positive half of the x-axis (counterclockwise rotation for $\theta > 0$ and clockwise rotation for $\theta < 0$). This ray intersects the unit circle with center $O = (0, 0)$ at the point $A = (x_A, y_A)$. Let the line extended to by Γ intersects $x = 1$ at $B = (1, y_B)$ and $y = 1$ at $C = (x_C, 1)$. The tangent line to the unit circle at A intersects the x- and y-axes at points $D = (x_D, 0)$ and $E = (0, y_E)$.

Function	Symbol	Definition	Domain	Range	Period	Odd or even	Reflectional symmetric about	Rotationally symmetric of order 2 about
Sine	sin	y_A	$\mathbb{R}$	$[-1, 1]$	2π	odd	$x \in \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\}$	
Cosine	cos	x_A	$\mathbb{R}$	$[-1, 1]$	2π	even	$x \in \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\}$	
Tangent	tan	y_B	$\mathbb{R} \setminus \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\}$	$\mathbb{R}$	π	odd		$(x, y) \in \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\} \times \{0\}$
Cotangent	cot	x_C	$\mathbb{R} \setminus \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\}$	$\mathbb{R}$	π	odd		$(x, y) \in \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\} \times \{0\}$
Secant	sec	x_D	$\mathbb{R} \setminus \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\}$	$(-\infty, -1] \cup [1, \infty)$	2π	even	$x \in \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\}$	$(x, y) \in \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\} \times \{0\}$
Cosecant	csc	y_E	$\mathbb{R} \setminus \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\}$	$(-\infty, -1] \cup [1, \infty)$	2π	odd	$x \in \left\{x \middle\| \left(\frac{x}{\pi} - \frac{1}{2}\right) \in \mathbb{Z}\right\}$	$(x, y) \in \left\{x \middle\| \frac{x}{\pi} \in \mathbb{Z}\right\} \times \{0\}$

iii Exponential definitions

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i}$$
$$\cos x = \frac{e^{ix} + e^{-ix}}{2}$$
$$\tan x = -i\frac{e^{2ix} - 1}{e^{2ix} + 1}, \quad x \neq \frac{\pi}{2} + k\pi, k \in \mathbb{Z}$$
$$\cot x = i\frac{e^{2ix} + 1}{e^{2ix} - 1}, \quad x \neq k\pi, k \in \mathbb{Z}$$
$$\sec x = \frac{2e^{ix}}{e^{2ix} + 1}, \quad x \neq \pi + 2k\pi, k \in \mathbb{Z}$$
$$\csc x = i\frac{2e^{ix}}{e^{2ix} - 1}, \quad x \neq 2k\pi, k \in \mathbb{Z}$$

iv Power notation

$$\sin^n x := \begin{cases} (\sin x)^n, & n \geq 0 \\ \arcsin x, & n = -1 \end{cases}$$
$$\cos^n x := \begin{cases} (\cos x)^n, & n \geq 0 \\ \arccos x, & n = -1 \end{cases}$$
$$\tan^n x := \begin{cases} (\tan x)^n, & n \geq 0 \\ \arctan x, & n = -1 \end{cases}$$
$$\cot^n x := \begin{cases} (\cot x)^n, & n \geq 0 \\ \operatorname{arccot} x, & n = -1 \end{cases}$$
$$\sec^n x := \begin{cases} (\sec x)^n, & n \geq 0 \\ \operatorname{arcsec} x, & n = -1 \end{cases}$$
$$\csc^n x := \begin{cases} (\csc x)^n, & n \geq 0 \\ \operatorname{arccsc} x, & n = -1 \end{cases}$$

v Trigonometric functions of important angles

Radian	Angle	sin	cos	tan
0	0°	0	1	0
$\frac{\pi}{2}$	90°	1	0	
π	180°	0	-1	0
$\frac{3\pi}{2}$	270°	-1	0	
$\frac{\pi}{4}$	45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$\frac{3\pi}{4}$	135°	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	-1
$\frac{\pi}{6}$	30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$

$\frac{\pi}{3}$	60 °	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$\frac{2\pi}{3}$	120°	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$-\sqrt{3}$
$\frac{5\pi}{6}$	150°	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{3}$
$\frac{\pi}{12}$	15°	$\frac{\sqrt{6}-\sqrt{2}}{4}$	$\frac{\sqrt{6}+\sqrt{2}}{4}$	$2-\sqrt{3}$
$\frac{5\pi}{12}$	75°	$\frac{\sqrt{6}+\sqrt{2}}{4}$	$\frac{\sqrt{6}-\sqrt{2}}{4}$	$2+\sqrt{3}$
$\frac{\pi}{10}$	18°	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{\sqrt{10+2\sqrt{5}}}$
$\frac{2\pi}{10}$	36°	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{\sqrt{5}+1}$
$\frac{3\pi}{10}$	54°	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{\sqrt{10-2\sqrt{5}}}$
$\frac{4\pi}{10}$	72°	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{\sqrt{5}-1}$
	37°	≈ 0.6018	≈ 0.7986	≈ 0.7536
	53°	≈ 0.7986	≈ 0.6018	≈ 1.3270

vi Sinusoid

A function $f(t) = A\cos(\omega t + \theta)$ for some real non-negative constants A, ω, θ is called a sinusoid.

vii Sinc function

The unnormalized sinc function is defined for $x \neq 0$ as

$$\operatorname{sinc}(x) = \frac{\sin(x)}{x}.$$

The normalized sinc function is defined for $x \neq 0$ as

$$\operatorname{sinc}(x) = \frac{\sin(\pi x)}{\pi x}.$$

II Phasor or complex amplitude

i Definition

The phasor or complex amplitude of a sinusoid $A\cos(\omega t + \theta)$ for some real non-negative constants A, ω, θ is a complex number

$$Ae^{i\theta}, \quad i = \sqrt{-1},$$

also denoted as $A\angle\theta$ or a two-dimensional vector

$$(A\cos\theta, A\sin\theta),$$

which satisfies

$$\Re(Ae^{i\theta}e^{i\omega t}) = A\cos(\omega t + \theta).$$

The $A\angle\theta$ notation is often called phasor and the $Ae^{i\theta}$ notation if often called complex number or complex amplitude.

Phasor denoting $A\cos(\omega t + \theta)$ is called rotating phasor. Phasor denoting $A\cos(\theta)$ is called non-rotating phasor.

ii Addition

Consider $X_1\cos(\omega t + \varphi_1)$ and $X_2\cos(\omega t + \varphi_2)$ and associated phasors $\underline{X}_1 = X_1\angle\varphi_1$ and $\underline{X}_2 = X_2\angle\varphi_2$.

$$X_1\cos(\omega t + \varphi_1) + X_2\cos(\omega t + \varphi_2) = X_m\cos(\omega t + \varphi),$$

where

$$X_m = \left|\underline{X}_1 + \underline{X}_2\right|$$
$$\varphi = \arg\left(\underline{X}_1 + \underline{X}_2\right)$$

iii Multiplication

$$(A\angle\theta)(B\angle\varphi) = (AB)\angle(\theta + \varphi)$$

iv Differentiation

The phasor of $\frac{\mathrm{d}}{\mathrm{d}t}(A\cos(\omega t + \theta))$ is $i\omega Ae^{i\theta}$.

v Integration

The phasor of the sinusoid part of $\int A\cos(\omega\tau + \theta)\,\mathrm{d}\tau$ is $-\frac{i}{\omega}Ae^{i\theta}$.

vi Phasor diagram

A phasor as a two-dimensional vector on a plane is called phasor diagram.

III Inverse trigonometric functions (反三角函數)

i Definition

Function	Symbols	Definition	Domain	Range
Inverse sine (反正弦)	$y = \arcsin x = \sin^{-1}(x) = \operatorname{asin}(x)$	$x = \sin y$	$[-1, 1]$	$[-\frac{\pi}{2}, \frac{\pi}{2}]$
Inverse cosine (反餘弦)	$y = \arccos x = \cos^{-1}(x) = \operatorname{acos}(x)$	$x = \cos y$	$[-1, 1]$	$[0, \pi]$

Inverse tangent (反正切)	$y = \arctan x = \tan^{-1}(x) = \operatorname{atan}(x)$	$x = \tan y$	$\mathbb{R}$	$(-\frac{\pi}{2}, \frac{\pi}{2})$
Inverse cotagent (反餘切)	$y = \operatorname{arccot} x = \cot^{-1}(x) = \operatorname{acot}(x)$	$x = \cot y$	$\mathbb{R}$	$(0, \pi)$
Inverse secant (反正割)	$y = \operatorname{arcsec} x = \sec^{-1}(x) = \operatorname{asec}(x)$	$x = \sec y$	$(-\infty, -1] \cup [1, +\infty)$	$[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$
Inverse cosecant (反餘割)	$y = \operatorname{arccsc} x = \csc^{-1}(x) = \operatorname{acsc}(x)$	$x = \csc y$	$(-\infty, -1] \cup [1, +\infty)$	$[-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}]$

ii atan2 函數


atan2 or arctan2(y, x): $\mathbb{R}^2 \setminus \{(0,0)\}$ 在 $x > 0$ 時返還 $\tan(\theta) = \frac{y}{x}$ 在 $(-\frac{\pi}{2}, \frac{\pi}{2})$ 中的解，在 $x < 0$、$y \geq 0$ 時返還 $\tan(\theta) = \frac{y}{x}$ 在 $(\frac{\pi}{2}, \pi)$ 中的解，在 $x < 0$、$y < 0$ 時返還 $\tan(\theta) = \frac{y}{x}$ 在 $(-\pi, -\frac{\pi}{2})$ 中的解，在 $x = 0$、$y \neq 0$ 時返還 $\frac{y}{|y|}\frac{\pi}{2}$，在 $x = y = 0$ 時未定義。

iii Trigonometric functions of inverse trigonometric functions

θ	$\sin\theta$	$\cos\theta$	$\tan\theta$	$\cot\theta$	$\sec\theta$	$\csc\theta$
$\arcsin(x)$	x	$\sqrt{1-x^2}$	$\frac{x}{\sqrt{1-x^2}}, \quad \lvert x\rvert < 1$	$\frac{\sqrt{1-x^2}}{x}, \quad x \neq 0$	$\frac{1}{\sqrt{1-x^2}}, \quad \lvert x\rvert < 1$	$\frac{1}{x}, \quad x \neq 0$
$\arccos(x)$	$\sqrt{1-x^2}$	x	$\frac{\sqrt{1-x^2}}{x}, \quad x \neq 0$	$\frac{x}{\sqrt{1-x^2}}, \quad \lvert x\rvert < 1$	$\frac{1}{x}, \quad x \neq 0$	$\frac{1}{\sqrt{1-x^2}}, \quad \lvert x\rvert < 1$
$\arctan(x)$	$\frac{x}{\sqrt{1+x^2}}$	$\frac{1}{\sqrt{1+x^2}}$	x	$\frac{1}{x}, \quad x \neq 0$	$\sqrt{1+x^2}$	$\frac{\sqrt{1+x^2}}{x}, \quad x \neq 0$
$\operatorname{arccot}(x)$	$\frac{1}{\sqrt{1+x^2}}$	$\frac{x}{\sqrt{1+x^2}}$	$\frac{1}{x}, \quad x \neq 0$	x	$\frac{\sqrt{1+x^2}}{x}, \quad x \neq 0$	$\sqrt{1+x^2}$
$\operatorname{arcsec}(x)$	$\frac{\sqrt{x^2-1}}{\lvert x\rvert}$	$\frac{1}{x}$	$\sqrt{x^2-1}\operatorname{sgn}(x)$	$\frac{\operatorname{sgn}(x)}{\sqrt{x^2-1}}, \quad \lvert x\rvert > 1$	x	$\frac{\lvert x\rvert}{\sqrt{x^2-1}}$
$\operatorname{arccsc}(x)$	$\frac{1}{x}$	$\frac{\sqrt{x^2-1}}{\lvert x\rvert}$	$\frac{\operatorname{sgn}(x)}{\sqrt{x^2-1}}, \quad \lvert x\rvert > 1$	$\sqrt{x^2-1}\operatorname{sgn}(x)$	$\frac{\lvert x\rvert}{\sqrt{x^2-1}}$	x

IV Identities

i 三角函數基本關係



- 名稱：左側三者為正；右側三者為餘；上面二者為弦；中間二者為切；下面二者為割。

- 餘角關係：以鉛直軸為對稱軸，位於線對稱位置者，左 (t) = 右 $\left(\frac{\pi}{2}-t\right)$。

$$\sin\theta=\cos\left(\frac{\pi}{2}-\theta\right)$$
$$\cos\theta=\sin\left(\frac{\pi}{2}-\theta\right)$$
$$\tan\theta=\cot\left(\frac{\pi}{2}-\theta\right)$$
$$\cot\theta=\tan\left(\frac{\pi}{2}-\theta\right)$$
$$\sec\theta=\csc\left(\frac{\pi}{2}-\theta\right)$$
$$\csc\theta=\sec\left(\frac{\pi}{2}-\theta\right)$$

- 倒數關係：三條通過中心點的線，其兩端者互為倒數，相乘為 1。

$$\sin\theta\csc\theta=1$$
$$\cos\theta\sec\theta=1$$
$$\tan\theta\cot\theta=1$$

- 商數關係：六邊形周上，連續三個頂點形成的連線，其兩端者相乘等於中間者。

$$\tan\theta\cos\theta=\sin\theta$$
$$\sin\theta\cot\theta=\cos\theta$$
$$\sin\theta\sec\theta=\tan\theta$$
$$\cos\theta\csc\theta=\cot\theta$$
$$\tan\theta\csc\theta=\sec\theta$$
$$\cot\theta\sec\theta=\csc\theta$$

- 平方（Pythagorean）關係：圖中有三個倒正三角形，其上方兩頂點者之平方和等於在下方頂點者之平方。

$$\sin^2\theta+\cos^2\theta=1$$
$$\tan^2\theta+1=\sec^2\theta$$
$$\cot^2\theta+1=\csc^2\theta$$

ii 平移關係

$$\cos(x)=\sin\left(x+\frac{\pi}{2}\right)=-\sin\left(x-\frac{\pi}{2}\right)$$
$$\cot(x)=-\tan\left(x+\frac{\pi}{2}\right)$$
$$\csc(x)=\sec\left(x-\frac{\pi}{2}\right)=-\sec\left(x+\frac{\pi}{2}\right)$$
$$\arccos(x)=-\arcsin(x)+\frac{\pi}{2}$$
$$\operatorname{arccot}(x)=-\arctan(x)+\frac{\pi}{2}$$
$$\operatorname{arccsc}(x)=-\operatorname{arcsec}(x)+\frac{\pi}{2}$$

iii 奇變偶不變，正負看象限

今有函數 f，已知其為 sin、cos、tan、sec、csc、cot 之一，且已知 $f(\theta)$。欲求 $f(\phi)$，其中 $\phi = \pm\theta \pm n\frac{\pi}{2}, \quad n \in \mathbb{Z}$。

- 判斷方法：奇變偶不變，正負看象限。
- 上句：奇偶指 n 之奇偶，變指倒數，即：若 n 為奇數則令 $g(\theta) = \frac{1}{f(\theta)}$，否則令 $g(\theta) = f(\theta)$，則 $|f(\phi)| = |g(\theta)|$。
- 下句：象限指假設 $[r, \theta]$ 在第一象限時，$[r, \phi]$ 之象限。令該象限中任意角度為 ω。令 $k = \frac{f(\phi)}{g(\theta)}$。則 $k = \frac{f(\omega)}{|f((\omega)|}$，即：

象限 / f	一	二	三	四
sin	+	+	-	-
cos	+	-	-	+
tan	+	-	+	-
csc	+	+	-	-
sec	+	-	-	+
cot	+	-	+	-

iv 歐拉關係

$$\cos x + i \sin x = e^{ix}$$
$$\cos x - \sin x = e^{-ix}$$

v 正切萬能公式

$$\sin\theta = \frac{2\tan\frac{\theta}{2}}{1+\tan^2\frac{\theta}{2}}$$
$$\cos\theta = \frac{1-\tan^2\frac{\theta}{2}}{1+\tan^2\frac{\theta}{2}}$$
$$\tan\theta = \frac{2\tan\frac{\theta}{2}}{1-\tan^2\frac{\theta}{2}}$$

vi　二倍角公式

$$\begin{aligned}\sin 2\theta &= 2\sin\theta\cos\theta\\ \cos 2\theta &= 1-2\sin^2\theta\\ &= 2\cos^2\theta-1\\ &= \cos^2\theta-\sin^2\theta\end{aligned}$$

vii　半角公式或平方化倍角公式

$$\begin{aligned}\sin^2\frac{\theta}{2} &= \frac{1-\cos\theta}{2}\\ \cos^2\frac{\theta}{2} &= \frac{1+\cos\theta}{2}\\ \tan^2\frac{\theta}{2} &= \frac{1-\cos\theta}{1+\cos\theta}\\ \tan\frac{\theta}{2} &= \frac{\sin\theta}{1+\cos\theta}\\ &= \frac{1-\cos\theta}{\sin\theta}\\ &= \frac{1+\sin\theta-\cos\theta}{1+\sin\theta+\cos\theta}\\ &= \csc\theta-\cot\theta\end{aligned}$$

viii　三倍角公式

$$\begin{aligned}\sin 3\theta &= 3\sin\theta-4\sin^3\theta\\ \cos 3\theta &= 4\cos^3\theta-3\cos\theta\\ \tan 3\theta &= \frac{3\tan\theta-\tan^3\theta}{1-3\tan^2\theta}\end{aligned}$$

ix　和差角公式

$$\begin{aligned}\sin(\alpha+\beta) &= \sin\alpha\cos\beta+\cos\alpha\sin\beta\\ \sin(\alpha-\beta) &= \sin\alpha\cos\beta-\cos\alpha\sin\beta\\ \cos(\alpha+\beta) &= \cos\alpha\cos\beta-\sin\alpha\sin\beta\\ \cos(\alpha-\beta) &= \cos\alpha\cos\beta+\sin\alpha\sin\beta\\ \tan(\alpha+\beta) &= \frac{\tan\alpha+\tan\beta}{1-\tan\alpha\tan\beta}\\ \tan(\alpha-\beta) &= \frac{\tan\alpha-\tan\beta}{1+\tan\alpha\tan\beta}\\ \cot(\alpha+\beta) &= \frac{\cot\alpha\cot\beta-1}{\cot\alpha+\cot\beta}\\ \cot(\alpha-\beta) &= \frac{\cot\alpha\cot\beta+1}{\cot\beta-\cot\alpha}\end{aligned}$$

$$\sec(\alpha+\beta)=\frac{\sec\alpha\sec\beta}{1-\tan\alpha\tan\beta}=\frac{\csc\alpha\csc\beta}{\cot\alpha\cot\beta-1}$$
$$\sec(\alpha-\beta)=\frac{\sec\alpha\sec\beta}{1+\tan\alpha\tan\beta}=\frac{\csc\alpha\csc\beta}{\cot\alpha\cot\beta+1}$$
$$\csc(\alpha+\beta)=\frac{\csc\alpha\csc\beta}{\cot\alpha+\cot\beta}=\frac{\sec\alpha\sec\beta}{\tan\alpha+\tan\beta}$$
$$\csc(\alpha-\beta)=\frac{\csc\alpha\csc\beta}{\cot\beta-\cot\alpha}=\frac{\sec\alpha\sec\beta}{\tan\alpha-\tan\beta}$$

x　平方化正切平方公式

$$\sin^2\theta=\frac{\tan^2\theta}{1+\tan^2\theta}$$
$$\cos^2\theta=\frac{1}{1+\tan^2\theta}$$
$$\cot^2\theta=\frac{1}{\tan^2\theta}$$
$$\sec^2\theta=1+\tan^2\theta$$
$$\csc^2\theta=\frac{1+\tan^2\theta}{\tan^2\theta}$$

xi　三角形內角正切公式

$$\tan\alpha+\tan\beta+\tan(\pi-\alpha-\beta)=\tan\alpha\cdot\tan\beta\cdot\tan(\pi-\alpha-\beta)$$

xii　正餘弦函數疊合

$$\left(a\sin\theta+b\cos\theta\right)^2\le a^2+b^2,\quad a,b\in\mathbb{R}$$
$$\begin{aligned}a\sin x+b\cos x&=\sqrt{a^2+b^2}\sin\left(x+\tan^{-1}\left(\frac{b}{a}\right)\right)\\&=\sqrt{a^2+b^2}\cos\left(x-\tan^{-1}\left(\frac{a}{b}\right)\right)\end{aligned}$$

xiii　和差化積公式

$$\sin\alpha+\sin\beta=2\sin\frac{\alpha+\beta}{2}\cos\frac{\alpha-\beta}{2}$$
$$\sin\alpha-\sin\beta=2\cos\frac{\alpha+\beta}{2}\sin\frac{\alpha-\beta}{2}$$
$$\cos\alpha+\cos\beta=2\cos\frac{\alpha+\beta}{2}\cos\frac{\alpha-\beta}{2}$$
$$\cos\alpha-\cos\beta=-2\sin\frac{\alpha+\beta}{2}\sin\frac{\alpha-\beta}{2}$$

xiv 積化和差公式

$$2\sin\alpha\cos\beta = \sin(\alpha+\beta)+\sin(\alpha-\beta)$$
$$2\cos\alpha\sin\beta = \sin(\alpha+\beta)-\sin(\alpha-\beta)$$
$$2\cos\alpha\cos\beta = \cos(\alpha+\beta)+\cos(\alpha-\beta)$$
$$2\sin\alpha\sin\beta = -\cos(\alpha+\beta)+\cos(\alpha-\beta)$$

xv 連加公式

$$\sum_{k=1}^{n}\sin(k\theta) = \frac{\sin\left(\frac{n\theta}{2}\right)\sin\left(\frac{(n+1)\theta}{2}\right)}{\sin\left(\frac{\theta}{2}\right)}$$
$$\sum_{k=1}^{n}\cos(k\theta) = \frac{\sin\left(\frac{n\theta}{2}\right)\cdot\cos\left(\frac{(n+1)\theta}{2}\right)}{\sin\left(\frac{\theta}{2}\right)}$$

xvi 正餘切和等於正餘割積公式

$$\tan\theta+\cot\theta = \sec\theta\csc\theta$$

xvii Inverse tangent addition formula

$$\arctan(u)+\arctan(v) = \arctan\left(\frac{u+v}{1-uv}\right) \mod \pi, \quad uv \neq 1$$
$$\arctan(u)-\arctan(v) = \arctan\left(\frac{u-v}{1+uv}\right) \mod \pi, \quad uv \neq -1$$

Proof. Take $\alpha = \arctan(u), \beta = \arctan(v)$ and take arctan on both sides of

$$\tan(\alpha+\beta) = \frac{\tan\alpha+\tan\beta}{1-\tan\alpha\tan\beta}$$
$$\tan(\alpha-\beta) = \frac{\tan\alpha-\tan\beta}{1+\tan\alpha\tan\beta}$$

□

xviii 正餘弦四次方和公式

$$\sin^4\theta+\cos^4\theta = 1-2\sin^2\cos^2\theta = 1-\frac{1}{2}\sin^2(2\theta)$$

xix 正餘弦四次方差公式

$$\sin^4\theta-\cos^4\theta = \sin^2\theta-\cos^2\theta = -\cos(2\theta)$$

xx 正餘弦六次方和公式

$$\sin^6\theta+\cos^6\theta = 1-3\sin^2\cos^2\theta = 1-\frac{3}{4}\sin^2(2\theta)$$

V　正弦連乘、餘切連加、餘割平方級數與餘切平方級數公式

$$\prod_{k=0}^{n-1} \sin\left(x+\frac{\pi k}{n}\right) = 2^{1-n}\sin(nx)$$

$$\sum_{k=0}^{n-1} \cot\left(x+\frac{\pi k}{n}\right) = n\cot(nx)$$

$$\sum_{k=0}^{n-1} \csc^2\left(x+\frac{\pi k}{n}\right) = n^2\csc^2(nx)$$

$$\sum_{k=1}^{n-1} \csc^2\frac{\pi k}{n} = \frac{(n-1)(n+1)}{3}$$

$$\sum_{k=1}^{n-1} \cot^2\frac{\pi k}{n} = \frac{(n-1)(n-2)}{3}$$

Proof.

$$\begin{aligned}
&\prod_{k=0}^{n-1} \sin\left(x+\frac{\pi k}{n}\right) \\
&\qquad = \prod_{k=0}^{n-1} \frac{i}{2}\left(e^{-i\left(x+\frac{\pi k}{n}\right)} - e^{i\left(x+\frac{\pi k}{n}\right)}\right) \\
&\qquad = i^n 2^{-n} \prod_{k=0}^{n-1} e^{-i\left(x+\frac{\pi k}{n}\right)} \prod_{k=0}^{n-1}(1-e^{2i\left(x+\frac{\pi k}{n}\right)}) \\
&\qquad = i^n 2^{-n} e^{-inx} e^{-i\pi\left(\frac{n-1}{2}\right)} \prod_{k=0}^{n-1}(1-e^{2i\left(x+\frac{\pi k}{n}\right)}) \\
&\qquad = i^n 2^{-n} e^{-inx} i^{1-n} \prod_{k=0}^{n-1}(1-e^{2i\left(x+\frac{\pi k}{n}\right)})
\end{aligned}$$

考慮：

$$f(t) = t^n - e^{2inx}$$

$f(t)=0$ 的根為：

$$t = e^{2i\left(x+\frac{\pi k}{n}\right)}, \quad k \in \mathbb{N}_0 \wedge k < n$$

故：

$$f(t) = \prod_{k=0}^{n-1}(t-e^{2i\left(x+\frac{\pi k}{n}\right)})$$

$$\prod_{k=0}^{n-1}(1-e^{2i\left(x+\frac{\pi k}{n}\right)}) = 1-e^{2inx}$$

代回：

$$\prod_{k=0}^{n-1} \sin\left(x+\frac{\pi k}{n}\right)$$
$$= i^n 2^{-n} e^{-inx} i^{1-n}(1-e^{2inx})$$
$$= 2^{-n} i(e^{-inx}-e^{inx})$$
$$= 2^{1-n}\sin(nx)$$

$$\sum_{k=0}^{n-1} \ln\left|\sin\left(x+\frac{\pi k}{n}\right)\right| = (1-n)\ln(2)+\ln|\sin(nx)|$$

微分兩次：

$$\sum_{k=0}^{n-1} \cot\left(x+\frac{\pi k}{n}\right) = n\cot(nx)$$

$$\sum_{k=0}^{n-1} \csc^2\left(x+\frac{\pi k}{n}\right) = n^2\csc^2(nx)$$

$$\sum_{k=1}^{n-1} \csc^2\left(x+\frac{\pi k}{n}\right) = n^2\csc^2(nx)-\csc^2(x)$$

$$\sum_{k=1}^{n-1} \csc^2\left(\frac{\pi k}{n}\right) = \lim_{x\to 0} n^2\csc^2(nx)-\csc^2(x) = \frac{(n-1)(n+1)}{3}$$

$$\cot^2(x) = \csc^2(x)-1$$

$$\sum_{k=1}^{n-1} \cot^2\frac{\pi k}{n} = \sum_{k=1}^{n-1} \csc^2\frac{\pi k}{n} - n + 1 = \frac{n^2-3n+2}{3}$$

□