

Algorithms HW

1. Let G be a group, and let $\Gamma^\bullet G$ be its descending central series. Show that:

- each $\Gamma^i G$ is a normal subgroup of G ;
- each $\Gamma^{i+1} G$ is contained in $\Gamma^i G$; and
- each $\Gamma^i G / \Gamma^{i+1} G$ is contained in the centre of $G / \Gamma^{i+1} G$.

- We show that $\Gamma^i G \trianglelefteq G$ by induction on i . Since $\Gamma^1 G = G$ our base case is $i = 1$. Recall that $\gamma^2 G$ is generated by commutators $[g, h]$ where $g, h \in G$. Now let $k \in G$ and consider

$$k[g, h]k^{-1} = kghg^{-1}h^{-1}k^{-1} = (kgk^{-1})(khk^{-1})(kg^{-1}k^{-1})(kh^{-1}k^{-1}) = [kgk^{-1}, khk^{-1}] \in \Gamma^2 G.$$

And so, $\Gamma^2 G \trianglelefteq G$ by definition.

Now suppose $\Gamma^i G \trianglelefteq G$ and consider $\Gamma^{i+1} G$, which is generated by elements $[g, h]$ where now $g \in G$ and $h \in \Gamma^i G$. Let $k \in G$ and notice that, since $\Gamma^i G$ is normal by the induction hypothesis, we have $khk^{-1} \in \Gamma^i G$. Thus the above calculation gives that $k[g, h]k^{-1} = [kgk^{-1}, khk^{-1}] \in \Gamma^{i+1} G$. Thus, each $\Gamma^i G \trianglelefteq G$ by induction.

- We show that $\Gamma^{i+1} G \subseteq \Gamma^i G$. Let $[g, h]$ be a generator of $\Gamma^{i+1} G$ so that $g \in G$ and $h \in \Gamma^i G$. Above we showed that $\Gamma^i G \trianglelefteq G$ and so $ghg^{-1} \in \Gamma^i G$. Moreover, $h \in \Gamma^i G$ implies $h^{-1} \in \Gamma^i G$ since $\Gamma^i G$ is a group. Thus

$$[g, h] = ghg^{-1}h^{-1} = (ghg^{-1})h \in \Gamma^i G.$$

Then, since each generator of $\Gamma^{i+1} G$ is contained in $\Gamma^i G$, we have $\Gamma^{i+1} G \subseteq \Gamma^i G$. ■

2

■

3

■

4. Let

$$1 \rightarrow N \rightarrow G \xrightarrow{\pi} H \rightarrow 1$$

be an extension of groups. Show that there is a homomorphism

$$\rho: H \rightarrow \text{Out}(N)$$

sending an element $h \in H$ to the outer automorphism of N given by conjugation by any $\tilde{h} \in G$ such that $\pi(\tilde{h}) = h$. In the particular case that $G = N \rtimes_{\theta} H$ is the semidirect product of H by N via θ , show that ρ is equal to the composition

$$H \xrightarrow{\theta} \text{Aut}(N) \rightarrow \text{Out}(N).$$

Firstly, we will show that ρ is a well defined map $H \rightarrow \text{Out}(N)$. Let $h \in H$ and $\tilde{h}_1, \tilde{h}_2 \in G$ such that $\pi(\tilde{h}_1) = \pi(\tilde{h}_2) = h$. We have $\rho(\tilde{h}_1) = f := (n \mapsto \tilde{h}_1 n \tilde{h}_1^{-1})$ and $\rho(\tilde{h}_2) = g := (n \mapsto \tilde{h}_2 n \tilde{h}_2^{-1})$. Note that these are indeed automorphisms of N , as in the previous homework we showed that conjugation by a fixed element is an automorphism. If we show that $\rho(\tilde{h}_1)$ and $\rho(\tilde{h}_2)$ lie in the same coset of $\text{Inn}(N)$ then ρ is well-defined. (Note: I believe this map is not well defined as a map $H \rightarrow \text{Aut}(N)$).

Recall that two elements g, h of a group lie in the same coset of a normal subgroup N if $g^{-1}h \in N$. For our automorphisms f, g we have $g^{-1} = (n \mapsto \tilde{h}_2^{-1} n \tilde{h}_2)$. And so we have $(g^{-1} \circ f)(n) = \tilde{h}_2^{-1} \tilde{h}_1 n \tilde{h}_1^{-1} \tilde{h}_2$. Recall that $N \trianglelefteq G$ and so is closed under conjugation by definition. In particular then $\tilde{h}_1 n \tilde{h}_1^{-1} \in N$ and $\tilde{h}_2^{-1}(\tilde{h}_1 n \tilde{h}_1^{-1}) \tilde{h}_2 \in N$ since $\tilde{h}_1, \tilde{h}_2 \in G$. Thus f, g have the same image in $\text{Out}(N)$ and so ρ is well defined with respect to the choice of \tilde{h} .

Next we show that ρ is a group homomorphism. Let $h_1, h_2 \in H$ and $\tilde{h}_1, \tilde{h}_2 \in G$ such that $\pi(\tilde{h}_1) = h_1$ and $\pi(\tilde{h}_2) = h_2$. Moreover, since π is a group homomorphism we have $\pi(\tilde{h}_1 \tilde{h}_2) = \tilde{h}_1 \tilde{h}_2$. Following a similar, calculation to last week's homework, consider the following

$$\begin{aligned}
\rho(h_1 h_2) &= \gamma_{\tilde{h}_1 \tilde{h}_2} \\
&= (n \mapsto \tilde{h}_1 \tilde{h}_2 n (\tilde{h}_1 \tilde{h}_2)^{-1}) \\
&= (n \mapsto \tilde{h}_1 \tilde{h}_2 n \tilde{h}_2^{-1} \tilde{h}_1^{-1}) \\
&= \gamma_{\tilde{h}_1} \circ \gamma_{\tilde{h}_2} \\
&= \rho(h_1) \rho(h_2).
\end{aligned}$$

Thus, the given ρ is indeed a group homomorphism.

Now suppose $G = N \rtimes_{\theta} H$. We can state more precisely the outer automorphism given by ρ . Let $h \in H$ and then all lifts are of the form $\tilde{h} = (m, h)$ for some $m \in N$. Then, being explicit about the details of the semidirect product, our map $\rho(h) : \iota(N) \rightarrow \iota(N)$ acts as follows

$$\begin{aligned}
\rho_h(n) &= (m, h) \cdot_{\theta} (n, e_H) \cdot_{\theta} (m, h)^{-1} \\
&= (m, h)(n, e_H)(\theta_{h^{-1}}(m^{-1}), h^{-1}) \\
&= (m\theta_h(n), h)(\theta_{h^{-1}}(m^{-1}), h^{-1}) \\
&= (m\theta_h(n)(\theta_h \circ \theta_{h^{-1}}(m^{-1}), hh^{-1}) \\
&= (m\theta_h(n)m^{-1}, e_H).
\end{aligned}$$

Which induces the automorphism $f = (n \mapsto m\theta_h(n)m^{-1}) : N \rightarrow N$. Note that $(\theta_h \theta_{h^{-1}}) = id_H$ since θ is a group homomorphism $H \rightarrow Aut(N)$.

We show that this is the same as the composition $H \rightarrow Aut(N) \rightarrow Out(N)$. We have $h \mapsto \theta_h \mapsto \overline{\theta_h}$. Notice now that θ_h and f lie in the same coset of $Inn(N)$. In particular

$$\overline{\theta_h} = \overline{\gamma_m \theta_h} = \overline{f}$$

since $\gamma_m = (n \mapsto mn m^{-1})$ is one of the inner automorphisms of N . Hence, in the case where $G = N \rtimes_{\theta} H$ we have ρ and $H \rightarrow Aut(N) \rightarrow Out(N)$ give the same map.

One interpretation of this is that, whilst ρ is a well defined map $H \rightarrow Out(N)$, it is not a well defined map $H \rightarrow Aut(N)$. However, in the case where G is a semidirect product of

N and H via θ , we have a preferred lift $h \mapsto (e_N, h) \in G$, and in fact there is a well defined map $H \rightarrow \text{Aut}(N)$, namely θ , whose projection gives the same map as ρ .

■

5. (Aluffi Exercise IV.5.15) Let G be a group of order 28.

- Prove that G contains a subgroup of order 4, and a normal subgroup of order 7. Deduce that G is either a split extension of C_4 by C_7 , or is a split extension of $C_2 \times C_2$ by C_7 .
- Prove that there are only two homomorphisms $C_4 \rightarrow \text{Aut}(C_7)$ and only two homomorphisms $C_2 \times C_2 \rightarrow \text{Aut}(C_7)$, up to changing the choice of generators for C_4 and $C_2 \times C_2$.
- Deduce that there are exactly four groups of order 28, up to isomorphism.

- Sylow's theorem I gives us that there exists a subgroup of order 7 in G , since $|H| = 7^1 \cdot 4$ and $7 \nmid 4$. Alternatively, Cauchy's theorem gives us that there exists an element $g \in G$ with $|g| = 7$, hence we have $|\langle g \rangle| \leq G$. Moreover, Sylow III gives us that there's only a single Sylow 7 group. Consider, if n_7 is the number of Sylow 7 groups in G then Sylow III gives us that $n_7 \equiv 1 \pmod{7}$ and $n_7 \mid 4$. The only integer solving both these conditions is $n_7 = 1$. Likewise if we write $|G| = 28 = 2^2 \cdot 7$ and notice $2 \nmid 7$ then Sylow I gives us that there exists a subgroup of order $2^2 = 4$.

Next we argue that N is normal. If $g \in G$ then recall $\gamma_g = (\ell \mapsto g\ell g^{-1}) \in \text{Aut}(G)$. Therefore $|\gamma_g(N)| = |N|$. However, there's a unique subgroup of order 7 in G and so the image $\gamma_g(N) = N$ for all $g \in G$. That is, N is closed under conjugation by elements in G and so N is normal by definition. We have shown that G has a normal subgroup of order 7 and in fact we have found that $N \cong C_7$.

- Recall **or perhaps I shall prove** that $\text{Aut}(N) = \text{Aut}(C_7) \cong C_6$. Consider C_4 , once we have specified where a generator $\sigma \in C_4$ is mapped to in C_6 then we have determined the homomorphism $C_4 \rightarrow C_6$. Since $|\sigma| = 4$ we must have $|\theta(\sigma)| = 4$ or $|\theta(\sigma)| = 2$, for θ non-trivial, since a homomorphism must map an element to an element whose order divides the original order. Notice that there's only a single element of order 2 in C_6 . And so there's one trivial map and one non-trivial map $\bar{\theta} : C_4 \rightarrow N$. Since $\bar{\theta}(\sigma)$ has order two we can deduce that it is the automorphism which sends each element of C_7 to its inverse. That is $\bar{\theta}(\sigma) = (n \mapsto 7 - n)$. And, of course, the trivial map $\theta_{\text{triv}}(\sigma) = (n \mapsto 0)$ for each $\sigma \in C_4$.

We use similar reasoning to determine the maps $\theta : C_2 \times C_2 \rightarrow \text{Aut}(N) \cong C_6$.

One generating set of $C_2 \times C_2$ is $\{(0, 1), (1, 0)\}$ and again, once we determine where these elements are mapped to by θ we have determined the entire homomorphism $\theta : C_2 \times C_2 \rightarrow C_6$. Now each generating element has order two, and so any non-trivial θ maps both the generating elements to the unique element of order 2 in C_6 . And so, again, we have one trivial map $\theta_{\text{triv}} : C_2 \times C_2 \rightarrow C_6$ and one non-trivial map $\tilde{\theta} : C_2 \times C_2 \rightarrow C_6$. The automorphisms $\tilde{\theta}((0, 1)) = \tilde{\theta}(1, 0)$ are both the same as the one described above — $(n \mapsto 7 - n \equiv -n)$.

- Determining all the possible semi-direct products $C_7 \rtimes H$ with $H = C_4$ or $H = C_2 \times C_2$ will tell us the possible group laws on G . Notice that $N \cap H = \{e\}$ for $H = C_4$ or $C_2 \times C_2$, this follows since every element of $N \cong C_7$ is the identity or is order 7, meanwhile there are no elements of order 7 in either C_4 or $C_2 \times C_2$. **We also need to show that $NH = G$.** Then it follows that $G \cong N \rtimes_{\theta} H$ for $H = C_4$ or $H = C_2 \times C_2$ and one of the θ described above.

With all possible homomorphisms $H \rightarrow \text{Aut}(N)$ described above, we can determine all the semi-direct products $N \rtimes H$. First suppose $H = C_4$ and $\theta : C_4 \rightarrow C_6$ the trivial map. That is $\theta(h) = (n \mapsto n)$ for each $h \in H$. We have the following group product for $N \rtimes_{\theta} H$:

$$\begin{aligned} (n_1, h_1) \cdot_{\theta} (n_2, h_2) &= (n_1 \theta_{h_1}(n_2), h_1 h_2) \\ &= (n_1 n_2, h_1 h_2). \end{aligned}$$

That is, then $N \rtimes_{\theta} H$ is isomorphic to $C_7 \times C_4 \cong G$. The same calculation will give us that when $H = C_2 \times C_2$ and $\theta : C_2 \times C_2 \rightarrow \text{Aut}(N)$ is the trivial map, we also have $G \cong C_7 \times C_2 \times C_2$.

Now we determine the products given by the non-trivial $H \rightarrow \text{Aut}(N)$.

■

4

■

4

■

4

■