

# Two electron integrals - Particle in a cube

Jellium Team

March 11, 2017

## Abstract

Analytical expressions for two-electron repulsion integrals in particle in a cube basis

## 1 Introduction

Two-electron integrals for particle in a cube:

Each integral is of the form:

$$(ab|cd) = \int \frac{\phi_a(r_1)\phi_b(r_1)\phi_c(r_2)\phi_d(r_2)}{|r_1 - r_2|} dr_1 dr_2 \quad (1)$$

where  $\phi_a(r_1) = \left(\frac{2}{\pi}\right)^{3/2} \sin(a_x x_1) \sin(a_y y_1) \sin(a_z z_1)$ .

Using the trigonometric identity  $2\sin(a_x x_1)\sin(b_x x_1) = \cos((a_x - b_x)x_1) - \cos((a_x + b_x)x_1)$  leads to the realization that the two electron integrals  $(ab|cd)$  can be expanded as linear combinations of integrals of the form

$$(p|q) = \frac{1}{\pi^6} \int \frac{\cos(p_x x_1)\cos(p_y y_1)\cos(p_z z_1)\cos(q_x x_2)\cos(q_y y_2)\cos(q_z z_2)}{|r_1 - r_2|} dr_1 dr_2. \quad (2)$$