

Sampling Plans and Initial Condition Problems For Continuous Time Duration Models

James J. Heckman
University of Chicago

Econ 312, Spring 2019

Sampling Plans and Initial Condition Problems For Continuous Time Duration Models

James J. Heckman
University of Chicago

Econ 312, Spring 2019

Sampling plans and initial condition problems: Duration Models

For interrupted spells, one of the following duration times may be observed:

- time in state up to sampling date (T_b)
- time in state after sampling date (T_a)
- total time in completed spell observed at origin of sample ($T_c = T_a + T_b$)

Duration of spells beginning after the origin date of the sample, denoted T_d , are not subject to initial condition problems. The intake rate, $k(-t_b)$, is the proportion of the population entering a spell at $-t_b$. Assume:

- A time homogenous environment, i.e. constant intake rate, $k(-t_b) = k, \forall b$
- A model without observed or unobserved explanatory variables.
- No right censoring, so $T_c = T_a + T_b$
- Underlying distribution is nondefective
- $m = \int_0^\infty xg(x)dx < \infty$

The proportion of the population experiencing a spell at $t = 0$, the origin date of the sample, is

$$\begin{aligned} P_0 &= \int_0^\infty k(-t_b)(1 - G(t_b))dt_b = k \int_0^\infty (1 - G(t_b))dt_b \\ &= k \left[t_b(1 - G(t_b)) \Big|_0^\infty - \int_0^\infty t_b d(1 - G(t_b)) \right] \\ &= k \int_0^\infty t_b g(t_b) dt_b = km \end{aligned}$$

where $1 - G(t_b)$ is the probability the spell lasts from $-t_b$ to 0 (or equivalently, from 0 to $-t_b$).

So the density of a spell of length t_b interrupted at the beginning of the sample ($t = 0$) is

$$\begin{aligned} f(t_b) &= \frac{\text{proportion surviving til } t = 0 \text{ from batch } t_b}{\text{total surviving til } t = 0} \\ &= \frac{k(-t_b)(1 - G(t_b))}{P_0} = \frac{1 - G(t_b)}{m} \neq g(t_b) \end{aligned}$$

The probability that a spell lasts until t_c given that it has lasted from $-t_b$ to 0, is

$$g(t_c|t_b) = \frac{g(t_c)}{1 - G(t_b)}$$

So the density of a spell that lasts for t_c is

$$\begin{aligned} f(t_c) &= \int_0^{t_c} g(t_c|t_b) f(t_b) dt_b \\ &= \int_0^{t_c} \frac{g(t_c)}{m} dt_b = \frac{g(t_c)t_c}{m} \end{aligned}$$

Likewise, the density of a spell that lasts until t_a is

$$\begin{aligned} f(t_a) &= \int_0^{\infty} g(t_a + t_b | t_b) f(t_b) dt_b \\ &= \int_0^{\infty} \frac{g(t_a + t_b)}{m} dt_b \\ &= \frac{1}{m} \int_{t_a}^{\infty} g(t_b) dt_b \\ &= \frac{1 - G(t_a)}{m} \end{aligned}$$

So the functional form of $f(t_b) = f(t_a)$: Consequences of stationarity.

Some useful results that follow from this model:

- ① If $g(t) = \theta e^{-t\theta}$, then $f(t_b) = \theta e^{-t_b\theta}$ and $f(t_a) = \theta e^{-t_a\theta}$.

Proof:

$$g(t) = \theta e^{-t\theta} \rightarrow m = \frac{1}{\theta},$$

$$G(t) = 1 - e^{-t\theta} \rightarrow f(t_a) = \frac{1 - G(t)}{m} = \theta e^{-t\theta}$$

$$\textcircled{1} E(T_a) = \frac{m}{2} \left(1 + \frac{\sigma^2}{m^2}\right).$$

Proof:

$$\begin{aligned} E(T_a) &= \int t_a f(t_a) dt_a = \int t_a \frac{1 - G(t_a)}{m} dt_a \\ &= \frac{1}{m} \left[\frac{1}{2} t_a^2 (1 - G(t_a)) \Big|_0^\infty - \int \frac{1}{2} t_a^2 d(1 - G(t_a)) \right] \\ &= \frac{1}{m} \int \frac{1}{2} t_a^2 g(t_a) dt_a = \frac{1}{2m} [\text{var}(t_a) + E^2(t_a)] \\ &= \frac{1}{2m} [\sigma^2 + m^2] \end{aligned}$$

- ① $E(T_b) = \frac{m}{2}(1 + \frac{\sigma^2}{m^2})$. **Proof:** See proof of Proposition 2.
- ② $E(T_c) = m(1 + \frac{\sigma^2}{m^2})$. **Proof:**

$$E(T_c) = \int \frac{t_c^2 g(t_c)}{m} dt_c = \frac{1}{m}(\text{var}(t_c) + E^2(t_c))$$

$$\rightarrow E(T_c) = 2E(T_a) = 2E(T_b), E(T_c) > m \text{ unless } \sigma^2 = 0$$

Some Additional Results:

$$h(t) = \text{hazard} : h(t) = \frac{F(t)}{1 - F(t)}.$$

- ① $h'(t) > 0 \rightarrow E(T_a) = E(T_b) < m$. **Proof:** See Barlow and Proschan.
- ② $h'(t) < 0 \rightarrow E(T_a) = E(T_b) > m$. **Proof:** See Barlow and Proschan.

Examples

Specification of the Distribution

Weibull Distribution

- Parameters: $\lambda > 0, k > 0$
- Probability Density Function (PDF):

$$\frac{\lambda}{k} \left(\frac{t}{\lambda}\right)^{k-1} \exp\left(-\left(\frac{t}{\lambda}\right)^k\right)$$

- Cumulative Density Function:


$$1 - \exp\left(-\left(\frac{t}{\lambda}\right)^k\right)$$

- Set of Parameters:


$$\begin{pmatrix} \lambda_1, k_1 = 0.5 \\ \lambda_2, k_1 = 1.0 \\ \lambda_3, k_1 = 2.0 \\ \lambda_3, k_1 = 3.0 \end{pmatrix}, \text{ respectively}$$

Basic Distribution Graphs

PDF for Weibull Distribution




CDF of Weibull Distribution




Basic Duration Graphs


Hazard Function for Weibull Distribution




Integrated Hazard Function for Weibull




Observed and Original Distribution for T_b (Example 1)




Observed and Original Distribution for T_b (Example 3)




Observed and Original Distribution for T_b (Example 4)




Observed and Original Distribution for T_c (Example 1)



Observed and Original Distribution for T_c (Example 2)



Observed and Original Distribution for T_c (Example 3)



Observed and Original Distribution for T_c (Example 4)

