Lecture 5, Theory Income, Fall 2018 Fernando Alvarez, UofC

Euler Equations and Transversality Conditions for Dynamic Problems, CRTS case

- ► This notes introduces the elements of discrete time dynamic optimization problems.
- Conditions under which Euler equations and Transversality conditions are necessary and sufficient for a path to be optimal are discussed.

► Most of this note follows RMED, "Recursive Methods in Economic Dynamics", by Stokey and Lucas with Prescott, Chapter 6.

▶ We use this material for different examples, including those of CRTS or Homogenous returns and growth models.

Alvarez (U. Chicago) EE and Trans. Fall 2018 2

Set up in discrete time

The elements of a Dynamic Programming problem are $[X, \Gamma, F, \beta]$. X is the set of states x. We typically let x be the current state and y the next period state. $\Gamma: X \to X$, is the correspondance describing the feasibility constraints. That is for each $x \in X$, $\Gamma(x)$ is the set of feasible values for the state variable next period if the current state is x, with its graph given by

$$Gr(\Gamma) \equiv \{(y,x) : x \in X, y \in \Gamma(x)\}.$$

The period return function F(x,y) is defined on $F:Gr(\Gamma)\to R$. Finally a discount factor $\beta\in(0,1)$. The sequence problem is

$$V^{*}(x_{0}) = \max_{\{x_{t+1}\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^{t} F(x_{t}, x_{t+1})$$

subject to

$$x_{t+1} \in \Gamma(x_t)$$
 for all $t \ge 0$

with x_0 given.

◆□▶◆□▶◆臣▶◆臣▶ 臣 釣9@

Alvarez (U. Chicago) EE and Trans. Fall 2018 3 / 1

$$olt = G(K, 1)$$

 $olt = G(K, 1)$

Example: Neoclassical growth model

$$V^{*}(k_{0}) = \max_{\{k_{t+1}\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^{t} U \underbrace{(f(k_{t}) - k_{t+1})}_{C_{t}})$$

subject to

$$0 \leq \textit{k}_{t+1} \leq \textit{f}\left(\textit{k}_{t}\right)$$

 k_0 given. This fits in the general notation by letting

$$F(x,y) = U(f(x) - y)$$

$$\Gamma(x) = [0, f(x)]$$

$$\text{non-negative consumption & capital}$$

$$f(k) = G(k,1) + (1-\delta)k$$

and

where $G(\cdot, \cdot)$ is a neoclassical constant return production function and δ the depreciation rate.

(ロ) (個) (重) (重) (重) のQで

4/1

Graph (17)

Alvarez (U. Chicago) EE and Trans. Fall 2018

A related notation distinguishes between controls, u_t , and states, x_t . In this notation the sequence problem is described by $[X, U, h, g, \beta]$. Where U is the set of feasible controls, h is the period return function and g is the law of motion of the state. The sequence problem is defined as:

$$V^*(x_0) = \max_{\{u_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^t h(x_t, u_t)$$

 $\begin{cases} \text{know how to} \\ \text{go back and} \\ \text{forth} \end{cases}$

subject to the law of motion of the state

$$x_{t+1} = g(x_t, u_t)$$
 and $u_t \in U$

for x_0 given.

To see that the control-state notation is equivalent to the previous one takes

$$F(x,y) = \max_{u} \{h(x,u) : u \in U, y = g(x,u)\}$$

 $\Gamma(x) = \{y : \text{ there is } u \in U, \text{ s.t. } y = g(x,u)\}$

Alvarez (U. Chicago) EE and Trans. Fall 2018 5/1

Exercise

- Consider the neoclassical growth model as describe above (in terms of F and Γ).
- ▶ Describe it in terms of functions *h* and *g*. What is the controls and which is the state? Hint: the list of variables are capital, consumption and/or investment.

Exercise

- Consider the neoclassical growth model with variable labor supply.
- ▶ Denote hours work by n and leisure buy ℓ . Let the period utility function depend on (c, ℓ) consumption and leisure, where we assume that there an endowment one of time, so that $\ell + n = 1$.
- Let the production function be, again, G(k, n) a function of capital k and labor n.
- ▶ Describe the problem in terms of functions *h* and *g*. What are the controls and which are (is) the state? Hint: the list of variables are capital, consumption, labor and investment.
- ightharpoonup Describe the problem in terms of the period return function F and the feasible correspondence Γ.

Euler Equations (EE) and Transversality conditions (TC).

Assume that $X \in R^m$, F is C^1 , $\beta \in (0,1)$. **Def.** The path $\{x_{t+1}\}_{t=0}^{\infty}$ satisfies EE if * \checkmark , ? we ver for S

$$F_y(x_t, x_{t+1}) + \beta F_x(x_{t+1}, x_{t+2}) = 0$$
 for $t \ge 0$

end-order implicit difference equation

Def. The path $\{x_{t+1}\}_{t=0}^{\infty}$ satisfies TC if

$$\lim_{t\to\infty}\beta^{t}F_{x}\left(x_{t},x_{t+1}\right)\cdot x_{t}=0.$$

Exercise. Write the Euler equations and TC for the neoclassical growth model.

General Principle: EE and TC are necessary and sufficient for the path $\{x_{t+1}\}_{t=0}^{\infty}$ to be optimal.

Alvarez (U. Chicago) EE and Trans. Fall 2018

Sufficiency of EE and TC. We now show that the EE and TC are sufficiency for optimality, if the problem is convex. Assume that F is concave in (x,y), that $F_x\left(x_t^*,x_{t+1}^*\right)\geq 0$, and $X=R_+^m$. Then if $\left\{x_{t+1}^*\right\}_{t=0}^\infty$ satisfies EE and TC, the path $\left\{x_{t+1}^*\right\}_{t=0}^\infty$ is optimal.

Proof. We use the fact that $f(x) \le f(x^0) + f'(x^0)(x - x^0)$ for all x, if f is concave.

Take an arbitrary $\{x_{t+1}\}_{t=0}^{\infty}$ with $x_0 = x_0^*$ and $x_{t+1} \ge 0$ for all t.

$$\lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} \left[F(x_{t}, x_{t+1}) - F(x_{t}^{*}, x_{t+1}^{*}) \right]$$

$$\leq \lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} \left[F_{x} \left(x_{t}^{*}, x_{t+1}^{*} \right) \left(x_{t} - x_{t}^{*} \right) + F_{y} \left(x_{t}^{*}, x_{t+1}^{*} \right) \left(x_{t+1} - x_{t+1}^{*} \right) \right]$$

where the inequality follows by concavity. Developing the summation in the right side:



Alvarez (U. Chicago) EE and Trans. Fall 2018 9 /

$$= \lim_{T \to \infty} \left\{ F_{x} \left(x_{0}^{*}, x_{1}^{*} \right) \left(x_{0} - x_{0}^{*} \right) + F_{y} \left(x_{0}^{*}, x_{1}^{*} \right) \left(x_{1} - x_{1}^{*} \right) + \right. \\
\left. + \beta \left[F_{x} \left(x_{1}^{*}, x_{2}^{*} \right) \left(x_{1} - x_{1}^{*} \right) + F_{y} \left(x_{1}^{*}, x_{2}^{*} \right) \left(x_{2} - x_{2}^{*} \right) \right] + \\
+ \dots + \\
\left. + \beta^{t} \left[F_{x} \left(x_{t}^{*}, x_{t+1}^{*} \right) \left(x_{t} - x_{t}^{*} \right) + F_{y} \left(x_{t}^{*}, x_{t+1}^{*} \right) \left(x_{t+1} - x_{t+1}^{*} \right) \right] \right. \\
\left. + \beta^{t+1} \left[F_{x} \left(x_{t+1}^{*}, x_{t+2}^{*} \right) \left(x_{t+1} - x_{t+1}^{*} \right) + F_{y} \left(x_{t+1}^{*}, x_{t+2}^{*} \right) \left(x_{t+2} - x_{t+2}^{*} \right) \right] \right. \\
+ \dots + \\
\left. \beta^{T} \left[F_{x} \left(x_{T}^{*}, x_{T+1}^{*} \right) \left(x_{T} - x_{T}^{*} \right) + F_{y} \left(x_{T}^{*}, x_{T+1}^{*} \right) \left(x_{T+1} - x_{T+1}^{*} \right) \right] \right\}$$

using $x_0 = x_0^*$

Alvarez (U. Chicago) EE and Trans. Fall 2018 10 / 1

$$= \lim_{T \to \infty} \{ [F_{y}(x_{0}^{*}, x_{1}^{*}) + \beta F_{x}(x_{1}^{*}, x_{2}^{*})] (x_{1} - x_{1}^{*}) \\ + \beta [F_{y}(x_{1}^{*}, x_{2}^{*}) + \beta F_{x}(x_{1}^{*}, x_{2}^{*})] (x_{2} - x_{2}^{*}) \\ + \dots + \\ + \beta^{t} [F_{y}(x_{t}^{*}, x_{t+1}^{*}) + \beta F_{x}(x_{t+1}^{*}, x_{t+2}^{*})] (x_{t+1} - x_{t+1}^{*}) \\ + \beta^{t+1} [F_{y}(x_{t+1}^{*}, x_{t+2}^{*}) + \beta F_{x}(x_{t+2}^{*}, x_{t+3}^{*})] (x_{t+2} - x_{t+2}^{*}) \\ + \dots + \\ \beta^{T} F_{y}(x_{T}^{*}, x_{T+1}^{*}) (x_{T+1} - x_{T+1}^{*}) \}$$

Using EE:

$$= \lim_{T \to \infty} \beta^{T} F_{y} \left(x_{T}^{*}, x_{T+1}^{*} \right) \left(x_{T+1} - x_{T+1}^{*} \right)$$

$$= -\lim_{T \to \infty} \beta^{T+1} F_{x} \left(x_{T+1}^{*}, x_{T+2}^{*} \right) \left(x_{T+1} - x_{T+1}^{*} \right)$$

Alvarez (U. Chicago)

using
$$x_{T+1} \ge 0$$
, $F_x\left(x_{T+1}^*, x_{T+2}^*\right) \ge 0$,
$$= -\lim_{T \to \infty} \beta^{T+1} F_x\left(x_{T+1}^*, x_{T+2}^*\right) x_{T+1} + \lim_{T \to \infty} \beta^{T+1} F_x\left(x_{T+1}^*, x_{T+2}^*\right) x_{T+1}^*$$

$$\le \lim_{T \to \infty} \beta^{T+1} F_x\left(x_{T+1}^*, x_{T+2}^*\right) x_{T+1}^*$$

thus, if the TC holds:

$$\lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} \left[F(x_{t}, x_{t+1}) - F(x_{t}^{*}, x_{t+1}^{*}) \right]$$

$$\leq \lim_{T \to \infty} \beta^{T} F_{x} \left(x_{T}^{*}, x_{T+1}^{*} \right) x_{T}^{*} = 0$$

which finishes the proof.

Alvarez (U. Chicago) EE and Trans. Fall 2018 12 / 1

concavity
$$\xrightarrow{\times}$$
 does not imply unique steady state

"permanent meme" \longrightarrow multiple steady state!

We need concavity!

Def. Let \bar{x} be a steady state, i.e. a solution to

$$F_{y}\left(\bar{x},\bar{x}\right)+\beta F_{x}\left(\bar{x},\bar{x}\right)=0.$$

Exercise. For what kind of problems does $x_{t+1} = \bar{x}$ for $t \ge 0$ is optimal if $x_0 = \bar{x}$?

Exercise. Find the steady state(s) for the neoclassical growth model. Assume that G, the production function, satisfies Inada conditions.

Alvarez (U. Chicago) EE and Trans. Fall 2018 1

Necessity of EE and TC. Assume that F is C^1 . We will consider adding a variation around the optimal path $\{x\}$, denoted by ε . Let

$$\mathbf{x}_{t}(\alpha, \varepsilon) = \mathbf{x}_{t} + \alpha \varepsilon_{t}$$

for $\alpha \in R$ and $\varepsilon = \{\varepsilon_t\}_{t=0}^{\infty}$ with $\varepsilon_t \in R^m$ and $\varepsilon_0 = 0$. Then

$$V^{*}(x_{0}) = v(0) \equiv \lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} F(x_{t}(0, \varepsilon), x_{t+1}(0, \varepsilon))$$

$$\geq v(\alpha) \equiv \lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} F(x_{t}(\alpha, \varepsilon), x_{t+1}(\alpha, \varepsilon))$$

for any α, ε such that $x_{t+1}(\alpha, \varepsilon) \in \Gamma(x_t(\alpha, \varepsilon))$ for all $t \ge 0$.

◆ロト ◆回 ト ◆ 差 ト ◆ 差 ・ か Q (*)

Alvarez (U. Chicago) EE and Trans. Fall 2018 14

Since $\alpha = 0$ maximizes ν , if ν is differentiable, it must be that

$$\frac{\partial v(0)}{\partial \alpha} = 0.$$

Assuming that the limits involved in the derivative (with respect to α) and in the summation (with respect to T) can be exchanged we obtained:

$$\frac{\partial v(0)}{\partial \alpha} = \lim_{T \to \infty} \sum_{t=0}^{T} \beta^{t} [F_{x}(x_{t}, x_{t+1}) \varepsilon_{t} + F_{y}(x_{t}, x_{t+1}) \varepsilon_{t+1}]$$

$$= \lim_{T \to \infty} \sum_{t=0}^{T-1} \beta^{t} [F_{y}(x_{t}, x_{t+1}) + \beta F_{x}(x_{t+1}, x_{t+2})] \varepsilon_{t+1}$$

$$+ \lim_{T \to \infty} \beta^{T} F_{y}(x_{T}, x_{T+1}) \varepsilon_{T+1}$$

Exercise: Show the second equality above, i.e. fill the intermediate steps (Hint: imitate the sufficiency case).

 4 □ ▷ 4 □ ▷ 4 □ ▷ 4 □ ▷ 4 □ ▷ 4 □ ▷ 2 □ 4 ○ ○

 Alvarez (U. Chicago)
 EE and Trans.

 Fall 2018
 15/1

Necessity of the EE. Consider the case where $\varepsilon_s = 0$ all s, except at time t+1. In this case x_{t+1} (α, ε) will be feasible if $(x_{t+1}, x_t) \in Int(Gr(\Gamma))$. Also assume that, v is differentiable and the limits can be interchanged. Then, direct computation gives

$$\frac{\partial V(0)}{\partial \alpha} = \left[F_{y}\left(x_{t}, x_{t+1}\right) + \beta F_{x}\left(x_{t+1}, x_{t+2}\right)\right] \varepsilon_{t+1} = 0,$$

so if ε_{t+1} can be anywhere in a neighborhood of 0, we get EE

$$F_{y}(x_{t}, x_{t+1}) + \beta F_{x}(x_{t+1}, x_{t+2}) = 0.$$

Alvarez (U. Chicago) EE and Trans. Fall 2018 16 /

Necessity of TC. As explained above, assuming that v is differentiable, that interchanging the limits is valid, and using the EE:

$$\frac{\partial v(0)}{\partial \alpha} = \lim_{T \to \infty} \beta^{T} F_{y}(x_{T}, x_{T+1}) \varepsilon_{T+1}$$
$$= -\lim_{T \to \infty} \beta^{T+1} F_{x}(x_{T+1}, x_{T+2}) \varepsilon_{T+1}$$

if $\varepsilon_{T+1} = -x_{T+1}$ is feasible

$$0 = \frac{\partial v(0)}{\partial \alpha} = \lim_{T \to \infty} \beta^{T} F_{x}(x_{T}, x_{T+1}) x_{T}$$

i.e. TC must hold.

◆ロト ◆回 ト ◆ 重 ト ◆ 重 ・ 夕 Q (~)

Alvarez (U. Chicago) EE and Trans. Fall 2018 17/1

Uses of EE and TC.

Notice that EE can be regarded as a second order difference equation, i.e. define $x_{t+2} = \psi(x_{t+1}, x_t)$

$$F_{V}(x_{t}, x_{t+1}) + \beta F_{X}(x_{t+1}, \psi(x_{t+1}, x_{t})) = 0$$

There is an initial condition, x_0 , and a boundary condition, namely TC.

Alvarez (U. Chicago) EE and Trans. Fall 2018 18 / 1

Exercise. Assume that F is C^2 . What condition will suffice to uniquely define ψ ?

Exercise. Write down convexity conditions on X, F, Γ so that the dynamic problem has, at most, one solution.

Shooting algorithm. This algorithm is described as follows. Given x_0 select x_1 arbitrarily. Generate a sequence $\{x\}$ using $x_{t+2} = \psi\left(x_{t+1}, x_t\right)$ for all $t \geq 2$. Compute if the limit of this sequence satisfies the TC for the arbitrary choice of x_1 . If not, try a different one.

Exercise. For what type of problems does the shooting algorithm works? Why does it work?

Exercise: Transversality

Consider a problem with

$$F(x,y) = U(w + x(1+r) - y)$$
 with $(1+r)\beta = 1$

- ▶ This is a saving problem with constant income w and interest rate r.
- Question: Is the solution of this problem unique?
- Question: How many steady states does this problem has?
- ▶ Solution: $c_t^* = w + r x_t$ and $x_{t+1}^* = x_t^* = x_0$. Interpret it.
- Check that the proposed policy satisfied EE and Transversality.
- Give an interpretation to the EE.

Alvarez (U. Chicago) EE and Trans. Fall 2018 20 / 1

Exercise: Transversality (cont)

- ▶ Consider and alternative policy $\tilde{c}_t = \tilde{c}_0 < c_0^* = c_t^*$ for all $t \ge 0$. We are keeping the same initial condition.
- Can this policy be optimal?
- ▶ Does the path satisfies EE? Interpret it.
- \triangleright Compute the implied sequence of x_t for this policy.
- Does the implied path satisfies Transversality?
- Use the result so far to give an interpretation to the Transversality condition.

 Alvarez (U. Chicago)
 EE and Trans.
 Fall 2018
 21 / 1

Takeaway: Reaching steady-state depends on:

- (1) Curvature of the utility function (2) The rate at which MPK drops

Exercise. Linear utility in the neoclassical growth model. maximize PV of dividends = invest until 1 = discounted MPK

▶ Let U(c) = c and

$$f(k) = G(k, 1) + (1 - \delta)k$$

where G is a neoclassical production function: strictly increasing and strictly concave in k, satisfying Inada conditions.

EE (doesn't need linear utility)

▶ Show that, as long as k_0 is such that $f(k_0) - \bar{k} \ge 0$ for $\frac{\beta f'(\bar{k}) = 1}{k_0}$, then capital converges to steady state \bar{k} in one period, i.e $\bar{k} = a(x_t)$ were adenotes the optimal policy.

(Hint: use the sufficiency of EE and Transversality). unique steady state

- If consumption is non-negative and $f(k_0) < \bar{k}$ what will be the optimal policy? Hint: trickier question, since you have to consider corners.
- cross-derivative is zero (like the case when we didn't have dynamic problems)
- so there's only one solution.
- polar opposite of permanent income (where you stayed where you began; now you go to the steady state right away)

Alvarez (U. Chicago) EE and Trans. Fall 2018 22 / 1

Exercise: Adjustment cost model

Let the adjustment cost model be:

$$F(x,y) = -\frac{a}{2}y^2 - \frac{b}{2}(y-x)^2$$

 $\Gamma(x) = R$

- ▶ What is the interpretation of $b/a \ge 0$.
- ▶ Suppose that $x_0 = 0$. What is the optimal path after that initial condition?
- Write the EE and evaluate them at the steady state. What is that value?
- ▶ What is the optimal policy if a = 0?
- ▶ Show that the optimal policy is $x_{t+1} = g(x_t) = \gamma x_t$ for some $0 < \gamma < 1$. Characterize γ in terms of b/a and β . You should obtain a quadratic equation for γ in terms of the parameters.
- Give an economic interpretation of the results.

Alvarez (U. Chicago) EE and Trans. Fall 2018 23/1

Exercise: constant saving rate.

► Consider the Neoclassical growth model with log utility, Cobb-Douglas production function and 100% depreciation: i.e.

$$F(x,y) = \log(x^{\alpha} - y)$$

$$\Gamma(x) = [0, x^{\alpha}]$$

Show that the optimal policy is of the form

$$k_{t+1} = g(k_t) = s x^{\alpha}$$
.

Find an expression for s in terms of α and β . Hints: Use EE and replace the optimal policy for consumption.

◆□▶ ◆■▶ ◆■▶ ◆■ ◆9<0</p>

Alvarez (U. Chicago) EE and Trans. Fall 2018 24 / 1

Exercise: constant savings rate.

Consider the Neoclassical growth model with 100% depreciation,

$$f(k) = \left[\alpha k^{1-\frac{1}{\rho}} + (1-\alpha)^{1-\frac{1}{\rho}}\right]^{1/(1-1/\rho)}$$

$$U(c) = \left(c^{1-1/\sigma} - 1\right) / (1-1/\sigma)$$

▶ Look for the relationship between parameters ρ and σ such that the optimal policy is to have a constant savings rate:

$$k_{t+1} = g(k_t) = s f(k_t)$$

for some number $s \in (0, 1)$.

▶ Hint: The previous exercise is an special case of this. In the previous case the elasticity of substitution of capital is one, and the intertemporal elasticity of substitution σ is also 1.

4 D > 4 B > 4 E > 4 E > E 900

Homogeneous of degree 1 case (CRTS)

- ▶ Assume *X* is a cone, $x \in X \implies \lambda x \in X$ for all scalar $\lambda > 0$.
- $y \in \Gamma(x) \implies \lambda y \in \Gamma(\lambda x)$ for all scalar $\lambda > 0$.
- ► $F(\lambda x, \lambda y) = \lambda F(x, y)$ for all scalar λ and $(x, y) \in Graph(\Gamma)$
- ► Result: Optimal policy homogeneous of degree one, $y = g(x) \implies \lambda y = g(\lambda x)$.
- We will specialize on one dimensional case, so $y = g(x) = \bar{g}x$ for some constant \bar{g} .

Fall 2018

26 / 1

Alvarez (U. Chicago) EE and Trans.

Homogeneous of degree 1 case (Exercise)

Using homogeneity on Euler Equation:

$$0 = F_{y}(x_{t}, x_{t+1}) + \beta F_{x}(x_{t+1}, x_{t+2})$$
$$= F_{y}\left(1, \frac{x_{t+1}}{x_{t}}\right) + \beta F_{x}\left(1, \frac{x_{t+2}}{x_{t+1}}\right)$$

• Guessing $\bar{g} = \frac{\chi_{t+1}}{\chi_t} = \frac{\chi_{t+2}}{\chi_{t+1}}$

Alvarez (U. Chicago)

$$0 = F_{y}(1, \bar{g}) + \beta F_{x}(1, \bar{g})$$

- Write Transversality using homogeneity & form of decision rule.
- We require that $\beta |\bar{q}| < 1$. Why?
- Why is it OK to guess? What result are we using?

4□ → 4周 → 4 重 → 4 重 → 9 Q P **EE** and Trans Fall 2018

27 / 1

Homogeneous of degree 1 case (Exercise)

- ▶ Suppose that *F* is strictly quasi-concave, can the Euler equation be satisfied for multiple values of \bar{a} ?
- Use concavity implies $F_{xx} < 0, F_{yy} < 0$ and jointly with homogeneity of degree 1, $F_{xy}^2 = F_{xx}F_{yy} > 0$.
- Differentiate Euler equation w.r.t. \bar{g} and use $F_{xx}F_{yy}=F_{xy}^2$:

$$F_{yy}(1,\bar{g}) + \beta F_{xy}(1,\bar{g}) = F_{yy}(1,\bar{g}) + \beta \sqrt{F_{xx}(1,\bar{g})F_{yy}(1,\bar{g})}$$
$$= |F_{yy}(1,\bar{g})| \left[-1 + \beta \sqrt{\frac{F_{xx}(1,\bar{g})}{F_{yy}(1,\bar{g})}} \right]$$

- \triangleright So, for β small enough the derivative is negative.
- It is also negative if $F_{xx}(1,\bar{g})/F_{yy}(1,\bar{g}) < 1$ for all \bar{g} since $\beta < 1$.

4 D > 4 B > 4 E > 4 E > E 900 Alvarez (U. Chicago) EE and Trans.

Fall 2018

General Homogeneity, Exercise

▶ Result extend to the case of homogeneity of degree 1 $-\gamma$:

$$\frac{F(\lambda x, \lambda y)}{1 - \gamma} = \lambda^{1 - \gamma} \frac{F(x, y)}{1 - \gamma} \text{ for all } x, y \text{ and } \lambda > 0$$

with the same assumptions on X and Γ .

- ▶ Alternatively $F(x, y) = H(x, y)^{1-\gamma}/(1-\gamma)$ for H homog. of degree one.
- ▶ In this case we also have g(x) homogeneous of degree one, i.e.: $y = g(x) \implies y\lambda = g(x\lambda)$.
- ▶ Strict concavity requires $\gamma > 0$. The case of $\gamma = 1$ is the log case.

This case is used a lot in growth theory. Simple example is Ak model: $c_t + i_t = A k_t$, standard l.o.m. for capital, and $u(c_t) = c_t^{1-\gamma}/(1-\gamma)$.

Transversality and Euler are a bit different. Left as $\underline{\textbf{exercise}}$ for the one dimensional case. Must use properties of derivatives of homogeneous of degree 1 $-\gamma$ function.

Alvarez (U. Chicago) EE and Trans. Fall 2018 29/1

General Homogeneity, solutions

- F(x, y) are homogeneous of degree $-\gamma$.
- ▶ Differentiate $\lambda^{1-\gamma}F(x,y) = F(\lambda x, \lambda y)$ with respect to x:

$$F_x(x,y) = \lambda^{\gamma} F_x(\lambda x, \lambda y)$$
 and $F_y(x,y) = \lambda^{\gamma} F_y(\lambda x, \lambda y)$

Apply to Euler Equations:

$$F_{x}\left(x_{t}, x_{t+1}\right) = \left(\frac{1}{x_{t}}\right)^{\gamma} F_{x}\left(1, \frac{x_{t+1}}{x_{t}}\right)$$

$$F_{y}\left(x_{t+1}, x_{t+2}\right) = \left(\frac{1}{x_{t+1}}\right)^{\gamma} F_{y}\left(1, \frac{x_{t+2}}{x_{t+1}}\right)$$

▶ Use $x_{t+1} = g x_t$ and $x_{t+2} = g x_{t+1}$:

$$0 = \left(\frac{1}{x_t}\right)^{\gamma} F_x(1,g) + \beta \left(\frac{1}{x_{t+1}}\right)^{\gamma} F_y(1,g)$$
$$0 = F_x(1,g) + \beta g^{-\gamma} F_y(1,g)$$

 4□ ▶ 4□ ▶ 4□ ▶ 4□ ▶ 4□ ▶ 4□ ▶ 4□ ▶ 4□ №

 Alvarez (U. Chicago)
 EE and Trans.

 Fall 2018
 30 / 1

Ak, solutions

- ▶ Use $F_y(x,y) = -U(f(x) y)$ and $F_x(x,y) = U'(f(x) y)f'(x)$
- ▶ Specialize to $x = 1, y = g, U'(c) = c^{-\gamma}$ and f'(x) = A:

$$0 = -\left(A-g
ight)^{-\gamma} + eta g^{-\gamma} \left(A-g
ight)^{-\gamma} ext{ or } 1 = g^{-\gamma} eta A$$

- ▶ Solution: $g = (\beta A)^{1/\gamma}$
- ▶ Taking logs, recall $log(1 + x) \approx x$:

$$\log g = \frac{1}{\gamma} \log (\beta A)$$

- ▶ Higher value of γ , more curvature, reduces growth given $\beta A > 1$. What is the economic intuition for this result?
- Higher value of βA, increases growth, given γ.
 What is the economic intuition for this result?



31 / 1

Adjustment cost and Investment

- Maximize discounted profit net of investment expenditures.
- ▶ Problem of a firm, or for economy with u(c) = c.
- ▶ Production function f(k).
 - Case f(k) strictly concave.
 - Case f(k) linear.
- ▶ Capital Law of motion $k_{t+1} = i_t + (1 \delta)k_t$
- ▶ Case w/additional cost of installing capital, in terms of final goods $\phi(i/k)k$ for some function ϕ .
- ▶ Problem: $\max_{\{i_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^t \left[f(k_t) i_t \phi\left(\frac{i_t}{k_t}\right) k_t \right]$ subject to law of motion capital.

 Alvarez (U. Chicago)
 EE and Trans.
 Fall 2018
 32 / 1

Concave f and NO Adj. Costs ("old news")

- ▶ Consider the case of f(k) strictly concave and satisfies Inada conditions.
- ▶ There is NO adjustment cost $\phi(\cdot) = 0$.
- Assume that investment can be positive or negative.
- ▶ Write F(x, y) for this case.
- ▶ Write F(x, y) the Euler Equation for this case.
- Show that steady state is achieve immediately.
- Can f be linear instead of strictly concave in this case?

Alvarez (U. Chicago) EE and Trans. Fall 2018 33 / 1

Linear f w/Adj. Costs.

- ▶ Let f(k) = Ak
- ▶ Use I.o.m. capital and define cost of adjustment function *a* as:

$$a\left(\frac{k_{t+1}}{k_t}\right) \equiv \phi\left(\frac{k_{t+1} - k_t(1-\delta)}{k_t}\right)$$

- ▶ Write F(x, y) in using the constant A and the function a.
- We will assume that:
 - a is positive (so any change implies cost) and strictly concave (so cost are increasing in size of change)
 - a'(1) = a(1) = 0 both marginal and per unit cost are zero if capital stays constant.
 - $a(1/\beta) < A$, i.e. large changes are costly.

4 D > 4 A > 4 E > 4 E > E 904 A

Alvarez (U. Chicago) EE and Trans. Fall 2018 34 / 1

Linear f w/Adj. Costs. (Exercise)

- 1. Compute F_x and F_y in terms of A and $a(\cdot)$. Make sure your expressions depend only on the ration y/x.
- 2. Write the Euler equation for this model. Use that the optimal policy is homogeneous of degree one (Why?) and denote $y = \bar{g}x$. Your expression should be a function of A, $a(\cdot)$, $a'(\cdot)$, β and \bar{g} .
- 3. Differentiate the Euler equation with respect to \bar{g} . What is the sign of this expression for values $g < 1/\beta$?
- 4. Plot the constant βA against $\beta a(g) + a'(g)(1 \beta g)$, with g in the horizontal axis. Indicate in your graph the value of \bar{g} , where both curves intersect. How is \bar{g} compared with $1/\beta$? How is \bar{g} compared with 1?
- 5. What happen with \bar{g} if A increases?
- 6. What happens if we replace the function a but another one, say \tilde{a} , with higher cost, i.e. $\tilde{a}(g) > a(g)$ for all $g \neq 1$, also with $\tilde{a}(1) = \tilde{a}'(1) = 0$ and $\tilde{a}(1/\beta) > A$.

Linear f w/Adj. Costs. (answers)

- 1. F(x,y) = Ax a(y/x)x, $F_x(x,y) = A a(y/x) + a'(y/x)(y/x)$ and $F_y(x,y) = -a'(y/x)$
- 2. Euler: $0 = -a'(\bar{g}) + \beta [A a(\bar{g}) + a'(\bar{g})\bar{g}].$
- 3. $-a''(g) + \beta [-a(g) + a'(g) + a''(g)g] = -a''(g)(1 \beta g).$
- 4. Differentiating $\beta a(g) + a'(g)(1-\beta g)$ we get $\beta a' + a''(1-\beta) a'\beta = a''(1-\beta g)$ so it is strictly increasing until $g = 1/\beta$. At g = 1, we have $\beta A > \beta a(1) + a'(1)(1-\beta) = 0$. At g = 1/beta we have $\beta A < \beta a(1/beta) + a'(g)(1-\beta/beta) = \beta a(1/beta)$.
- 5. If A increases, the value of \bar{g} increases.
- 6. Higher adjustment cost implies lower \bar{g} .

◆□▶ ◆□▶ ◆■▶ ◆■ ◆○○○

36 / 1

Alvarez (U. Chicago) EE and Trans. Fall 2018

First order condition, again

- ► Rewrite $a'(\bar{g}) = \beta \left[A a(\bar{g}) a'(\bar{g})\bar{g} \right]$ as $a'(\bar{g}) \left[1 \beta \bar{g} \right] = \beta \left[A a(\bar{g}) \right]$
- ▶ Can be expressed as $a'(\bar{g}) = \beta \frac{[A-a(\bar{g})]}{1-\beta\bar{g}}$
- ▶ What is the interpretation of $A a(\bar{g})$?
- ▶ What is the interpretation of $1/(1 \beta \bar{g})$?
- ▶ Redefine net growth rate $\gamma = \bar{g} 1$ and net interest $1 + r = 1/\beta$, rewrite

$$a'(1+\gamma) = \frac{A - a(1+\gamma)}{1/\beta - (1+\gamma)} = \frac{A - a(1+\gamma)}{r - \gamma}$$
 or
 $A = a(1+\gamma) + a'(1+\gamma)[r - \gamma]$

What happens with γ if r increases?

◆ロ > ◆ 個 > ◆ 差 > ◆ 差 > り へ ②

Alvarez (U. Chicago) EE and Trans. Fall 2018 37 / 1

Tobin's q

 \triangleright Since growth is constant at \bar{q} , we can write expected discounted profits of firms with k, or its total market value as:

$$V(k) = k \left[A - a(\bar{g}) \right] + \beta V(k\bar{g})$$

 \triangleright The function V is also homogenous of degree one. (Why?), so $\lambda V(k) = V(\lambda k)$ for all k and λ , thus

$$V(k) = [A - a(\bar{g})] k + \beta \bar{g} V(k)$$
 or $V(k) = V(1)k = \frac{A - a(\bar{g})}{1 - \beta \bar{g}} k$

Note that Market to Book value is V(k)/k = V(1) with:

$$q \equiv \frac{V(k)}{k} = V(1) = \frac{A - a(\bar{g})}{1 - \beta \bar{g}}$$

4 D > 4 B > 4 E > 4 E > E 900 **FE and Trans** Fall 2018

38 / 1

Tobin's q (cont.)

From above:

$$q \equiv \frac{V(k)}{k} = V(1) = \frac{A - a(\bar{g})}{1 - \beta \bar{g}}$$

Compare with Euler Equation:

$$\beta \left[A - a(\bar{g}) \right] = a'(\bar{g})(1 - \beta \bar{g}) \implies q \equiv V(1) = \frac{A - a(\bar{g})}{1 - \beta \bar{g}} = \frac{1}{\beta} a'(\bar{g})$$
$$q = (1 + r) a'(1 + \gamma)$$

Tobin's q equals derivative of adjustment cost $a'(\cdot)$, evaluated at the optimal growth rate \bar{g} .

Since in principle, Tobin's q is observable, as market capitalization divided by book value of capital, to forecast growth rate of investment \bar{g} .

39 / 1

Alvarez (U. Chicago) EE and Trans. Fall 2018