



Title

Finnbar Wilson - 22031076

April 19, 2024

Abstract

1 Introduction

NGC 3201 is a globular cluster, discovered by James Dunlop on the 28th of May 1826, located at $10^{\text{h}} 17^{\text{m}} 36.82^{\text{s}} / -46^{\circ} 24' 44.9''$ (in RA/Dec) 5.0 kpc away from the Sun [Paust et al., 2010]. NGC 3201 has a large sub-cluster of black hole in its core making it an interesting source for observing the interactions in large populations of black holes [Vital et al., 2022].

Globular clusters are among the oldest stellar populations in the universe, providing key insights into how stars and galactic structures evolve. In the Milky Way, some clusters are thought to have originated outside the galaxy due to their similar properties to satellite dwarf galaxies, whereas others are believed to have evolved within the Milky Way itself due to the observable effects of tidal forces and shocks in the inner galaxy. This allows for globular clusters to be classified by their characteristics as shown by Mackey and Van Den Bergh [2005] into three types: 'Young' halo(YH) which are thought to have been formed in external galaxies, 'Old' halo(OH) and 'Bulge/Disc'(BD) which are formed in the Milky Way. According to Mackey and Van Den Bergh [2005], their study on globular clusters classified NGC 3201 as a YH cluster based on the metallicity and redder horizontal branch stars. NGC 3201 stands out from other clusters classified by Mackey and Van Den Bergh [2005] due to its irregular radial velocity and differential reddening across its face [Kravtsov et al., 2010]. This makes it one of the few known clusters with an inhomogeneous stellar population for its size, which could affect how it has been classified.

In this report the structure of NGC 3201 will be analysed by finding the stellar populations inside the cluster and comparing them to isochrones to determine their age. These stellar populations will then allow a greater insight into the internal structure of the cluster creating a more accurate analysis of its classification. Given that NGC 3201 is an abnormal cluster this report will also test various methods of determining the classification of abnormal clusters.

2 Procedure

2.1 Calibration

Two images of NGC 3201 were taken in the V and B filters on a 1.0m diameter telescope. Five stars were found in each filter to calibrate the zero point magnitude in each image and their data can be found in Table 1 & 2. To find the calibration stars a catalog of local stars from SIMBAD [2000] was overlaid in each image and 10 stars were selected in total and their known magnitudes recorded. An aperture photometry of each star was performed and recorded as well as their error.

Table 1: Calibration stars in B filter

ID	RA	Dec	B _{instrument}	B _{simbad}
CI* NGC 3201 CWFD 3-109	10:17:23.65	-46:24:17.31	-12.566±0.004	16.216
CI* NGC 3201 CWFD 3-198	10:17:34.49	-46:25:36.15	-12.947±0.003	15.800
CI* NGC 3201 CWFD 3-224	10:17:36.86	-46:23:11.97	-15.062±0.001	13.601
NGC 3201 3401	10:17:38.12	-46:22:39.29	-14.787±0.001	14.080
NGC 3201 4319	10:17:42.55	-46:27:15.42	-14.812±0.001	13.999

ID is the stars identification searchable on the SIMBAD database, B_{instrument} is the magnitude recorded in this experiment and B_{simbad} is the known magnitude found on SIMBAD [SIMBAD, 2000]

Table 2: Calibration stars in V filter

ID	RA	Dec	V _{instrument}	V _{simbad}
2MASS J10173339-4620241	10:17:33.39	-46:20:24.16	-13.579±0.003	15.650
CI* NGC 3201 CWFD 3-296	10:17:42.26	-46:19:47.92	-13.904±0.002	15.230
CI* NGC 3201 CWFD 3-255	10:17:38.86	-46:22:56.86	-14.536±0.002	14.730
CI* NGC 3201 CWFD 3-235	10:17:37.72	-46:22:53.50	-14.410±0.002	14.910
CI* NGC 3201 CWFD 3-195	10:17:34.01	-46:23:26.20	-13.225±0.003	16.030

ID is the stars identification searchable on the SIMBAD database, V_{instrument} is the magnitude recorded in this experiment and V_{simbad} is the known magnitude found on SIMBAD [SIMBAD, 2000]

The equation to find the zero point magnitude is shown by Equation 1.

$$m_{\text{zero point}} = m_{\text{simbad}} - m_{\text{instrument}} \quad (1)$$

Where m is the magnitude (B or V). This produced a zero point magnitude of: B_{zero point} = 28.774 ± 0.005 and V_{zero point} = 29.2407 ± 0.0010. The errors associated with these values are from the error in the $m_{\text{instrument}}$ recordings as well as the error in the m_{simbad} which are not shown in Table 1 & 2 but can be found on SIMBAD [2000].

2.2 Automated detection

2.3 Manual detection

3 Analysis

4 Discussion

5 Conclusion

References

Kravtsov, V., Alcaíno, G., Marconi, G. and Alvarado, F. [2010], ‘Evidence of the inhomogeneity of the stellar population in the differentially reddened globular cluster ngc 3201’, *Astronomy and Astrophysics* **512**, L6.

URL: <https://doi.org/10.1051/0004-6361/200913749>

Mackey, A. D. and Van Den Bergh, S. [2005], ‘The properties of Galactic globular cluster subsystems’, *Monthly Notices of the Royal Astronomical Society* **360**(2), 631–645.

URL: <https://doi.org/10.1111/j.1365-2966.2005.09080.x>

Paust, N. E. Q., Reid, I. N., Piotto, G., Aparicio, A., Anderson, J., Sarajedini, A., Bedin, L. R. and et al. [2010], ‘The acs survey of galactic globular clusters. viii. effects of environment on globular cluster global mass functions’, *The Astronomical Journal* **139**(2), 476.

URL: <https://dx.doi.org/10.1088/0004-6256/139/2/476>

SIMBAD [2000], ‘Simbad astronomical database’, Website.

URL: <https://simbad.u-strasbg.fr/simbad/sim-basic?ident=m33&submit=SIMBAD+search>

Vitral, E., Kremer, K., Libralato, M., Mamon, G. A. and Bellini, A. [2022], ‘Stellar graveyards: clustering of compact objects in globular clusters NGC 3201 and NGC 6397’, *Monthly Notices of the Royal Astronomical Society* **514**(1), 806–825.

URL: <https://doi.org/10.1093/mnras/stac1337>