# **Chapter 1** Section 1 Exercises

**1.** With  $S_1 = \{ 2, 3, 5, 7 \}$ ,  $S_2 = \{ 2, 4, 5, 8, 9 \}$ , and  $U = \{ 1 : 10 \}$ , compute  $\overline{S}_1 \cup S_2$ . *Solution.* 

$$\overline{S}_1 = \{ 1, 4, 6, 8, 9, 10 \} \implies \overline{S}_1 \cup S_2 = \{ 1, 2, 4, 5, 6, 8, 9, 10 \}.$$

**2.** With  $S_1 = \{ 2, 3, 5, 7 \}$ ,  $S_2 = \{ 2, 4, 5, 8, 9 \}$ , compute  $S_1 \times S_2$  and  $S_2 \times S_1$ . **Solution.** 

$$S_{1} \times S_{2} = \{ (2,2), (2,4), (2,5), (2,8), (2,9), \\ (3,2), (3,4), (3,5), (3,8), (3,9), \\ (5,2), (5,4), (5,5), (5,8), (5,9), \\ (7,2), (7,4), (7,5), (7,8), (7,9) \}.$$

$$S_{2} \times S_{1} = \{ (2,2), (2,3), (2,5), (2,7), \\ (4,2), (4,3), (4,5), (4,7), \\ (5,2), (5,3), (5,5), (5,7), \\ (8,2), (8,3), (8,5), (8,7), \\ (9,2), (9,3), (9,5), (9,7) \}.$$

**3.** For  $S = \{ 2, 5, 6, 8 \}$  and  $T = \{ 2, 4, 6, 8 \}$ , compute  $|S \cap T| + |S \cup T|$ . *Solution.* 

$$S \cap T = \{\ 2,6,8\ \}, \quad S \cup T = \{\ 2,4,5,6,8\ \} \quad \Rightarrow \quad |S \cap T| + |S \cup T| = 3 + 5 = 8.$$

**4.** What relation between two sets S and T must hold so that  $|S \cup T| = |S| + |T|$ . **Solution.** 

$$|S \cup T| = |S| + |T| - |S \cap T| = |S| + |T| \quad \Rightarrow \quad |S \cap T| = 0 \quad \Rightarrow \quad S \cap T = \emptyset.$$

Therefore, S and T are disjoint.

**5.** Show that for all sets S and T,  $S - T = S \cap \overline{T}$ .

Proof.

$$S-T=\{\ x:x\in S\ \text{and}\ x\notin T\ \}$$
 
$$\iff S-T=\{\ x:x\in S\ \text{and}\ x\in \overline{T}\ \}$$
 
$$\iff S-T=S\cap \overline{T}.$$

6. Prove DeMorgan's laws,

$$\overline{S_1 \cup S_2} = \overline{S_1} \cap \overline{S_2},$$

$$\overline{S_1 \cap S_2} = \overline{S_1} \cup \overline{S_2}.$$

by showing that if an element x is in the set on one side of the equality, then it must also be in the set on the other side of the equality.

## Proof.

$$S_1 \cup S_2 = \{ x : x \in S_1 \text{ or } x \in S_2 \} \iff \overline{S_1 \cup S_2} = \{ x : x \notin S_1 \text{ and } x \notin S_2 \}.$$

$$\overline{S_1} = \{ x : x \notin S_1 \}, \quad \overline{S_2} = \{ x : x \notin S_2 \} \iff \overline{S_1} \cap \overline{S_2} = \{ x : x \notin S_1 \text{ and } x \notin S_2 \}.$$

Therefore,

$$\overline{S_1 \cup S_2} = \overline{S_1} \cap \overline{S_2}.$$

$$S_1 \cap S_2 = \{ \ x : x \in S_1 \text{ and } x \in S_2 \ \} \qquad \Longleftrightarrow \qquad \overline{S_1 \cap S_2} = \{ \ x : x \notin S_1 \text{ or } x \notin S_2 \ \}.$$

$$\overline{S_1} = \{ \ x : x \notin S_1 \ \}, \quad \overline{S_2} = \{ \ x : x \notin S_2 \ \} \qquad \Longleftrightarrow \qquad \overline{S_1} \cup \overline{S_2} = \{ \ x : x \notin S_1 \text{ or } x \notin S_2 \ \}.$$

Therefore,

$$\overline{S_1 \cap S_2} = \overline{S_1} \cup \overline{S_2}.$$

7. Show that if  $S_1 \subseteq S_2$ , then  $\overline{S_2} \subseteq \overline{S_1}$ . **Proof.** 

$$S_{1} \subseteq S_{2}$$

$$\iff x \in S_{1} \Rightarrow x \in S_{2}$$

$$\iff x \notin S_{2} \Rightarrow x \notin S_{1}$$

$$\iff x \in \overline{S_{2}} \Rightarrow x \notin \overline{S_{1}}$$

$$\iff \overline{S_{2}} \subseteq \overline{S_{1}}.$$

**8.** Show that  $S_1 = S_2$  if and only if  $S_1 \cup S_2 = S_1 \cap S_2$ . *Proof.* 

1. 
$$S_1 = S_2 \implies S_1 \cup S_2 = S_1 \cap S_2$$
.  
 $S_1 = S_2 \implies S_1 \cup S_2 = S_1 \cup S_1 = S_1$   
 $S_1 = S_2 \implies S_1 \cap S_2 = S_1 \cap S_1 = S_1$   $\Rightarrow S_1 \cup S_2 = S_1 \cap S_2$ .

2.  $S_1 \cup S_2 = S_1 \cap S_2 \implies S_1 = S_2$ .

Assume that  $S_1 \cup S_2 = S_1 \cap S_2$  and  $S_1 \neq S_2$ ,

•  $\exists x \in S_1 \text{ and } x \notin S_2 \quad \Rightarrow \quad x \in S_1 \cup S_2 \text{ and } x \notin S_1 \cap S_2 \quad \Rightarrow \quad S_1 \cup S_2 \neq S_1 \cap S_2.$ 

•  $\exists x \in S_2 \text{ and } x \notin S_1 \quad \Rightarrow \quad x \in S_1 \cup S_2 \text{ and } x \notin S_1 \cap S_2 \quad \Rightarrow \quad S_1 \cup S_2 \neq S_1 \cap S_2.$ 

The result contradicts with the permise. Therefore,  $S_1 \cup S_2 = S_1 \cap S_2 \implies S_1 = S_2$ . To sum up,  $S_1 = S_2$  if and only if  $S_1 \cup S_2 = S_1 \cap S_2$ .

**9.** Use induction on the size of S to show that if S is a finite set, then  $|2^S| = 2^{|S|}$ . **Proof.** 

#### 1. Basis

If |S| = 1, assume that  $S = \{a\}$ . Then

$$2^S = \{ \varnothing, \{ a \} \}.$$

Therefore,  $|2^S| = 2^{|S|} = 2$ .

## 2. Inductive Assumption

Assume that  $|2^{S}| = 2^{|S|}$ , for  $|S| = 1, 2, \dots, n$ .

## 3. Inductive Step

For |S| = n+1, assume that  $S = \{a_1, a_2, \dots, a_n, a_{n+1}\}$ . Let  $T = \{a_1, a_2, \dots, a_n\}$ , then

$$2^T = \{ T_1, T_2, \cdots, T_{2^n} \}.$$

For  $\forall i = 1, 2, \dots, 2^n$  where  $i \in \mathbb{N}^*$ 

$$T_i \subseteq T \\ T \subseteq S$$
  $\iff$   $T_i \subseteq S$ .

However,

$$S-T=\{\ a_{n+1}\ \} \quad \Rightarrow \quad a_{n+1}\notin T \quad \Rightarrow \quad a_{n+1}\notin T_i.$$

In addition

$$T_i \subseteq S$$

$$a_{n+1} \in S_i \quad \Longleftrightarrow \quad \{ a_{n+1} \} \subseteq S$$

$$\iff \quad T_i \cup \{ a_{n+1} \} \subseteq S.$$

Let

$$T'_{i} = T_{i} \cup \{ a_{n+1} \}, \qquad U = \{ T'_{1}, T'_{2}, \cdots, T'_{2^{n}} \}.$$

Now, for  $\forall S_i \subseteq S$ 

- If  $a_{n+1} \notin S_i$ , then  $S_i \subseteq T$ , so  $S_i \in 2^T$ .
- If  $a_{n+1} \in S_i$ , then  $S_i \{a_{n+1}\} \subseteq T$ , so  $S_i \{a_{n+1}\} \in 2^T$ . Assume that

$$S_i - \{a_{n+1}\} = T_j \quad \Rightarrow \quad S_i = T_j \cup \{a_{n+1}\} \quad \Rightarrow \quad S_i \in U.$$

Moreover,  $2^T$  and U are disjoint. Therefore,

$$2^{S} = 2^{T} \cup U$$
,  $|2^{S}| = |2^{T}| \cup |U| = 2^{n} + 2^{n} = 2^{n+1} = 2^{|S|}$ 

To sum up, if S is a finite set, then  $|2^S| = 2^{|S|}$ .

10. Show that if  $S_1$  and  $S_2$  are finite sets with  $|S_1| = n$  and  $|S_2| = m$ , then

$$|S_1 \cup S_2| \leqslant n + m$$
.

**Proof.** Assume that

$$S_1 = \{ a_1, a_2, \dots, a_n \}, \qquad S_2 = \{ b_1, b_2, \dots, b_m \}.$$

1.  $S_1$  and  $S_2$  are disjoint. Then

$$S_1 \cup S_2 = \{ a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_m \}.$$

Therefore,

$$|S_1 \cup S_2| = n + m.$$

2.  $S_1$  and  $S_2$  are not disjoint. Assume that

$$c_1, c_2, \cdots, c_k \in S_1 \text{ and } c_1, c_2, \cdots, c_k \in S_2.$$

where  $k \leq n, \ k \leq m, \ k \in \mathbb{N}^*$ . Assume that

$$b_{i_1}=c_1,\ b_{i_2}=c_2,\ \cdots,\ b_{i_k}=c_k.$$

Now

$$S_1 \cup S_2 = \{ a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_{i_1-1}, b_{i_1+1}, \dots, b_{i_k-1}, b_{i_k+1}, \dots, b_m \}.$$

Therefore,

$$|S_1 \cup S_2| = n + m - k < n + m.$$

To sum up, if  $S_1$  and  $S_2$  are finite sets with  $|S_1| = n$  and  $|S_2| = m$ , then

$$|S_1 \cup S_2| \leq n+m$$
.

11. If  $S_1$  and  $S_2$  are finite sets, show that  $|S_1 \times S_2| = |S_1||S_2|$ . **Proof.** Assume that

$$S_1 = \{ a_1, a_2, \dots, a_n \}, \qquad S_2 = \{ b_1, b_2, \dots, b_m \}.$$

Therefore,

$$S_1 \times S_2 = \{ (a_1, b_1), (a_2, b_1), \cdots, (a_n, b_1), (a_1, b_2), (a_2, b_2), \cdots, (a_n, b_2), \vdots \\ (a_1, b_m), (a_2, b_m), \cdots, (a_n, b_m) \}.$$

Thus,

$$|S_1 \times S_2| = nm = |S_1||S_2|.$$