

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/238842971>

# Value of the Steinitz constant

Article in *Functional Analysis and Its Applications* · April 1980

DOI: 10.1007/BF01086559

CITATIONS

51

READS

789

2 authors:



Victor S. Grinberg

17 PUBLICATIONS 166 CITATIONS

SEE PROFILE



Sergey Sevastyanov

Sobolev Institute of Mathematics, Siberian Branch of the Russian Academy of Sciences

67 PUBLICATIONS 1,258 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Scheduling [View project](#)



Compact vector summation and its applications [View project](#)

We sketch the proof that  $p_\alpha(\gamma) = 0$ , where  $\gamma = \gamma_\tau \otimes I$ . It can be shown that there exists a partial isometry  $A$  in  $C_{M_1}^U$  such that  $\sigma_i^U(A) = \lambda^{it}A$  ( $t \in \mathbb{R}$ ) and  $\theta_1^U(A) = A$  (cf. the proof of [5, Theorem 4.4.2]). Then  $A \in C_M^U$ ,  $\sigma_i^U(A) = \lambda^{it}A$  ( $t \in \mathbb{R}$ ), and  $\gamma_\tau^U(A) = \lambda^{it}A$ . On the other hand,  $C_{R_\infty}^U$  contains a partial isometry  $B$  such that  $\sigma_i^U(B) = \lambda^{it}B$  (cf. [5]). But then,  $AB^* \in (C_{M \otimes R_\infty}^U)_{\theta_U}$  and  $\gamma_\tau^U(AB^*) = \lambda^{it}AB^*$ , and therefore  $p_\alpha(\gamma) = 0$  (cf. [5, Sec. 4.2]).

## LITERATURE CITED

1. A. Connes, J. Funct. Anal., **16**, 415-445 (1974).
2. H. Araki and E. J. Woods, Publ. RIMS, Kyoto Univ., Ser. A, **3**, 51-130 (1968).
3. A. Connes, J. Funct. Anal., **18**, 318-327 (1975).
4. V. Ya. Golodets, Izv. Akad. Nauk SSSR, Ser. Mat., **39**, 635-656 (1975).
5. V. Ya. Golodets, Usp. Mat. Nauk, **33**, No. 1, 43-94 (1978).
6. M. Takesaki, Acta Math., **131**, Nos. 3-4, 249-310 (1973).

## VALUE OF THE STEINITZ CONSTANT

V. S. Grinberg and S. V. Sevast'yanov

UDC 513.82

A well-known lemma of Steinitz [1] states that for every normal  $m$ -dimensional space there exists a constant  $B$  such that for every collection of vectors  $x_1, x_2, \dots, x_n$  satisfying  $x_1 + x_2 + \dots + x_n = 0$ ,  $\|x_i\| \leq 1$  ( $i = 1, 2, \dots, n$ ) there exists a permutation  $\pi$  such that for all positive integers  $k \leq n$  we have  $\left\| \sum_{i=1}^k x_{\pi(i)} \right\| \leq B$ . The smallest possible value of  $B$  is denoted by  $C$  and will be called the Steinitz constant.  $C$  depends only on  $m$  and the given norm.

It is shown in [2] that for a Euclidean space,

$$C \leq V^{(4^m - 1)/3}. \quad (1)$$

A discussion of the Steinitz lemma and its applications is given in [3-5] and [6], where estimate (1) was re-discovered.

Up to now, only upper estimates exponential with respect to  $m$  have been known for  $C$ . We show that  $C \leq m$  for every norm (not necessarily even symmetric).

**LEMMA.** Let  $K$  be a polyhedron in  $\mathbb{R}^n$  defined by a system

$$\begin{cases} f_i(x) = a_i, & i = 1, 2, \dots, p, \\ g_j(x) \leq b_j, & j = 1, 2, \dots, q, \end{cases}$$

where the  $f_i$  and  $g_j$  are linear functions. Let  $x_0$  be a vertex of  $K$  and  $A = \{j: g_j(x_0) = b_j\}$ .

Then  $|A| \geq n - p$ , where  $|A|$  is the cardinality of the set  $A$ .

**Proof.** Assume the contrary. Then the system

$$\begin{cases} f_i(x) = 0, & i = 1, 2, \dots, p, \\ g_j(x) = 0, & j \in A, \end{cases}$$

has a nontrivial solution  $x_1$ . The vectors  $x_0 - \varepsilon x_1$  and  $x_0 + \varepsilon x_1$  belong to  $K$  for sufficiently small  $\varepsilon > 0$ , which contradicts the fact that  $x_0$  is a vertex.

**THEOREM 1.** Let an arbitrary norm be given in  $\mathbb{R}^n$  and assume that  $\|x_i\| \leq 1$  ( $i = 1, 2, \dots, n$ ) and  $x_1 + \dots + x_n = x$ .

Ukrainian Scientific-Research Gas Institute. Mathematics Institute, Siberian Branch, Academy of Sciences of the USSR. Translated from Funktsional'nyi Analiz i Ego Prilozheniya, Vol. 14, No. 2, pp. 56-57, April-June, 1980. Original article submitted May 5, 1979.

Then there exists a permutation  $\pi$  such that for all positive integers  $k \leq n$

$$\left\| \sum_{i=1}^k x_{\pi(i)} - \frac{k-m}{n} x \right\| \leq m. \quad (2)$$

Proof. By induction, we construct a chain of sets

$$A_m \subset A_{m+1} \subset \dots \subset A_n = \{1, 2, \dots, n\}$$

and numbers  $\lambda_k^i$  ( $k = m, m+1, \dots, n$ ;  $i \in A_k$ ) with the properties:

$$\begin{aligned} |A_k| = k, \quad 0 \leq \lambda_k^i \leq 1, \quad \sum_{i \in A_k} \lambda_k^i = k - m, \quad \sum_{i \in A_k} \lambda_k^i x_i = \frac{k-m}{n} x. \\ k = n. \quad A_n = \{1, 2, \dots, n\}, \quad \lambda_n^i = \frac{n-m}{n}. \end{aligned}$$

Inductive Step  $k+1 \rightarrow k$ . Consider the set  $K$  of collections  $\{\mu_i, i \in A_{k+1}\}$  with the properties:

$$0 \leq \mu_i \leq 1, \quad \sum_{i \in A_{k+1}} \mu_i = k - m, \quad \sum_{i \in A_{k+1}} \mu_i x_i = \frac{k-m}{n} x. \quad (3)$$

$K$  is convex and compact in  $R^{k+1}$  and nonempty (we can take  $\mu_i = \frac{k-m}{k+1-m} \lambda_{k+1}^i$ ). Let  $\{\bar{\mu}_i, i \in A_{k+1}\}$  be a vertex of  $K$ . By the lemma  $|\{i: 0 < \bar{\mu}_i < 1\}| \leq m+1$ . Using (3), we get that  $\{i: \bar{\mu}_i = 0\} \neq \emptyset$ . Let  $\bar{\mu}_j = 0$ . We put  $A_k = A_{k+1} \setminus \{j\}$ ,  $\lambda_k^i = \bar{\mu}_i$  ( $i \in A_k$ ), which completes the construction.

We put  $\{\pi(i)\} = A_i \setminus A_{i-1}$  ( $i = m+1, \dots, n$ ), with  $\pi$  otherwise arbitrary. For  $k \leq m$ , inequality (2) to be proved is obvious. For  $k \geq m+1$ ,

$$\left\| \sum_{i=1}^k x_{\pi(i)} - \frac{k-m}{n} x \right\| = \left\| \sum_{i \in A_k} (1 - \lambda_k^i) x_i \right\| \leq \sum_{i \in A_k} (1 - \lambda_k^i) = m.$$

Remark. Symmetry of the norm has not been used anywhere.

For the case  $x = 0$  and a symmetric norm (Steinitz' lemma), a more cumbersome proof is given in [7].

The following example shows that the estimate obtained is best possible. Take the unit ball to be a regular (unsymmetric) simplex with center at zero. Put  $n = m+1$ , and let  $x_i$  be the  $i$ -th vertex of the simplex. Then  $\sum x_i = 0$  and  $\|x_{\pi(1)} + \dots + x_{\pi(n-1)}\| = m$  for every permutation  $\pi$ .

The maximal lower estimate for the Steinitz constant known to the author for the case of a symmetric norm ( $(m+1)/2$  in the space  $l_1$ ,  $C \geq (m+3)^{1/2}/2$  for a Euclidean space) is attained on the collection of vectors  $B_k = \{e_i (i = 1, \dots, m-1); a_i^k (i = 1, \dots, k); b_i^k (i = 1, \dots, k)\}$ ,  $k \geq (m-1)/2$  as  $k \rightarrow \infty$ , where the  $e_i$  are unit vectors;  $a_i^k(j) = b_i^k(j) = -1/2k$  ( $j = 1, \dots, m-1, i = 1, \dots, k$ ),  $a_i^k(m) = -b_i^k(m) = 1 - (m-1)/2k$  ( $i = 1, \dots, k$ ).

We state without proof another assertion relevant to the topic considered.

THEOREM 2. Let  $x_i \in R^m$ ,  $\|x_i\| \leq 1$  ( $i = 1, 2, \dots, n$ ),  $\sum x_i = x$ . Then  $\forall \rho (0 \leq \rho \leq 1) \exists A \subseteq \{1, 2, \dots, n\}$ , for which

$$\left\| \sum_{i \in A} x_i - \rho x \right\| < m, \quad \text{and in the case of a symmetric norm} \quad \left\| \sum_{i \in A} x_i - \rho x \right\| \leq m/2.$$

Both the estimates are exact. For the proof put  $n = m$  and let  $\{x_i\}$  be the standard basis in  $R^m$ . In the symmetric case, we define an  $l_1$ -norm and set  $\rho = 1/2$ . For the unsymmetric case, let the unit ball be the convex hull of the vectors  $\{x_1, \dots, x_n, -\varepsilon x_1, \dots, -\varepsilon x_n\}$  and  $\rho = \varepsilon$ , where  $\varepsilon$  is a small positive number.

#### LITERATURE CITED

1. E. Steinitz, J. Reine Angew. Math., **143**, 128-175 (1913); **144**, 1-40 (1914).
2. V. Bergström, Abhandl. Math. Sem. Hamb. Univ., **8**, 148-152 (1931).
3. H. Hadwiger, Math. Z., **41**, No. 5, 732-738 (1936).
4. H. Hadwiger, Math. Z., **46**, No. 1, 70-79 (1940).
5. H. Hadwiger, Math. Z., **47**, No. 4, 663-668 (1942).
6. M. I. Kadets, Usp. Mat. Nauk, **8**, No. 1, 139-143 (1953).
7. S. V. Sevast'yanov, Discrete Analysis [in Russian], No. 32, Novosibirsk (1978).