

Spacetime as Escrow Bookkeeping: A Conceptual Translation of General Relativity into the Static Entropy Escrow Vocabulary

Grant Whitmer

Windstorm Institute, Fort Ann, New York 12827, USA

grant@windstorminstitute.org • ORCID: 0009-0007-3224-755X

May 2026

Abstract

We recast four standard results of general relativity—gravitational time dilation, the Tolman temperature law, the Bekenstein–Hawking entropy, and Jacobson’s (1995) thermodynamic derivation of the Einstein equations—in the vocabulary of the static gravitational entropy escrow framework [1]. A single ratio, $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ (gravitational binding energy over local Unruh temperature), reproduces all four: it equals the Bekenstein–Hawking entropy of a Schwarzschild horizon exactly, supplies the entropy flux in Jacobson’s first law, and governs the Tolman redshift. No underlying physics is modified; the contribution is interpretive—a thermodynamic re-reading, not a new equation of motion. We are explicit about the limits: $|U_{\text{grav}}|$ takes regime-specific forms not yet unified by a single covariant construction, and T_U appears in two conventions related by the Tolman shift, so the unification is exact at the algebraic level $S = |U|/T$ but only partial as a single observable. Lattice tests [2] recover the Bisognano–Wichmann asymptote partially in 1+1D (with an unexplained 1/30 prefactor) and show a qualitatively different form in 3+1D. Deriving that prefactor from first principles is the open calculation that would turn this conceptual synthesis into a constraining proposal.

1 Introduction

1.1 What this paper is

The static gravitational entropy escrow framework [1] proposes that gravitational binding energy is entropy held “in escrow” against the local Unruh temperature: for any gravitationally bound configuration with binding energy $|U_{\text{grav}}|$ at a location where the local Unruh temperature is T_U , the escrow entropy is $S_{\text{esc}} = |U_{\text{grav}}|/T_U$. The framework paper develops the consequences of this single relation, including the recovery of Newton’s law via an entropic-force argument and a reproduction of the Bekenstein–Hawking entropy formula at horizons. Companion work [2] tests the modular-Hamiltonian interpretation of the postulate against free scalar field theory on regulated lattices in 1+1 and 3+1 dimensions; that work reports approximate recovery of the Bisognano–Wichmann linear asymptote in 1+1D at small distances from the cut, with a prefactor suppression of approximately 1/30 of the literal BW value, and non-recovery in 3+1D within the d1 range the present lattice can resolve.

What the framework paper does not do, and what the lattice paper does not do, is connect the escrow postulate to the established literature on the thermodynamic derivation of Einstein’s equations — specifically to Jacobson’s (1995) derivation [3] from $\delta Q = T \cdot dS$ at local Rindler horizons, to the Tolman–Ehrenfest relation [4] for thermal equilibrium in static gravitational potentials, and to the Faulkner–Guica–Hartman–Myers–Van Raamsdonk (FGHMV) result [5,6] that demanding the entanglement-entropy first law for vacuum-state perturbations gives linearized Einstein equations in asymptotically AdS spacetimes. These three results, plus the Bekenstein–Hawking formula [7,8], constitute a now-substantial body of work establishing that general relativity has the standing of a thermodynamic equation of state. The static escrow postulate, as stated, lives squarely inside this body of work. The connection has not previously been made explicit.

The present paper makes the connection explicit. We translate each of these four standard results into the escrow vocabulary, show that the translation in each case is mathematically empty (no equation of standard GR is modified or supplemented), and demonstrate that the escrow postulate provides a single interpretive object — S_{esc} — that lets the four results be read as four faces of one identity. We are explicit (as developed in §V.G–H) that this ‘single object’ description is partly notational: the $|U_{\text{grav}}|$ feeding into S_{esc} takes regime-specific forms across the four translation legs, and T_U is used with two related but distinct conventions; the unification is real at the level of the algebraic form $S = |U|/T$, and partial at the level of identifying a single covariant observable. The contribution is a conceptual re-reading, not new physics. We are explicit throughout that ‘escrow’ is interpretive vocabulary, not a claim about a literal account-keeping mechanism in the vacuum. $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ is a defined ratio; the bookkeeping language (held in escrow, released, flow) is suggestive shorthand for the thermodynamic structure of that ratio, not a claim about ontologically separate accounts or ledgers. The framework’s scientific content is the dimensional and structural relation $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ and its consequences, not the linguistic frame in which we describe it.

1.2 Why a translation paper is worth writing

Three observations motivate the present paper. First, the escrow framework’s published presentation [1] does not state its relationship to Jacobson’s derivation explicitly, with the consequence that readers familiar with [3] have no easy path to placing the escrow postulate within the existing thermodynamic-gravity literature. The framework is at risk of being read as a parallel proposal competing with [3] when it is, more accurately, a specification of what the S in Jacobson’s $\delta Q = T \cdot dS$ is the entropy of. The present paper makes this specification explicit and addresses the parallel-proposal misreading.

Second, the lattice paper [2] tests the modular-Hamiltonian leg of this connection numerically and finds partial recovery in 1+1D and non-recovery in 3+1D. Without an explicit statement of what is being tested, readers of [2] may be unclear which interpretation of the escrow postulate the lattice results actually constrain. The present paper supplies the explicit interpretation that [2]’s tests target.

Third, the framework’s history includes one documented failed extension — the C8 entropy-current attempt described in [9] — in which an attempt to make the postulate dynamically active reduced to a rate-equation reformulation of the saturated Bekenstein bound. The failure mode there was an overreach beyond what the framework’s interpretive

content actually supports. The present paper consolidates what the framework can and cannot claim at the interpretive level, with explicit guardrails (§V) against the failure modes documented in [9] and in the present authors' subsequent reviewer-feedback iterations on [2]. These guardrails are the principal pedagogical contribution of the paper, and they include (i) explicit rejection of a separate "escrow stress-energy tensor" added to the matter source of Einstein's equation, (ii) explicit rejection of any "vacuum computational bandwidth" reading of the framework, and (iii) explicit observer-dependence of the escrow entropy (different observers, different escrows).

1.3 Scope and reading order

Sections II, III, and IV present the three legs of the translation: time dilation as Tolman temperature shift (§II), the Jacobson derivation as an escrow-flow identity (§III), and the modular-Hamiltonian connection to vacuum entanglement structure (§IV). Each section follows a common template: (A) standard statement of a known GR result, (B) translation into escrow vocabulary, (C) why the two readings reduce to the same identity, (D) explicit scope (what the section does and does not establish), (E) explicit guardrails against the local failure modes specific to that section. Section V consolidates the guardrails from §II.E, §III.F, and §IV.E into a single statement of what the framework does not commit to, with subsections G and H explicitly naming the framework's unresolved scope-limitations (§V.G: $|U_{\text{grav}}|$ lacks a single covariant definition across regimes; §V.H: T_U is used with two related but distinct conventions related by the Tolman shift). Section VI states the conclusion and identifies the principal open theoretical problems.

Three contributions of this paper are worth flagging at the outset. First, equation (8) of §II isolates the dimensionless geometric quantity $2\pi r/\lambda_C$ — the ratio of a circumference to the test particle's reduced Compton wavelength — as the dimensionless content of the escrow entropy for a single test mass; we introduce the working notation $\mathcal{N}_{\text{esc}} \equiv 2\pi r/\lambda_C$ for this quantity in §II.B and identify it as the candidate organizing variable for the test-mass leg of the translation. This ratio is the unique dimensionless invariant constructible from the Newtonian-limit ingredients (binding energy GMm/r divided by an acceleration-based Unruh temperature), so its appearance in equation (8) is dimensionally forced rather than a discovery; what is non-trivial is that the same dimensionless quantity recurs in independently-derived corners of semiclassical gravity (saturation of the Bekenstein bound for a test mass of energy mc^2 confined to radius r , Euclidean horizon periodicity, modular flow generators), and the recurrence is the smallest piece of evidence the framework is tracking a real organizing variable rather than performing a sophisticated change of variables. Second, equations (17)–(18) of §III show that the escrow postulate's prediction for the entropy of a Schwarzschild black hole equals the Bekenstein–Hawking entropy to all displayed digits, with no fudge factor; we describe this as a structural-consistency check rather than an independent prediction (the equality follows algebraically from the choice $|U_{\text{grav}}|^{\text{Schw}} = Mc^2/2$ and the standard Hawking temperature formula), and we extend the consistency check in §III.D to Reissner–Nordström and Kerr horizons via the Smarr formula. Third, §IV's connection to the FGHMV result places the lattice paper's empirical content [2] inside the FGHMV theoretical framework: the 1/30 prefactor and the 3+1D non-recovery become specific calculational questions about how lattice-regulated free QFT approaches its continuum BW limit, rather than free-floating

empirical curiosities.

The framework's broader empirical anchors live in [1] and depend on the static escrow postulate's horizon-thermodynamic content, which is unchanged by the conceptual reframing the present paper provides. We do not revisit those anchors here; the present paper concerns the GR-to-escrow translation.

2 Time Dilation As Tolman Temperature Shift

2.1 The standard statement

A clock at radius r outside a non-rotating mass M runs slow relative to a clock at infinity by the Schwarzschild redshift factor

$$\frac{d\tau}{dt} = \sqrt{1 - \frac{2GM}{rc^2}} \quad (1)$$

where τ is proper time at r and t is coordinate time (the proper time of an observer at infinity). In the weak-field limit $GM/rc^2 \ll 1$,

$$\frac{d\tau}{dt} \approx 1 - \frac{GM}{rc^2} \quad (2)$$

This is the standard expression for gravitational time dilation.

A separate but related fact about static gravitational potentials is the Tolman relation [4]. A thermal bath in equilibrium across a static spacetime does not have a uniform local temperature; the local temperature $T_{\text{local}}(r)$ measured by a stationary observer at radius r is related to the temperature measured at infinity T_∞ by

$$T_{\text{local}}(r) \sqrt{-g_{tt}(r)} = T_\infty = \text{constant} \quad (3)$$

For the Schwarzschild metric this is

$$T_{\text{local}}(r) \sqrt{1 - \frac{2GM}{rc^2}} = T_\infty \quad (4)$$

Comparing equations (1) and (4) shows that the local temperature is boosted relative to the temperature at infinity by exactly the same factor by which the local clock runs slow:

$$\frac{T_{\text{local}}(r)}{T_\infty} = \frac{dt}{d\tau} = \frac{1}{\sqrt{1 - 2GM/rc^2}} \quad (5)$$

This is a known result. It says that gravitational time dilation and the Tolman temperature shift are the same equation, expressed in two different vocabularies. What the standard derivation does not provide is an interpretation of why these two should be the same equation. The escrow framework provides one.

2.2 The escrow translation

The static escrow postulate [1] asserts that for any gravitationally bound configuration with binding energy $|U_{\text{grav}}|$ at a location where the local Unruh temperature is T_U , the entropy held in escrow against that temperature is

$$S_{\text{esc}} = \frac{|U_{\text{grav}}|}{T_U} \quad (6)$$

For a test mass m at radius r outside mass M , the binding energy is $|U_{\text{grav}}| = GMm/r$ in the Newtonian limit, and the proper acceleration required to hold the test mass static is $a = GM/r^2$. The Unruh temperature at this acceleration is

$$T_U = \frac{\hbar a}{2\pi c k_B} = \frac{\hbar GM}{2\pi c k_B r^2} \quad (7)$$

Substituting (7) into (6):

$$S_{\text{esc}} = \frac{GMm/r}{\hbar GM/(2\pi c k_B r^2)} = \frac{2\pi c k_B r m}{\hbar} \quad (8)$$

This is a real geometric quantity: $S_{\text{esc}}/k_B = 2\pi r/\lambda_C$ where $\lambda_C = \hbar/(mc)$ is the reduced Compton wavelength of the test mass — that is, the number of Compton wavelengths that fit into the circumference $2\pi r$. The dimensionless escrow counts how many Compton circumferences of the test mass fit around a circle of radius r . This is the quantity the escrow postulate identifies as the entropy held in reserve by the gravitational binding.

The same dimensionless quantity has a separate physical meaning that is worth stating explicitly. The Bekenstein bound states that the entropy of any system of total energy E confined within a sphere of radius R is bounded above by

$$S \leq 2\pi ER/(\hbar c)$$

For a system of energy $E = mc^2$ and confinement radius $R = r$, the bound takes the value

$$2\pi(mc^2)r/(\hbar c) = 2\pi mcr/\hbar = 2\pi r/\lambda_C \quad (\text{in units of } k_B)$$

— numerically identical to the escrow entropy of a test mass of rest mass m at orbital radius r given by equation (8). We are explicit that this is in the first instance a dimensional coincidence: the Bekenstein bound is a statement about a confined system, and the escrow translation considers an orbital configuration; reading the equality as ‘the escrow entropy saturates the Bekenstein bound’ requires identifying orbital radius with confinement radius, which is an interpretive step rather than a settled physical equivalence. With that interpretive identification made, the equality has a suggestive reading: gravitational binding holds the test particle’s entropy budget at the same value that the Bekenstein bound would assign to a system of its rest energy and orbital scale. The framework treats this reading as a physical motivation, independent of the postulate’s dimensional construction, for considering S_{esc} rather than some other dimensional combination as the relevant entropy; the dimensional coincidence itself is real, and its interpretive content is what the framework proposes.

Because this dimensionless ratio is the candidate organizing variable for the test-mass leg of the translation, we introduce a symbol for it. Let

$$\mathcal{N}_{\text{esc}} \equiv 2\pi r/\lambda_C = 2\pi mcr/\hbar \quad (8')$$

denote the dimensionless escrow count of a test mass m at radius r . $\mathcal{N}_{\text{esc}}$ is the number of reduced Compton wavelengths of the test mass that fit into the circumference of a circle of radius r , and equivalently — by the saturation observation above — the Bekenstein-bound entropy of the test mass at that radius in units of k_B . Within the framework, $\mathcal{N}_{\text{esc}}$ serves as the candidate organizing variable for the test-mass thermodynamic-gravity dictionary: every other quantity in §II is expressed in terms of $\mathcal{N}_{\text{esc}}$ and one local kinematic factor (T_U or $m c^2$). The naming is provisional; the dimensionless ratio it labels has appeared implicitly in semiclassical-gravity calculations for decades (e.g., in Bekenstein-bound saturation arguments, Euclidean horizon periodicity, and modular flow generators), but its identification as the escrow count of a gravitationally bound test mass is, to our knowledge, first stated in the framework paper [1] and made explicit here. If the framework is eventually adopted, the dimensionless escrow count is the most natural quantity to receive a permanent name; we have used ' $\mathcal{N}_{\text{esc}}$ ' as a working notation rather than attaching a personal label, in deference to the community's prerogative to settle such naming through citation rather than authorial declaration.

We can now rewrite the Newtonian-limit time dilation (equation 2) in escrow language. Multiplying numerator and denominator on the right side of (2) by T_U and recognizing $|U_{\text{grav}}|/T_U = S_{\text{esc}}$:

$$\frac{d\tau}{dt} \approx 1 - \frac{GM}{rc^2} = 1 - \frac{|U_{\text{grav}}|}{mc^2} = 1 - \frac{S_{\text{esc}}T_U}{mc^2} \quad (9)$$

This is the translation. The fractional slowdown of a clock at radius r is the product of the local escrow entropy and the local Unruh temperature, divided by the rest energy of the clock.

Equation (9) is not a new equation. It is equation (2) — the Newtonian limit of the Schwarzschild redshift — with the binding energy $|U_{\text{grav}}|$ replaced by its escrow decomposition $S_{\text{esc}} \cdot T_U$. Its only content beyond equation (2) is interpretive: it identifies why the clock slows down at exactly the rate it does. The clock slows because a clock is an apparatus for converting energy into distinguishable states at a rate set by its rest mass, and the energy available for state-distinction at radius r is reduced by the amount $|U_{\text{grav}}|$ that is held in escrow against the local Unruh temperature.

Numerical verification: at the Earth's equatorial surface ($M = 5.972 \times 10^{24}$ kg, $r = 6.378 \times 10^6$ m using WGS84), the right side of equation (9) evaluates to a fractional clock slowdown of 6.953×10^{-10} , which matches the published Schwarzschild redshift exactly. The corresponding accumulated clock difference is approximately 22 milliseconds per year, consistent with measured values.

C. Why the Tolman shift and the clock slowdown are the same equation

The Tolman relation (equation 4) and the redshift relation (equation 1) are the same equation because of a basic thermodynamic identity. A thermal bath in equilibrium across a static spacetime is, by definition, a configuration in which energy flows freely between regions without net change. For this to be consistent with gravitational redshift — which causes photons emitted in a region of low gravitational potential to lose energy as they climb out — the local temperature must increase to compensate. Tolman (1930) derived equation (3) from

this requirement. Equivalently, the Tolman relation is the statement that a thermal photon's energy redshifts according to (1) and a thermal photon's temperature is its characteristic energy.

In the escrow vocabulary the identity is immediate. The local Unruh temperature $T_U(r)$ is fixed by the proper acceleration $a(r) = GM/r^2$. The escrow held against that temperature is $|U_{\text{grav}}|/T_U$. The fractional clock slowdown is $|U_{\text{grav}}|/(mc^2)$. Eliminating $|U_{\text{grav}}|$ between these two relations immediately yields

$$\frac{d\tau}{dt} = 1 - \frac{S_{\text{esc}} T_U}{mc^2} = 1 - \frac{T_U S_{\text{esc}}}{E_{\text{rest}}} \quad (10)$$

so the clock slowdown is set by the ratio of (local Unruh temperature times local escrow) to the clock's rest energy. The Tolman relation, in turn, is the statement that T_{local} scales as the inverse of the clock slowdown factor. Both statements reduce to the same algebraic identity: the local Unruh temperature, the local clock rate, and the local escrow are three faces of one quantity, namely the local proper acceleration $a(r)$.

In standard GR this identity is established through the metric: $\sqrt{(-g_{tt})}$ appears in the redshift, in the Tolman factor, and in the Unruh temperature through its relationship to surface gravity. The escrow framework makes the same identity transparent by routing all three through S_{esc} and T_U explicitly, without requiring the metric as an intermediate object.

2.3 What this section does and does not establish

This section does not derive equation (1) from the escrow postulate. Equation (1) is a standard result of general relativity and has many derivations independent of any thermodynamic interpretation. What this section establishes is a dictionary: the standard expression for gravitational time dilation can be rewritten in escrow language as equation (10), and the rewriting makes manifest the relationship between time dilation and the Tolman temperature shift that is otherwise an apparent coincidence requiring separate derivation. We note explicitly that the escrow translation in equation (9) operates at the level of the weak-field expansion ($1 - GM/rc^2$), not the exact Schwarzschild factor $\sqrt{1 - 2GM/rc^2}$. It is important to distinguish two separate statements: the Tolman–Ehrenfest relation $T_{\text{local}} \cdot \sqrt{(-g_{tt})} = T_{\infty}$, equation (3), is exact for any static spacetime (it follows from the existence of a timelike Killing vector and from Tolman–Ehrenfest photon-redshift equilibrium); the weak-field approximation enters only when one inserts the linearized Schwarzschild metric factor as $\sqrt{1 - 2GM/rc^2} \approx 1 - GM/rc^2$. The escrow translation of the Schwarzschild redshift in equation (9) is therefore weak-field, but the underlying Tolman relation is not. The exact Schwarzschild form can be obtained by inserting the full metric factor into the Tolman relation rather than by extending the escrow translation itself. The framework's structural commitment at this level is to the weak-field reading of equation (9), with the strong-field exact form available through the (exact) Tolman relation.

The escrow framework's contribution at this level is interpretive, not predictive. No measurement of time dilation distinguishes the escrow reading from the standard GR reading; the equations are identical. The framework's value, here, is a conceptual re-reading: the postulate $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ provides a single object — in the partly-notational sense developed in §V.G–H — through which time dilation, Tolman temperature, Unruh temperature, and

gravitational binding energy are all expressed, replacing four separately-stated relations with one.

This conceptual re-reading is what we will exploit in Section III, where we apply the same dictionary to the Jacobson derivation of Einstein’s equations and the Bekenstein–Hawking entropy.

2.4 A note on what the framework does not commit to

We do not invoke a “vacuum computational bandwidth” or “information processing rate” reading of equation (10). Such readings appear in some popularizations of entropic gravity, and they are not part of the escrow postulate as stated in [1]. The framework’s commitment is to the thermodynamic identity (6) and its consequences; it does not commit to a digital or computational substrate for the vacuum. The local rate of escrow production at radius r is a thermal quantity, set by $T_U(r)$, and is correctly described in the vocabulary of equilibrium thermodynamics. No additional ontological structure is required to make equation (10) hold.

Similarly, we do not introduce a separate “escrow stress-energy tensor” $T_{\mu\nu}^{\text{escrow}}$ to be added to the matter stress-energy on the right side of Einstein’s equation. The reason is structural: Einstein’s equations are an algebraic identity between the matter stress-energy tensor $T_{\mu\nu}^{\text{matter}}$ and the geometric Einstein tensor $G_{\mu\nu}$, and the Bianchi identities ($\nabla^\mu G_{\mu\nu} = 0$) require $\nabla^\mu T_{\mu\nu}^{\text{matter}} = 0$ — the matter content already saturates the structure of conserved sources. Gravitational binding energy is not a localizable tensorial quantity in general relativity; it is encoded non-locally in the geometry itself (the difference, for an isolated system, between the ADM mass at infinity and the integrated rest mass on a spatial slice), and it cannot be added to $T_{\mu\nu}^{\text{matter}}$ as an independent source. The escrow framework’s S_{esc} is a thermodynamic interpretation of a configuration’s gravitational binding content; it does not introduce a new tensor and does not augment Einstein’s equation with any new stress-energy contribution. Attempting to do so would either violate the Bianchi identities or double-count: the geometry on the left of Einstein’s equation already responds to the full gravitational configuration, and a separate escrow source would constitute a second response to the same physical content. This point will be developed further in Section III.

3 The Jacobson Translation

3.1 The standard statement

In 1995, Jacobson showed [3] that Einstein’s field equations follow from demanding that the thermodynamic relation $\delta Q = T \cdot dS$ hold across every local Rindler horizon, where T is the Unruh temperature of an observer accelerating just outside the horizon and dS is the area-law entropy of the horizon. The derivation is now standard and we recapitulate only the load-bearing steps. Consider a point p in spacetime and a small local Rindler horizon through p , defined by the future causal horizon of an accelerated observer with proper acceleration a passing through p . The Unruh temperature associated with this horizon is

$$T_U = \frac{\hbar a}{2\pi c k_B} \quad (11)$$

The entropy associated with the horizon is its area in Planck units,

$$S_{\text{horizon}} = \frac{A}{4\ell_p^2} k_B \quad (12)$$

where $\ell_p = (\hbar G/c^3)^{1/2}$ is the Planck length. A small area element of the horizon expands or contracts as matter falls into the wedge behind it, and the rate of expansion is governed by the Raychaudhuri equation applied to the null geodesics generating the horizon.

Jacobson's central insight was that demanding

$$\delta Q = T_U dS_{\text{horizon}} \quad (13)$$

hold at every such point p , for every choice of local Rindler horizon, forces the Einstein field equations on the spacetime:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (14)$$

where $T_{\mu\nu}$ is the matter stress-energy tensor whose flux δQ across the horizon enters the left side of (13). The Raychaudhuri equation supplies the geometric content; the Clausius relation $\delta Q = T \cdot dS$ supplies the thermodynamic content; the combination is Einstein's equation. This derivation is not specific to Schwarzschild or to any particular solution — it holds locally everywhere, for every observer.

Jacobson's result establishes that Einstein's equation has the standing of a thermodynamic equation of state of the vacuum, analogous to the ideal-gas law $PV = NkT$ emerging from molecular kinetics rather than being postulated fundamentally. The metric, in this reading, is what you get when the vacuum's entropy obeys the second law locally everywhere; it is not a separate object whose dynamics need to be derived from a Lagrangian.

3.2 The escrow translation

The static escrow postulate identifies the entropy held against the local Unruh temperature as the gravitational binding energy divided by that temperature: $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ (equation 6). Jacobson's derivation uses $T = T_U$ and $S = S_{\text{horizon}}$ (the horizon area entropy). The question is: what is the relationship between S_{horizon} as it appears in Jacobson's derivation and S_{esc} as it appears in the escrow postulate?

The answer is that they are the same quantity, evaluated at different scales. Jacobson's S_{horizon} is the entropy associated with a particular local Rindler horizon — a coarse-grained quantity defined at the level of the horizon's area. The escrow entropy S_{esc} is the entropy held against the same Unruh temperature by whatever mass-energy is in the wedge behind that horizon. The first law of horizon thermodynamics, applied to a small flux of mass-energy crossing the horizon, equates these two: a flux δQ of energy at temperature T_U across the horizon produces a change δS in the horizon entropy, and that δS is exactly $\delta Q/T_U$ — the change in escrow entropy associated with the flux. (The naive substitution $\delta Q/T_U = \delta|U_{\text{grav}}|/T_U$ would be off by a factor of two under $|U_{\text{grav}}| = Mc^2/2$; the escrow differential $dS_{\text{esc}} = d(|U_{\text{grav}}|/T_U)$ reproduces $\delta Q/T_U$ correctly because $T_U \propto 1/M$ makes the temperature-variation term contribute equally, whereas the one-term reading $\delta|U_{\text{grav}}|/T_U$ alone omits it.) We note that the first law relates the two entropies at the differential level ($\delta S_{\text{horizon}} = \delta S_{\text{esc}}$);

the finite identification $S_{\text{horizon}} = S_{\text{esc}}$ for a Schwarzschild black hole of mass M follows from this differential relation together with integrability along the formation path, and is confirmed directly by the algebraic calculation in §III.C (equations 17–18). The finite-level confirmation does not depend solely on the differential argument.

In escrow language, Jacobson’s identity (13), with the substitution $S_{\text{horizon}} \rightarrow S_{\text{esc}}$ justified at the differential level by the first law and confirmed at the finite level by §III.C below, reads

$$\delta Q = T_U dS_{\text{esc}} \quad (15)$$

which is the differential version of the static escrow postulate (6). The static postulate says $S_{\text{esc}} = |U_{\text{grav}}|/T_U$; the differential form says that the change in escrow entropy across a horizon equals the heat flux across that horizon divided by the local Unruh temperature. These are the same statement. Equation (15) is therefore not a new equation. It is Jacobson’s (13) with S_{horizon} replaced by S_{esc} , justified by the observation that the horizon entropy is exactly the entropy held against the horizon’s Unruh temperature — which is the escrow definition.

The substitution buys nothing mathematically. It buys an interpretation: what the horizon is doing, thermodynamically, is bookkeeping the gravitational binding energy of the matter behind it. The horizon area grows when matter falls in because the matter brought additional binding energy to the wedge, and that additional binding energy must be held in escrow against the horizon’s Unruh temperature. The escrow framework supplies the answer to the question “what is the horizon’s entropy the entropy of?” — it is the entropy held in escrow against the local Unruh temperature by the gravitationally bound mass-energy in the wedge.

3.3 Why the substitution is legal

The identification $S_{\text{horizon}} = S_{\text{esc}}$ requires that two independently defined quantities coincide. The first is the horizon area entropy $A/(4\ell_P^2) \cdot k_B$, which is a property of the horizon geometry alone. The second is the escrow entropy $|U_{\text{grav}}|/T_U$, which is a property of the matter content and the local Unruh temperature. We now show that these are equal.

Consider a Schwarzschild black hole of mass M . The Schwarzschild radius and surface gravity at the horizon are

$$r_s = 2GM/c^2, \kappa = c^4/(4GM)$$

The Hawking temperature — the Tolman-redshifted (asymptotic) Unruh temperature of an observer hovering near the horizon, finite at infinity even though the local Unruh temperature of a static observer diverges as $r \rightarrow r_s$, with

$$T_H = \lim_{r \rightarrow r_s} T_{\text{Unruh}}(r) \cdot \sqrt{-g_{tt}(r)}$$

— is

$$T_U = \frac{\hbar \kappa}{2\pi c k_B} = \frac{\hbar c^3}{8\pi G M k_B} \quad (16)$$

The self-binding energy of the configuration at the horizon scale is $|U_{\text{grav}}| = GM^2/r_s =$

$Mc^2/2$ (a direct consequence of the Schwarzschild radius definition). This value also coincides with the Schwarzschild ($Q = J = 0$) limit of the Smarr identification $T_H S_{\text{BH}} = (M - \Phi Q - 2\Omega_H J)/2$ used in §III.D below, which gives the $Mc^2/2$ choice a thermodynamic motivation — it is the gravitational sector of the Smarr decomposition, with the electrostatic and rotational sectors vanishing for an uncharged non-rotating horizon — alongside the algebraic GM^2/r_s motivation. This choice of $|U_{\text{grav}}|$ is one of the four regime-specific identifications enumerated in §V.G; it is not derived from a single covariant binding-energy construction (such a construction is identified there as an open theoretical task) but is adopted because it produces the algebraic match with S_{BH} that the present subsection demonstrates. The escrow entropy is therefore

$$S_{\text{esc}}/k_B = |U_{\text{grav}}|/(k_B T_U) = (Mc^2/2) \cdot (8\pi GM k_B)/(\hbar c^3 k_B) =$$

$$\frac{4\pi GM^2}{\hbar c} \equiv \mathcal{N}_{\text{esc}}^{\text{Schw}} \quad (17)$$

where we have introduced the horizon analog of the dimensionless escrow count — $\mathcal{N}_{\text{esc}}^{\text{Schw}} \equiv 4\pi GM^2/(\hbar c) = 4\pi(M/m_P)^2$, with $m_P = \sqrt{\hbar c/G}$ the Planck mass — in analogy with the test-mass count $\mathcal{N}_{\text{esc}} \equiv 2\pi r/\lambda_C$ of equation (8') in §II.B. The horizon count $\mathcal{N}_{\text{esc}}^{\text{Schw}}$ measures the gravitationally bound mass-energy of a Schwarzschild black hole in units of the Planck-scale escrow quantum. We are explicit that $\mathcal{N}_{\text{esc}}$ and $\mathcal{N}_{\text{esc}}^{\text{Schw}}$ are not the same physical quantity in the literal sense: the test-mass $\mathcal{N}_{\text{esc}}$ has the geometric interpretation ‘reduced Compton wavelengths fitting into the orbital circumference’ (a mode count in the sense of counting wavelengths around a closed curve), while $\mathcal{N}_{\text{esc}}^{\text{Schw}}$ is the Bekenstein–Hawking entropy in dimensionless form (scaling as M^2 rather than as the linear mr of the test-mass case). They belong to the same family of dimensionless escrow constructions — each arising from the postulate $S = |U|/T$ applied in a different regime — but the family is bound by the common postulate, not by a single geometric mode-count formula. The shared notation marks the structural relationship; the regime label ($^{\text{Schw}}$) marks the difference in physical content. The Bekenstein–Hawking entropy of the same horizon is

$$\frac{S_{\text{BH}}}{k_B} = \frac{A}{4\ell_P^2} = \frac{4\pi r_s^2 c^3}{4\hbar G} = \frac{4\pi GM^2}{\hbar c} = \mathcal{N}_{\text{esc}}^{\text{Schw}} \quad (18)$$

Equations (17) and (18) are identical. The escrow entropy of a Schwarzschild black hole, computed from the self-binding energy and the Hawking temperature via the static escrow postulate, equals the Bekenstein–Hawking entropy exactly. There is no approximation and no fudge factor; the geometric 4π in the horizon area is recovered exactly by the geometric 4π in the Hawking temperature formula. This is the content of the static escrow postulate’s recovery of Bekenstein–Hawking, originally reported in the framework paper [1], restated here in the Jacobson-translation context.

The Bekenstein–Hawking formula by itself supplies the relation between area and entropy without supplying an interpretation of what the entropy is the entropy of. The escrow framework supplies that interpretation: the horizon entropy is the entropy held in escrow against the local Unruh temperature by the gravitationally bound mass-energy inside the horizon. This is a structural interpretive claim, equivalent in mathematical content to Bekenstein–Hawking but distinct in physical reading. The horizon’s area-as-entropy is not a coincidence

of dimensional analysis nor an axiom requiring separate justification — it is the escrow of the matter trapped behind the horizon, evaluated at the horizon's own Unruh temperature.

3.4 Extension to Reissner–Nordström and Kerr horizons

A natural question for the escrow postulate, raised in adversarial review of this paper, is whether the Schwarzschild recovery in equations (17)–(18) is the only horizon entropy the postulate gets right, or whether the same form $S_{\text{esc}} = |U_{\text{grav}}|/T_H$ continues to recover Bekenstein–Hawking entropies for charged (Reissner–Nordström) and rotating (Kerr) black holes. We report the result of that test, with explicit attention to what is and is not established by it.

For a Reissner–Nordström black hole of mass M and charge Q (in geometrized units $G = c = k_e = 1$, with $k_e = 1/(4\pi\epsilon_0)$ the Coulomb constant), the outer and inner horizons, area, and Hawking temperature are

$$r_{\pm} = M \pm \sqrt{M^2 - Q^2}, A = 4\pi r_+^2, T_H = (r_+ - r_-)/(4\pi r_+^2)$$

The Smarr formula relates these to the mass:

$$M = 2T_H S_{\text{BH}} + \Phi Q$$

where $\Phi = Q/r_+$ is the electrostatic potential at the horizon and $S_{\text{BH}} = A/4$ is the Bekenstein–Hawking entropy. Rearranged,

$$T_H S_{\text{BH}} = (M - \Phi Q)/2 = (M - Q^2/r_+)/2$$

For the escrow postulate $S_{\text{esc}} = |U_{\text{grav}}|/T_H$ to recover S_{BH} exactly, the gravitational binding energy must satisfy

$$|U_{\text{grav}}|^{\text{RN}} = (M - Q^2/r_+)/2 \text{ (geometrized); } (Mc^2/2) - k_e Q^2/(2r_+) \text{ (SI)}$$

i.e. the Schwarzschild binding minus the electrostatic energy stored at the horizon.

For Kerr (mass M , angular momentum $J = Ma$), the horizon radius, area, and Hawking temperature are

$$r_{\pm} = M \pm \sqrt{M^2 - a^2}, A = 4\pi(r_+^2 + a^2), T_H = (r_+ - r_-)/(4\pi(r_+^2 + a^2))$$

and the Smarr formula reads

$$M = 2T_H S_{\text{BH}} + 2\Omega_H J$$

where $\Omega_H = a/(r_+^2 + a^2)$ is the angular velocity of the horizon. Rearranged,

$$T_H S_{\text{BH}} = (M - 2\Omega_H J)/2$$

The required Kerr binding energy is therefore

$$|U_{\text{grav}}|^{\text{Kerr}} = (Mc^2/2) - \Omega_H J$$

the Schwarzschild binding minus the rotational work term.

Both extensions are exact: with these natural identifications of $|U_{\text{grav}}|$, the escrow postulate recovers S_{BH} for RN and Kerr to all displayed digits across the parameter range, including

near extremality (where $T_H \rightarrow 0$ and S_{esc} takes the $0/0$ form whose limit is the finite extremal entropy). The structural pattern is consistent across all three cases: $|U_{\text{grav}}|$ equals the Schwarzschild gravitational binding minus the matter-field chemical-potential terms (ΦQ for charge, ΩJ for rotation) that appear in the first law of black hole mechanics.

We are explicit about the interpretive status of these extensions. They are structural-consistency checks rather than independent predictions. The Smarr formula, which is itself derivable from the area law and the first law, already specifies $T_H S_{\text{BH}}$ as $(M - \Phi Q - 2\Omega_H J)/2$; the escrow postulate's apparent "recovery" of S_{BH} for RN and Kerr is the algebraic statement that $|U_{\text{grav}}|$ is identified with this same quantity. The framework does not independently specify $|U_{\text{grav}}|$ for charged or rotating configurations and then derive S_{BH} ; it adopts the chemical-potential decomposition that the first law already supplies, and reads off the consistency. What the consistency check does establish is that the escrow postulate's form $S_{\text{esc}} = |U_{\text{grav}}|/T_H$ survives the addition of matter fields and rotation without modification, which is a non-trivial structural fact: a postulate that produced S_{BH} only for Schwarzschild and failed for RN or Kerr would be ruled out by these calculations. The escrow framework is therefore consistent with the full first law of black hole mechanics, not just with the Schwarzschild special case.

3.5 What this section does and does not establish

This section does not derive Einstein's field equations from the escrow postulate. Jacobson did that in 1995 [3], using $\delta Q = T \cdot dS$ as input. What this section establishes is that the S appearing in Jacobson's input has the interpretation $S = S_{\text{esc}}$, with the consequence that the derived Einstein equation is the equation of state of the vacuum's escrow thermodynamics.

Two consequences follow. First, the escrow framework does not require any modification of Einstein's equations; it provides an interpretation of why those equations have the structure they do. The cosmological constant Λ appears on the left side of (14) without special handling because Jacobson's derivation produces it as an integration constant in the same way it produces the rest of the geometry. Second, attempts to add a separate "escrow stress-energy tensor" to the right side of Einstein's equation are explicitly ruled out by this derivation: the entire content of δQ on the right side of Jacobson's identity is already the matter stress-energy flux $T_{\mu\nu} k^\mu k^\nu d\lambda$ (where k^μ is the null generator of the horizon and λ is the affine parameter). The escrow contribution lives entirely in the interpretation of the S on the left side, not in any additional source on the right.

The framework's value at this level is the same as it was for time dilation: a conceptual re-reading. Jacobson's derivation, the Bekenstein–Hawking formula, the Tolman temperature law, the Unruh effect, and the static escrow postulate are five separately-derived results that the escrow framework expresses through a common thermodynamic object S_{esc} . None of these results is modified; their interrelation is made transparent.

3.6 A note on what the framework does not commit to

We follow Jacobson in treating equation (14) as "an" equation of state rather than asserting that spacetime is literally a thermodynamic system in some fundamental sense. The thermodynamic analogy is structural — the same Clausius relation, the same area-as-entropy identification, the same local equilibrium condition — not ontological. We do not claim

that spacetime is built out of more fundamental thermodynamic degrees of freedom in the way that an ideal gas is built out of molecules. Such a claim would require identifying the microscopic constituents whose statistical mechanics produces the area law; this is the project of quantum gravity and is not addressed by the escrow framework.

Similarly, we do not claim that the entropy in equation (15) is observer-independent. The Unruh temperature depends on the choice of accelerated observer; the horizon depends on the choice of local Rindler frame; the escrow entropy inherits both dependences. What is observer-independent is the relation $\delta Q = T_U \cdot dS_{\text{esc}}$, which holds for every observer in their own frame. This observer-dependence is a feature of the thermodynamic interpretation, not a bug: it is what distinguishes Jacobson's reading of GR from a global-entropy interpretation in which a unique S is assigned to a spacetime. Different observers see different escrow entropies because they have different local Unruh temperatures; the field equation (14) is what enforces consistency across observers.

The escrow framework's identification of S_{horizon} with S_{esc} therefore does not add a substantive claim beyond what Jacobson's derivation already commits to. It supplies a name and an interpretation for the entropy that Jacobson uses as input.

4 The Modular Hamiltonian Connection

4.1 The standard statement

Dimensional bookkeeping for this section: throughout §IV we follow the standard entanglement-entropy convention in which the modular Hamiltonian K is dimensionless (it appears as the exponent in the reduced density matrix $\rho = \exp(-K)/Z$, so dimensionlessness is required by the exponential's argument). Entanglement entropy S_{EE} in this convention is also dimensionless (a count, measured in nats or equivalently in units of k_B). Where this section writes $T_U \cdot \Delta S_{\text{EE}}$ or $T_U \cdot \Delta\langle K \rangle$, the implicit factor of k_B converting dimensionless entropy to thermodynamic entropy is understood: the energy associated with a modular-Hamiltonian-content change $\Delta\langle K \rangle$ at Unruh temperature T_U is $k_B T_U \cdot \Delta\langle K \rangle$, exactly as in §IV.A's calculation $\Delta K \cdot k_B \cdot T_U (d1) = mc^2$. Sections II and III used the alternative convention in which S is written in J/K from the start (with k_B explicit in each formula); the two conventions differ only by where the k_B is placed, and we use whichever is cleaner in each setting. Equations (19) and (20) below are written in the entanglement-entropy convention; explicit k_B factors are inserted where the equation crosses into thermodynamic-entropy quantities.

Jacobson's derivation [3] uses the horizon area entropy as the input S in $\delta Q = T \cdot dS$, with the area law itself taken as a given. A subsequent body of work, of which the most consequential single result is due to Faulkner, Guica, Hartman, Myers, and Van Raamsdonk (FGH MV, [5,6]), establishes a related but distinct identification: in asymptotically AdS spacetimes, demanding the entanglement-entropy first law $\delta S_{\text{EE}} = \delta\langle K \rangle$ for vacuum-state perturbations across spherical entangling surfaces forces the linearized Einstein equations in the bulk. Here S_{EE} is the entanglement entropy of a region of the boundary CFT, K is the modular Hamiltonian of that region in the vacuum state (the operator whose exponential gives the reduced density matrix on the region), and $\delta\langle K \rangle$ is the expectation value of K in the perturbed state minus its value in the vacuum.

The FGH MV result is structurally similar to Jacobson's but operationally distinct in two

respects. First, it uses entanglement entropy rather than horizon area as the input quantity, with the Ryu–Takayanagi formula [10] supplying the bulk-area dual of the boundary entanglement. Second, it produces linearized Einstein equations rather than the full nonlinear theory, with the linearization being around the AdS vacuum. Subsequent work has extended the result to broader settings, including non-spherical entangling regions and curved-background perturbations, with the unifying structural statement being that the gravitational equations of motion are equivalent to the requirement that relative entropy (a closely related and equally fundamental information-theoretic quantity) be non-negative for vacuum-state perturbations.

The vacuum modular Hamiltonian for a Rindler wedge has an exact form, originally derived by Bisognano and Wichmann [11]:

$$K = (2\pi/(\hbar c)) \int x_{\perp} T_{00}(x) dV$$

where $x_{\perp}$ is the perpendicular distance from the cut and T_{00} is the energy density. This is the same 2π that appears in the Unruh temperature — the modular flow generated by K is exactly the boost flow that defines the Rindler wedge, and the period 2π in boost parameter corresponds to the inverse Unruh temperature. The Bisognano–Wichmann formula gives a sharp prediction for the change in modular Hamiltonian content induced by inserting a localized energy $E = mc^2$ at perpendicular distance $d1$ from the cut:

$$\Delta K = 2\pi \cdot m \cdot d1 \text{ (natural units; } \hbar = c = k_B = 1\text{)}$$

or equivalently, restored to SI units,

$$\Delta K = 2\pi \cdot m \cdot c \cdot d1 / \hbar = \mathcal{N}_{\text{esc}}(d1)$$

(a dimensionless quantity, as required of a modular Hamiltonian content), where $\mathcal{N}_{\text{esc}}(d1)$ is the escrow count of equation (8') evaluated at $r = d1$. The associated energy at the Rindler-observer Unruh temperature is

$$T_U(d1) = \hbar c / (2\pi \cdot d1 \cdot k_B), \Delta K \cdot k_B \cdot T_U(d1) = mc^2$$

— the rest energy of the perturbation. This is the prediction that the lattice paper [2] tests numerically in free scalar field theory on regulated 1+1D and 3+1D lattices.

4.2 The escrow translation

The escrow postulate identifies $S_{\text{esc}} = |U_{\text{grav}}|/T_U$ as the entropy held against the local Unruh temperature by gravitationally bound mass-energy. The FGHMV result uses S_{EE} and the modular Hamiltonian K , neither of which is identical to S_{esc} as defined. The question is: what is the relationship between the escrow entropy and the modular Hamiltonian content?

The relationship is mediated by the local first law. For a vacuum-state perturbation localized in a Rindler wedge, the entanglement-entropy first law $\delta S_{\text{EE}} = \delta \langle K \rangle$ says that the change in entanglement entropy of the wedge (in dimensionless units; see the unit-convention statement at the top of §IV.A) equals the change in expectation value of the wedge's modular Hamiltonian. The product $k_B \cdot T_U \cdot \delta S_{\text{EE}}$ has units of energy and represents the heat content associated with the entanglement change. Identifying this with the gravitational binding energy of the perturbation gives

$$|U_{\text{grav}}| = k_B T_U \Delta S_{\text{EE}} = k_B T_U \Delta \langle K \rangle \quad (19)$$

which, by the escrow postulate (6) $S_{\text{esc}} = |U_{\text{grav}}|/T_U$, means

$$S_{\text{esc}} = k_B \Delta S_{\text{EE}} = k_B \Delta \langle K \rangle \quad (20)$$

with the explicit k_B factor converting the dimensionless entanglement-entropy quantities of the entanglement-gravity literature into the thermodynamic-entropy units in which S_{esc} was defined in §II. The dimensionless content of equation (20) is $S_{\text{esc}}/k_B = \Delta S_{\text{EE}} = \Delta \langle K \rangle$, which is what the lattice computations of [2] compute directly.

We are explicit about the regime-bridging assumptions that this identification carries. The FGHMV result [5,6] is formulated for asymptotically AdS spacetimes with spherical entangling surfaces on the boundary CFT, with the relevant entropy being the entanglement entropy of a boundary subregion and the relevant modular Hamiltonian being the operator generating modular flow for the vacuum reduced density matrix of that subregion. The escrow framework's identification in (19)–(20) applies the same first-law structure to a localized perturbation in a flat-space Rindler wedge, where T_U is the Unruh temperature of an accelerated observer. The bridge between these two settings carries three distinct technical assumptions, each well-established in the entanglement-entropy literature [5,6,11,12,13] but each worth stating individually: (a) the modular-Hamiltonian content of a Rindler wedge in Minkowski vacuum is given by the Bisognano–Wichmann formula $K = (2\pi/(\hbar c)) \int x_{\perp} T_{00} dV$, where $x_{\perp}$ is perpendicular distance from the cut; (b) the entanglement-first-law statement $\delta S_{\text{EE}} = \delta \langle K \rangle$ carries over from the boundary CFT setting to a flat-space Rindler wedge in the linear-response regime, with no curvature corrections to the first law at this order; (c) the gravitational binding energy of the localized perturbation, identified with the source energy mc^2 , plays the same role on the matter side that boundary-CFT energy density plays in the AdS setting. Each of (a), (b), (c) is standard in the literature but each is an additional assumption beyond the bare FGHMV statement. We treat all three as assumptions rather than as derived statements, and the framework's modular-Hamiltonian leg therefore inherits whatever caveats attend each of these standard identifications in the broader entanglement-gravity literature.

The escrow entropy of a localized mass perturbation, under this identification, equals both the change in entanglement entropy of the wedge containing it and the change in expectation value of the wedge's modular Hamiltonian. Equation (20) is the strongest interpretive content of the escrow framework: the escrow entropy is the modular-Hamiltonian content of a localized gravitational perturbation, with the FGHMV result then identifying the consistency requirement on this content with linearized Einstein equations.

If equation (20) held exactly, the modular Hamiltonian content of a localized mass m at perpendicular distance $d1$ from the cut would equal the escrow count $\mathcal{N}_{\text{esc}}(d1) = 2\pi \cdot m \cdot c \cdot d1 / \hbar$ (the dimensionless content of equation 8' with $r = d1$). The Bisognano–Wichmann formula gives exactly this. Equation (20) therefore predicts that lattice computations of ΔK in regulated free scalar field theory should yield the BW linear asymptote with prefactor unity, i.e. $\Delta K = \mathcal{N}_{\text{esc}}(d1)$ at small $d1$ in the half-space limit. This is the prediction the lattice paper [2] tests numerically.

4.3 What the lattice tests

The lattice paper [2] computes ΔK directly on regulated free scalar lattices in 1+1D (chain, N up to 3000) and 3+1D (cubic, N up to 14) using the Williamson-decomposition method, with two protocols: a two-mass protocol with masses at perpendicular distances d_1 and d_2 from the cut, and a single-mass-in-A protocol with a single mass at d_1 (the latter being the cleaner test of the BW prediction). The numerical results, summarized in [2], are:

In 1+1D at small d_1 ($d_1 \in [2, 6]$ at $m = 1$, $N = 1000$), the local power-law exponent of ΔK vs d_1 is $+1.02 \pm 0.05$, indistinguishable from the BW linear asymptote $\alpha = 1$. The prefactor is approximately 0.030–0.037, corresponding to a suppression of approximately 1/30 of the literal BW value. At larger d_1 the local exponent decreases smoothly toward zero, reflecting geometric corrections to the BW half-space assumption (Dirichlet boundary conditions, finite subsystem-A depth). The peak prefactor is independently reproducible from the lattice paper’s published methodology, with both V1 (relative-entropy positivity) and V2 (modal-versus-original-basis cross-check) sanity checks passing at every d_1 in the sweep; the analytic origin of the prefactor remains the open theoretical problem identified in this section.

In 3+1D at all d_1 the lattice can resolve ($N \leq 14$, $d_1 \leq 6$), $\Delta K(d_1)$ is peaked at $d_1 \approx 2$ and decays as d_1^{-2} to -3 for $d_1 \geq 3$. The BW linear asymptote is not recovered at any d_1 the lattice can resolve. The peak structure reproduces across $N = 12$ and $N = 14$, indicating that the result is not a finite-size artifact at the lattice sizes used.

Taken at face value, these results constrain equation (20) in two ways. In 1+1D, the equation holds approximately within a small- d_1 window with a prefactor suppression of 1/30; the prefactor has no presently-calculated origin, and finding one is the principal open theoretical question. In 3+1D, the equation does not hold within the d_1 range the lattice can resolve; whether it can be recovered at much larger d_1 inaccessible to the present lattice (corresponding to a slower rate of approach to the continuum BW asymptote in higher dimensions) is an open empirical question.

4.4 What this section does and does not establish

This section does not establish equation (20) as a derived identity. It identifies (20) as the natural escrow reading of the FGHMV result, with [2] supplying empirical content on whether the identification holds when both sides are computed independently. The 1+1D partial recovery and the 3+1D non-recovery within accessible d_1 are the present state of evidence; neither rules out (20) fundamentally, and neither establishes it without qualification.

Three distinct theoretical questions are sharpened by this section. First, what is the calculable origin of the 1/30 prefactor in 1+1D? A candidate explanation flagged in [2] is that the lattice-regularized free scalar at $m = 1$ has correlation length $\xi \sim 1/m \sim 1$ lattice spacing, making the localized mass a UV-scale perturbation by the lattice’s own measure, while the BW formula is an infrared / long-wavelength prediction. The 1/30 suppression may reflect a UV/IR mismatch correctable by a first-principles calculation of the lattice-regulated modular Hamiltonian. The Dirichlet boundary conditions used for the lattice differ from the free BCs assumed in the standard BW derivation, and the boundary-correction factor is another candidate contribution. Resolving the prefactor’s origin from first principles is the single most consequential follow-up calculation the framework presently faces.

Second, does the 3+1D non-recovery reflect a genuinely shorter characteristic length scale of the 3+1D modular Hamiltonian content, or a slower rate of approach to the continuum BW limit due to the smaller conformal-symmetry group of the 3+1D massless free scalar? The 1+1D massless scalar is a CFT with infinite-dimensional Virasoro symmetry; the 3+1D massless scalar has only the finite-dimensional conformal group, and the rate at which the lattice-regularized modular content approaches the continuum BW asymptote may differ between them for this reason. The 3+1D test on $N \gtrsim 20$ lattices (computationally feasible on dedicated GPU hardware but not on the present authors' CPU resources) would distinguish these two interpretations.

Third, what is the dimensional-dependence theorem, if any, connecting 1+1D BW partial recovery to the 3+1D structure observed in [2]? A 2+1D test on the same protocol would be the natural intermediate case and would determine whether the dimensional dependence is monotone. The present authors leave this as a future calculation.

4.5 A note on what the framework does not commit to

We do not claim that the lattice paper [2]'s partial recovery in 1+1D establishes equation (20) as a derived identity. The $1/30$ prefactor is an unexplained suppression, not a confirmed match; the framework's interpretive content at this level depends on that prefactor having a calculable origin, and the calculation has not been performed. The framework is at risk of being read as having empirical confirmation it does not yet have, and we explicitly disclaim that reading: [2] is consistent with the escrow framework's modular-Hamiltonian identification at the level of qualitative structure (positive sign, linear scaling at small d_1 , the right dimensionless quantity $\mathcal{N}_{\text{esc}}$), but it is not yet quantitatively confirmed at the level of the absolute prefactor.

We do not claim that the 3+1D non-recovery rules out the escrow framework. As stated above, the rate-of-approach question is unresolved by the present lattice sizes, and a finding that 3+1D BW is recovered at much larger d_1 inaccessible to the present lattice would resolve the non-recovery as a finite-size effect rather than a fundamental obstruction. The framework's commitment is to the postulate (6) and the interpretive content of equations (10), (15), (17), and (20); empirical confirmation of these at the lattice level is partial and dimensional, and we describe it as such.

We do not invoke any extension of equation (20) beyond the FGHMV setting. The FGHMV result is established for asymptotically AdS spacetimes with spherical entangling surfaces and vacuum-state perturbations. Extensions to other settings (flat-space, non-vacuum states, non-spherical regions, large perturbations) are areas of active research [12,13], and we do not claim that the escrow framework's identification (20) carries through to those extensions automatically. The framework's modular-Hamiltonian content as expressed in (20) is, like the FGHMV result, presently a statement about the linear-response / small-perturbation regime in asymptotically AdS-like settings.

5 Consolidated Guardrails

Sections II.E, III.F, and IV.E each conclude with explicit statements of what the escrow framework does not commit to in the local context of that section. We consolidate those statements here as a single list, for use in subsequent work and in defense against the failure modes documented in the framework's history.

5.1 No escrow stress-energy tensor

The escrow framework does not introduce a separate “escrow stress-energy tensor” $T_{\mu\nu}^{\text{escrow}}$ to be added to the matter stress-energy on the right side of Einstein’s equation. The reason is structural and matches the discussion in §II.E: Einstein’s equations are an algebraic identity between the matter stress-energy tensor $T_{\mu\nu}^{\text{matter}}$ and the geometric Einstein tensor $G_{\mu\nu}$, and the Bianchi identities ($\nabla^\mu G_{\mu\nu} = 0$) require $\nabla^\mu T_{\mu\nu}^{\text{matter}} = 0$ — the matter content already saturates the structure of conserved sources. Gravitational binding energy is not a localizable tensorial quantity in general relativity; it is encoded non-locally in the geometry itself (the difference, for an isolated system, between the ADM mass at infinity and the integrated rest mass on a spatial slice), and it cannot be added to $T_{\mu\nu}^{\text{matter}}$ as an independent source. Attempting to add a separate escrow source on the right side of Einstein’s equation would either violate the Bianchi identities or double-count the gravitational configuration: the geometry on the left side already responds to the full configuration through its non-local encoding of binding energy, and the matter content on the right side already saturates the conservation structure. The escrow framework supplies an interpretive dictionary for the existing fields; it does not augment the field equation with any new contribution. Jacobson’s derivation [3] makes this constraint particularly sharp: the entire content of δQ on the right side of his Clausius identity is already the matter stress-energy flux $T_{\mu\nu}^{\text{matter}} k^\nu dA^\mu$, with no remaining room for an additional escrow source. Attempts to extend the framework with additional source terms reduce to either trivial restatements of the matter source (no new physics) or violations of Jacobson’s derivation (which would require modifying the input $\delta Q = T \cdot dS$, not merely adding to the output).

5.2 No vacuum computational substrate

The escrow framework does not invoke a “vacuum computational bandwidth,” “information processing rate,” or digital substrate reading of the static escrow postulate. Such readings appear in some popularizations of entropic gravity — typically expressed as “the vacuum is a computer with finite processing capacity, and gravity emerges because the computer’s bandwidth is partially consumed by holding the escrow” — but they are not part of the postulate as stated in [1]. The local rate of escrow production at radius r is a thermal quantity, set by $T_U(r)$, and is correctly described in the vocabulary of equilibrium thermodynamics. No additional ontological structure is required to make the framework’s predictions hold. The framework’s interpretation of why time dilation gives the rate it gives (equation 10) is the thermodynamic Tolman-temperature reading, not a computational-bandwidth reading.

5.3 No claim about microscopic vacuum constituents

The escrow framework, like Jacobson’s derivation, treats the Einstein equation as “an” equation of state of the vacuum rather than asserting that spacetime is literally a thermodynamic system in some fundamental sense. The thermodynamic analogy is structural — the same Clausius relation, the same area-as-entropy identification, the same local equilibrium condition — not ontological. We do not claim that spacetime is built out of more fundamental thermodynamic degrees of freedom in the way that an ideal gas is built out of molecules. Such a claim would require identifying the microscopic constituents whose statistical mechanics produces the area law; this is the project of quantum gravity and is not addressed here. The escrow

framework operates entirely at the level of the macroscopic thermodynamic identity; it makes no commitment to the underlying microphysics.

5.4 Observer-dependence of S_{esc}

The escrow entropy S_{esc} is not observer-independent. The Unruh temperature depends on the choice of accelerated observer; the local Rindler horizon depends on the choice of local Rindler frame; the escrow entropy inherits both dependences. What is observer-independent is the relation $\delta Q = T_U \cdot dS_{\text{esc}}$ and its consequence (Einstein's equation, equation 14), which hold for every observer in their own frame. This observer-dependence is a feature of the thermodynamic interpretation, not a bug: it is what distinguishes the escrow framework's reading of GR from a global-entropy interpretation in which a unique S is assigned to a spacetime. Attempts to define a global, observer-independent S_{esc} reduce either to the global entropy of the universe (which is not what the static escrow postulate is about) or to the integrated horizon-flux entropy (which is Jacobson's input, not new content).

5.5 No empirical confirmation of equation (20)

The lattice paper [2]'s partial recovery of the Bisognano–Wichmann linear asymptote in 1+1D at small $d1$ with a $1/30$ prefactor is not a confirmation of equation (20). The $1/30$ suppression is an unexplained empirical observation; resolving its origin from first principles (UV/IR mismatch, Dirichlet boundary correction, lattice regularization artifact, or fundamental modification of the framework) is the principal open theoretical question. We disclaim any reading of [2] as providing empirical confirmation of the escrow framework's modular-Hamiltonian identification, and we similarly disclaim any reading of [2]'s 3+1D non-recovery as fundamentally ruling out the framework. The empirical situation is partial, dimensional, and unresolved; we describe it as such.

5.6 No claim of new equations of motion

The escrow framework does not produce new equations of motion. All gravitational dynamics in the framework's purview reduce to Einstein's equation (14), derived in the standard way via Jacobson [3] from $\delta Q = T \cdot dS$, with the escrow framework's contribution being the interpretation of the S as S_{esc} . Attempts to extract new equations of motion from the escrow postulate — such as the C8 entropy-current attempt documented in [9] — have reduced to either rate-equation reformulations of known results (no new physics) or to violations of the framework's interpretive constraints (no new defensible physics). The framework's mathematical content is the static escrow postulate (6) and the consequences derived in [1]; its interpretive content is the dictionary developed in the present paper. No new dynamical content is claimed.

5.7 $|U_{\text{grav}}|$ is not given a single covariant definition

Across the four legs of the translation, the symbol $|U_{\text{grav}}|$ denotes different quantities. In §II (test mass in a static gravitational potential), $|U_{\text{grav}}| = GMm/r$ is the Newtonian gravitational potential energy of a probe particle. In §III (Schwarzschild horizon), $|U_{\text{grav}}| = Mc^2/2$ is the self-binding energy of the configuration at horizon scale, defined via GM^2/r_s with r_s the Schwarzschild radius. In §III.D (Reissner–Nordström and Kerr), $|U_{\text{grav}}|$ is identified with the chemical-potential decomposition $(M - \Phi Q - 2\Omega_H J)/2$ that the Smarr formula already

supplies. In §IV (modular-Hamiltonian content of a localized perturbation), $|U_{\text{grav}}|$ is the energy of the localized perturbation, formally identified with mc^2 for a point mass of rest mass m . These four definitions are not connected by a single covariant, observer-independent construction in standard general relativity. Quasilocal energy proposals (Brown–York, Hawking mass, Komar integrals for stationary configurations) are candidate constructions that could in principle supply such a covariant definition, but the escrow framework does not currently adopt one. We are explicit that the framework’s $|U_{\text{grav}}|$ is presently a family of regime-specific identifications glued together by common notation, and that providing a single covariant definition is an open theoretical task. This limitation does not undermine the consistency checks within each regime, but it does mean that the framework’s claim to a ‘single thermodynamic object’ is partly notational: the object is S_{esc} in each regime, but the $|U_{\text{grav}}|$ feeding into S_{esc} is regime-specific.

5.8 T_U is used with two related but distinct conventions

The Unruh temperature T_U is used in this paper with two conventions that are related by the Tolman shift but are not identical. In §II (test mass in a Newtonian potential), T_U is the local Unruh temperature of a stationary observer at radius r , computed from the local proper acceleration $a = GM/r^2$ via

$$T_U = \hbar a / (2\pi c k_B) \text{ (local Unruh; §II convention)}$$

This T_U is what a local detector would register. In §III (Schwarzschild horizon), T_U is taken to equal the Hawking temperature

$$T_H = \hbar \kappa / (2\pi c k_B), \kappa = c^4 / (4GM) \text{ (asymptotic Hawking; §III convention)}$$

where κ is the surface gravity of the horizon. The Hawking temperature is the redshifted (asymptotic) Unruh temperature of an observer hovering just outside the horizon; the local Unruh temperature diverges at the horizon, but its product with the Tolman redshift factor approaches the finite asymptotic value T_H . The two T_U conventions are therefore related by the Tolman shift

$$T_{\text{local}} \cdot \sqrt{-g_{tt}} = T_{\infty}$$

(the same exact Tolman relation invoked in §II.A), and they agree in the appropriate limits. We use ‘ T_U ’ uniformly in the body for notational economy, with the understanding that the local vs. asymptotic distinction is contextually determined by the regime under discussion. This unification at the level of notation is not a unification at the level of distinct physical observables: the local Unruh temperature and the asymptotic Hawking temperature are different operational quantities, related by the Tolman relation rather than identical.

6 Conclusion

The static gravitational entropy escrow postulate of [1] admits an interpretive translation onto the established thermodynamic-gravity literature. The translation comprises four legs. In Section II, gravitational time dilation is recovered as the Tolman temperature shift expressed in escrow language, with the dimensionless escrow of a test mass identified as the escrow count

$$\mathcal{N}_{\text{esc}} \equiv 2\pi r/\lambda_C$$

— the number of reduced Compton wavelengths fitting into the orbit’s circumference. In Section III, Jacobson’s derivation of Einstein’s field equations from $\delta Q = T \cdot dS$ at local Rindler horizons becomes the differential form of the static escrow postulate, with the Bekenstein–Hawking entropy of a Schwarzschild black hole recovered exactly (to all displayed digits, with no fudge factor) as

$$S_{\text{BH}}/k_B = \mathcal{N}_{\text{esc}}^{\text{Schw}} = 4\pi(M/m_P)^2$$

the escrow entropy of the configuration computed from its self-binding energy and the Hawking temperature. In Section IV, the FGHMV identification of the entanglement-entropy first law with linearized Einstein equations becomes the statement that the escrow entropy of a localized mass perturbation equals the modular-Hamiltonian content of the wedge containing it,

$$\Delta K = \mathcal{N}_{\text{esc}}(d1) \quad (\text{BW prediction})$$

Section V consolidates the explicit guardrails against the failure modes that the framework’s history has surfaced.

None of these results modifies any standard equation of general relativity, of quantum field theory, or of horizon thermodynamics. The escrow framework’s contribution at the level treated in this paper is interpretive: it identifies a single thermodynamic object — S_{esc} — through which four otherwise-separately-derived results are expressed as faces of one identity. We note again (§V.G) that the ‘single object’ description is partly notational, as the $|U_{\text{grav}}|$ feeding into S_{esc} takes regime-specific forms across §II, §III, §III.D, and §IV that have not yet been connected by a single covariant construction. The framework’s broader empirical anchors live in [1] and depend on the horizon-thermodynamic content of the postulate, which is unchanged by the present paper.

Three open theoretical problems are sharpened by the present translation. First, the calculable origin of the $1/30$ prefactor that suppresses the lattice paper’s 1+1D modular-Hamiltonian content relative to the literal Bisognano–Wichmann asymptote remains unresolved. A first-principles calculation of the lattice-regularized modular Hamiltonian on the Dirichlet half-chain, accounting for the UV/IR mismatch between localized lattice masses and the long-wavelength BW prediction, is the calculation that would either explain the suppression (rehabilitating the framework’s modular-Hamiltonian interpretation) or fail to explain it (constraining the framework). This is the single most consequential follow-up calculation the framework presently faces, and we identify it as the principal short-term research priority.

Second, the 3+1D non-recovery of the BW asymptote within the lattice sizes accessible to [2] is unresolved as to whether it reflects a genuinely shorter characteristic length scale in higher dimensions or a slower rate of approach to the continuum BW asymptote due to the smaller conformal-symmetry group of the 3+1D massless free scalar. Lattice computations at $N \gtrsim 20$ (feasible on dedicated GPU hardware) would distinguish these interpretations

and would constitute a decisive next-generation test. A 2+1D test would supply a natural intermediate dimensional case.

Third, the dimensional-dependence theorem connecting 1+1D and 3+1D modular-Hamiltonian content is unidentified. Whether such a theorem exists — and what its form would be — is a question that the FGHMV framework, the Bisognano–Wichmann formula, and the lattice computations of [2] are all positioned to address jointly. The escrow framework’s contribution to this question is the explicit identification of the dimensionless quantity $\mathcal{N}_{\text{esc}} \equiv 2\pi r/\lambda_C$ as the candidate organizing variable; whether it is the right variable, and whether its dimensional dependence is calculable, are open.

A structural risk worth naming explicitly is that of retrofitted inevitability: the possibility that the escrow framework’s apparent unification of the four results translated in this paper is an artifact of having seen those four results first and then constructed a dimensional combination ($S_{\text{esc}} = |U_{\text{grav}}|/T_U$) that connects them in retrospect, rather than the proposal of an organizing variable that would have constrained the four results in advance had it been known. The framework’s escape from this failure mode is not interpretive but calculational. Equation (20) of §IV makes a sharp numerical prediction — the modular-Hamiltonian content of a localized mass perturbation should equal $\mathcal{N}_{\text{esc}}(d1) = 2\pi \cdot m \cdot d1$ in natural units, with prefactor unity — and the lattice paper [2] tests it. The 1+1D result is a partial confirmation (correct linear scaling, suppressed prefactor); the 3+1D result is a non-recovery within accessible $d1$. Whether equation (20) is a real identity with a calculable correction factor, or an arbitrary dimensional combination that the lattice partially fits in 1+1D and fails to fit in 3+1D for unrelated reasons, depends on whether the $1/30$ prefactor has a first-principles origin in lattice-regularization theory. A successful first-principles calculation would move the framework from “interpretive synthesis with retrofitted-inevitability risk” to “organizing variable with a calculable correction factor.” A failed calculation would constrain the framework’s modular-Hamiltonian leg. Either outcome is informative, and either outcome distinguishes the framework’s content from a pure relabeling exercise. The framework’s status as a candidate proposal rather than a derived theory is therefore reducible to a single open calculational question, which we name as such.

To make the falsification structure explicit, we identify five conditions under which specific legs of the framework would be ruled out. (i) If a first-principles lattice calculation accounting for Dirichlet boundary conditions, lattice UV cutoff effects, and the finite half-chain depth produces a prefactor that disagrees with the lattice-measured 0.030 – 0.037 at the same protocol, the modular-Hamiltonian leg expressed in equation (20) is constrained or ruled out in its present form. (ii) If a 3+1D lattice computation at $N \geq 24$ with $d1$ swept up to $d1 \leq N/3 \approx 8$ fails to show approach toward the BW linear asymptote (i.e., the peak-then-decay structure of $\Delta K(d1)$ observed at $N = 14$ persists with the peak location not migrating significantly toward larger $d1$ and the decay exponent not flattening toward zero), the modular-Hamiltonian leg of the framework is constrained in 3+1D, the 3+1D non-recovery becomes a structural feature requiring explanation rather than a finite-size artifact, and the framework retreats to its horizon-thermodynamic and Tolman-redshift content only. We name $N \geq 24$ with $d1 \leq N/3$ as the concrete threshold rather than leaving ‘any accessible $d1$ ’ unbounded; computation at larger N remains informative but the framework would already be constrained at this

threshold. (iii) If a charged or rotating black hole calculation reveals that the escrow postulate cannot recover S_{BH} for a configuration where the first law of black hole mechanics applies (i.e., a case not absorbed by the Smarr formula’s chemical-potential decomposition), the horizon-thermodynamic leg of the framework is constrained. (iv) If a strong-field calculation (e.g., the exact Schwarzschild redshift for a compact object, or the analogous Kerr or Reissner–Nordström redshift) shows that the escrow translation of equation (9) produces a quantitatively wrong prediction when extended beyond the Newtonian weak-field limit — i.e., the iterated escrow identity does not reproduce $\sqrt{(-g_{tt})}$ at strong field within reasonable precision — the Tolman-redshift leg is constrained to remain a weak-field effective description, and the framework explicitly disclaims any strong-field reading of §II’s translation. (v) If a rigorous theorem shows that Jacobson’s S_{horizon} cannot be identified with any quasilocal energy construction (such as Brown–York energy, Hawking mass, or a Komar integral on the horizon) in a manner consistent with the escrow postulate’s structural form $S = |U|/T$ — i.e., if the open theoretical task named in §V.G is shown to have no positive resolution — the framework’s claim that S_{esc} and S_{horizon} refer to the same underlying object is undermined, and the framework’s Jacobson-leg content reduces to a notational coincidence rather than a structural identification. In each case the falsification is partial, removing one leg of the five-condition translation without necessarily ruling out the others. We note these explicitly rather than relying on implicit logical structure.

We commit, in advance of the calculations named in conditions (i)–(v), to the following pre-registered retractions. If condition (i) is satisfied (the first-principles prefactor calculation disagrees with the lattice-measured 0.030–0.037 at the same protocol), the modular-Hamiltonian leg of the framework will be withdrawn, and any successor version of this paper will publish only the time-dilation (§II) and Jacobson/Bekenstein–Hawking (§III) legs. If condition (ii) is satisfied (the $N \geq 24$ lattice persistently shows the 3+1D peak-then-decay structure), the 3+1D applicability of the framework will be withdrawn; the modular-Hamiltonian leg will be scoped explicitly to 1+1D only. If condition (iii) is satisfied (a Smarr-resistant black-hole configuration fails to recover S_{BH}), the horizon-thermodynamic leg will be withdrawn. If condition (iv) is satisfied (the strong-field test produces a quantitatively wrong redshift prediction in the iterated escrow translation), §II will be explicitly scoped to the Newtonian weak-field limit and no strong-field claim will be defended. If condition (v) is satisfied (the §V.G open task is shown to have no positive resolution), the finite-level identification $S_{\text{horizon}} = S_{\text{esc}}$ for static black holes (§III.C and §III.D) will be retracted as a structural claim and retained only as a notational coincidence; the differential-level identification $\delta S_{\text{horizon}} = \delta S_{\text{esc}}$ (§III.B), which is the content of the escrow reading of Jacobson’s derivation and depends on the first law rather than on a covariant binding-energy construction, does not depend on the §V.G open task and would remain intact. These pre-registered commitments are intended to make the framework’s falsification structure costly rather than rhetorical: a Popperian theory carries with it the obligation to specify, in advance, what would constitute its failure and what one would do in response. We accept that obligation here and bind any future version of this paper to honor it.

The escrow framework joins an existing body of thermodynamic and entropic interpretations of gravity. Jacobson’s 1995 derivation [3], to which Section III is most directly indebted,

derives Einstein’s equations from $\delta Q = T \cdot dS$ at local Rindler horizons. Verlinde’s 2011 entropic gravity proposal and its 2017 emergent-gravity extension take a related but distinct approach, deriving Newton’s law from an entropy gradient on a holographic screen and relating the deep-MOND scale to the de Sitter horizon temperature. Padmanabhan’s emergent-gravity program has long emphasized that gravitational field equations have the standing of a thermodynamic equation of state and that horizon thermodynamics is the natural variable for understanding gravity. The covariant entropy bound (Bousso) constrains the entropy of light-sheets and provides a related but operationally distinct entropy/area relation. The escrow framework’s specific contribution within this literature is the identification of the dimensionless ratio $N_{\text{esc}} \equiv 2\pi r/\lambda_C$ (and its horizon analog $N_{\text{esc}}^{\text{Schw}} \equiv 4\pi(M/m_P)^2$) as a candidate organizing variable for the thermodynamic interpretation, and the explicit modular-Hamiltonian identification of equation (20) as the falsifiable empirical content. We do not claim that the escrow framework supersedes Jacobson, Verlinde, Padmanabhan, or Bousso, or that it occupies a unique position relative to them; we claim that it is a specific proposal within the broader thermodynamic-gravity program, with the conceptual translation of this paper showing how it connects to the established literature and with the lattice paper [2] supplying its principal falsifiable test. For transparency on the citation chain: references [3]–[8] and [10]–[13] are peer-reviewed publications, while references [1], [2], and [9] are the present author’s Zenodo preprints that have not yet been submitted to peer review; the framework’s empirical anchor (the lattice paper [2]) and its theoretical anchor (the framework paper [1]) are therefore the author’s own work, awaiting independent replication and peer review.

The framework as it presently stands is best characterized as a candidate organizing variable for the thermodynamic interpretation of gravity, with the static escrow postulate as the proposed organizing relation, the lattice paper as the empirical anchor for its modular-Hamiltonian interpretation, the framework paper as the empirical anchor for its horizon-thermodynamic interpretation, and the present paper as the explicit conceptual re-reading connecting these to the established thermodynamic-gravity literature. The translation is internally consistent and recovers standard results without modification. Whether the candidate organizing variable proves to be the right one is a question that the open theoretical problems above will determine, and that future empirical and analytical work will adjudicate.

Data, Code & Resources

Manuscript source and archived record: <https://github.com/Windstorm-Institute/escrow-spacetime> (Zenodo: 10.5281/zenodo.20126090). Experiments, datasets, and analysis code: <https://github.com/Windstorm-Labs/escrow-spacetime>. Windstorm Institute: <https://windstorminstitute.org>. Windstorm Labs: <https://windstormlabs.com>.

References

- [1] G. L. Whitmer III, Gravitational Entropy Escrow: An Interpretive Synthesis of Thermodynamic Approaches to Gravity, Zenodo (2026), doi:10.5281/zenodo.20031931.
- [2] G. L. Whitmer III, A Lattice QFT Test of the Static Escrow Postulate: 1+1D and 3+1D Falsification, with a Suggestive Modular-Hamiltonian Window in 1+1D, Zenodo (2026), doi:10.5281/zenodo.20057537.
- [3] T. Jacobson, Thermodynamics of Spacetime: The Einstein Equation of State, Phys. Rev. Lett. 75, 1260 (1995), arXiv:gr-qc/9504004.

- [4] R. C. Tolman and P. Ehrenfest, Temperature Equilibrium in a Static Gravitational Field, *Phys. Rev.* 36, 1791 (1930).
- [5] T. Faulkner, M. Guica, T. Hartman, R. C. Myers, and M. Van Raamsdonk, Gravitation from Entanglement in Holographic CFTs, *JHEP* 03, 051 (2014), arXiv:1312.7856.
- [6] N. Lashkari, M. B. McDermott, and M. Van Raamsdonk, Gravitational Dynamics from Entanglement “Thermodynamics”, *JHEP* 04, 195 (2014), arXiv:1308.3716.
- [7] J. D. Bekenstein, Black Holes and Entropy, *Phys. Rev. D* 7, 2333 (1973).
- [8] S. W. Hawking, Particle Creation by Black Holes, *Commun. Math. Phys.* 43, 199 (1975).
- [9] G. L. Whitmer III, On the Status of Local Entropy-Current Extensions of the Gravitational Entropy Escrow Framework: A Clarification Note, *Zenodo* (2026), doi:10.5281/zenodo.20041991.
- [10] S. Ryu and T. Takayanagi, Holographic Derivation of Entanglement Entropy from AdS/CFT, *Phys. Rev. Lett.* 96, 181602 (2006), arXiv:hep-th/0603001.
- [11] J. J. Bisognano and E. H. Wichmann, On the Duality Condition for Quantum Fields, *J. Math. Phys.* 17, 303 (1976).
- [12] H. Casini, M. Huerta, and R. C. Myers, Towards a derivation of holographic entanglement entropy, *JHEP* 05, 036 (2011), arXiv:1102.0440.
- [13] T. Faulkner, F. M. Haehl, E. Hijano, O. Parrikar, C. Rabideau, and M. Van Raamsdonk, Nonlinear Gravity from Entanglement in Conformal Field Theories, *JHEP* 08, 057 (2017), arXiv:1705.03026.