

A Lattice Quantum Field Theory Test of the Static Escrow

Postulate: 1+1D and 3+1D Falsification, with a Suggestive Modular-Hamiltonian Window in 1+1D

Grant Whitmer

Windstorm Institute, Fort Ann, New York 12827, USA

grant@windstorminstitute.org • ORCID: 0009-0007-3224-755X

May 2026

Abstract

We test the static gravitational entropy escrow postulate $S_{\text{esc}} = |U_{\text{grav}}|/T_{\text{Unruh}}$ of Whitmer [1] using lattice quantum field theory. In 1+1D, a 295-point scan of a free massless scalar (lattice sizes up to $N = 3000$) finds the dimensionless ratio $R = S_{\text{ent}}/S_{\text{esc}}$ spanning 10.56 orders of magnitude; the mass-induced entanglement $\Delta S = S_{\text{ent}} - S_{\text{vac}}$ is uniformly negative and lattice-converged. In 3+1D, an N^3 -site scan confirms the Bombelli–Srednicki area law ($S_{\text{vac}}/N^2 \rightarrow 0.0228$) and shows that R has no finite continuum limit, because the bipartition entropy is dominated by the area-law vacuum term while the mass-induced ratio R_{Δ} is bounded by 10^{-3} ; two independent code paths agree to five decimals. Mutual information decays as L^{-4} , opposite to the linear growth the postulate requires. The literal identification of S_{esc} with bipartition entanglement entropy is therefore ruled out in both dimensions. The modular Hamiltonian content ΔK , evaluated under the Bisognano–Wichmann conjecture, shows a positive, approximately linear-in- d_1 window at $m = 1$ (an empirical non-perturbative feature, *not* a linear-response recovery of the BW asymptote—in the linear-response regime ΔK is negative and d_1 -flat), with prefactor $\approx 1/30$ of the literal BW value ($\approx 1/7$ when energy-normalized). A companion paper [11] finds no BW recovery in 3+1D within the accessible d_1 range. The framework’s horizon-limit recoveries are independent of these flat-space tests.

1 Introduction

The Gravitational Entropy Escrow framework [1] proposes that gravitational binding energy is identified with entropy held in escrow against the local Unruh temperature,

$$S_{\text{esc}} = \frac{|U_{\text{grav}}|}{T_{\text{Unruh}}}, \quad (1)$$

where U_{grav} is the Newtonian binding energy of a configuration and $T_{\text{Unruh}} = \hbar a / (2\pi c k_B)$ is the Unruh temperature corresponding to the local acceleration a . For two equal point masses m at separation L , the prediction simplifies in lattice natural units ($c = \hbar = k_B = 1$, lattice spacing = 1) to

$$S_{\text{esc}}/k_B = 2\pi m L, \quad (2)$$

with G_{eff} and the source mass cancelling identically. The framework recovers Newton’s law via the Verlinde construction [2], Bekenstein–Hawking entropy at horizons via direct integration of Eq. (1) with surface gravity, and the deep-MOND acceleration scale $a_0 = c^2 \sqrt{\Lambda/3}/(2\pi)$ from the de Sitter horizon-temperature floor.

The central claim of [1] is interpretive: that $|U_{\text{grav}}|/T_{\text{Unruh}}$ has the dimensions of, and represents, an entropy held in escrow against the binding. A companion note [3] ruled out the natural-looking covariant local-current extension of Eq. (1) as a notational reformulation of the saturated Bekenstein bound [4]. With the dynamical extension closed off, the static identification (1) becomes the load-bearing claim that requires independent testing.

This paper reports such a test. We compute several QFT entropy quantities of a free massless scalar in the presence of two localized mass distributions and compare to the prediction (2), in 1+1D using Peschel’s exact formula for Gaussian states [5] and in 3+1D on cubic lattices. Three independent entropy measures are tested: bipartition entanglement entropy, mutual information between regions, and modular Hamiltonian content evaluated under the Bisognano–Wichmann conjecture [6]. The headline result is that the literal identification of S_{esc} with bipartition entanglement entropy fails in both 1+1D and 3+1D; the mutual information measure confirms this independently; and the modular Hamiltonian content gives a partial result—a positive, approximately linear ΔK window at $m = 1$ with the right shape but suppressed magnitude, and the opposite sign in the linear-response regime.

2 Methodology

2.1 Lattice setup (1+1D)

We discretize a free massless scalar field on a 1+1D lattice with N sites and Dirichlet boundary conditions at both walls (avoiding the periodic-BC zero mode). The Hamiltonian is

$$H = \frac{1}{2} \sum_i [\pi_i^2 + (\phi_{i+1} - \phi_i)^2 + m_i^2 \phi_i^2] = \frac{1}{2} [\pi^\top \pi + \phi^\top K \phi], \quad (3)$$

$$K = 2I - S - S^\top + \text{diag}(m_i^2), \quad (4)$$

where S is the lattice shift operator. Two equal localized masses are placed at sites $(N - L)/2$ and $(N + L)/2$, on opposite sides of the bipartition cut at site $N/2$ and at distance $L/2$ each from the cut. Single-site delta masses are used; Section IV.C verifies that the conclusions are invariant under Gaussian smearing. The parameter scan covers $N \in \{100, 300, 1000, 3000\}$, $L \in \{2, 4, 6, 10, 20, 40, 80, 160, 320, 640\}$, and $m \in [10^{-3}, 10^2]$ sampled across orders of magnitude, for 295 distinct (N, L, m) points.

2.2 Correlators and Peschel formula

The ground state of (3) is Gaussian, with closed-form correlators

$$\langle \phi_i \phi_j \rangle = \frac{1}{2} (K^{-1/2})_{ij}, \quad \langle \pi_i \pi_j \rangle = \frac{1}{2} (K^{1/2})_{ij}. \quad (5)$$

Restricting to subsystem A (sites $[0, N/2)$) gives reduced covariances $M_\phi = \langle \phi \phi^\top \rangle|_A$ and $M_\pi = \langle \pi \pi^\top \rangle|_A$. The symmetric product $M_{\text{sym}} = M_\phi^{1/2} M_\pi M_\phi^{1/2}$ has eigenvalues $\nu_k^2 \geq \frac{1}{4}$, and

the bipartition entanglement entropy follows from Peschel’s formula [5]:

$$S_{\text{ent}} = \sum_k [(\nu_k + \frac{1}{2}) \ln(\nu_k + \frac{1}{2}) - (\nu_k - \frac{1}{2}) \ln(\nu_k - \frac{1}{2})]. \quad (6)$$

2.3 Comparison metrics

Three diagnostic ratios are computed and compared to the prediction (2):

$$R_{\text{total}} = S_{\text{ent}}/S_{\text{esc}}, \quad R_{\Delta} = \Delta S/S_{\text{esc}}, \quad \Delta S \equiv S_{\text{ent}} - S_{\text{vac}}, \quad (7)$$

where S_{vac} is the entanglement entropy at the same N with no localized masses. If the postulate (2) holds with an order-unity prefactor, the diagnostic ratios should be approximately constant of order unity across the scan. Mutual information $I(M_1:M_2)$ and modular content ΔK are introduced in Sections V and VI.

2.4 Lattice setup (3+1D)

The 3+1D setup uses a cubic lattice of N^3 sites with Dirichlet boundary conditions on all six faces. Two point masses are placed on the central x -axis at $(i, N/2, N/2)$ with $i = (N \pm L)/2$, the bipartition cutting the cube at $i = N/2$. The K -matrix has shape $N^3 \times N^3$; for $N = 20$ this is 8000×8000 . The scan covers $N \in \{10, 12, 14, 16, 18, 20\}$, $L \in \{2, 4, 6, 8\}$, and $m \in \{0.1, 1.0, 10.0\}$. Two independent computational paths were used—a CPU implementation (NumPy/OpenBLAS, 24 cores) and a GPU implementation (`torch.linalg.eigh`, NVIDIA RTX 5090)—agreeing on ΔS at the $N = 14$ anchor to better than 2×10^{-5} .

3 Validations

Three validations against published results were performed before any scan was treated as established.

3.1 Cardy–Calabrese central charge (V1)

In the absence of localized masses, the bipartition entanglement entropy of a half-line on a Dirichlet-BC lattice obeys [7] $S_{\text{vac}}(N/2) \simeq (c/6) \ln(2N/\pi) + \text{const}$, with $c = 1$ for a free massless scalar. The slope from successive doublings converges to $(1/6) \ln 2 \simeq 0.11552$, matching the predicted central charge to better than 0.1% by $N = 1600$ (Table I); the non-universal additive constant $\simeq -0.030$ is the expected lattice-regularization piece.

N	S_{vac} (measured)	$(1/6) \ln(2N/\pi)$	Difference
50	0.55029	0.57674	−0.02645
100	0.66413	0.69226	−0.02814
200	0.77881	0.80779	−0.02897
400	0.89392	0.92331	−0.02939
800	1.00924	1.03884	−0.02960
1600	1.12466	1.15436	−0.02971

Table I. Vacuum bipartition entropy vs. the Calabrese–Cardy prediction ($c = 1$).

3.2 Heavy-mass Dirichlet limit (V2)

As $m \rightarrow \infty$ the localized-mass sites become perfect Dirichlet walls, and Calabrese–Cardy predicts $S_{\text{ent}}(m \rightarrow \infty) \rightarrow (1/6) \ln(2L/\pi) + \text{const.}$ For $L = 20$: predicted $S \rightarrow 0.394$; measured at $m = 100$: $S = 0.394$, agreement to four decimal places—an internal cross-check that the localized-mass machinery and the Peschel reduction agree at the analytically tractable limit.

3.3 Symbolic dimensional consistency (V3)

The escrow prediction (2) was derived symbolically (sympy) and verified to be the unique dimensionally-consistent form after cancellation of G_{eff} and the source mass. Per the lessons of [3], dimensional consistency is necessary but not sufficient; the substantive checks are V1 and V2.

3.4 3+1D area law (V4)

In 3+1D the vacuum entanglement entropy across a planar cut obeys the Bombelli–Srednicki area law [8,9] $S_{\text{vac}} \simeq a N^2 + \text{subleading}$, $a \simeq 0.022$. The measured coefficient S_{vac}/N^2 converges monotonically from below toward $\simeq 0.0228$ (Table II), consistent with published values.

N	S_{vac}	S_{vac}/N^2
10	2.140	0.0214
12	3.139	0.0218
14	4.331	0.0221
16	5.708	0.0223
18	7.290	0.0225
20	9.080	0.0227

Table II. Area-law coefficient S_{vac}/N^2 in 3+1D, converging toward $\simeq 0.0228$.

4 Result: bipartition entanglement entropy in 1+1D

4.1 Quantitative summary

Across the 295-point (N, L, m) scan in 1+1D: (i) the ratio R_{total} ranges from 2.3×10^{-9} to 83.3 at $N = 1000$ —a span of 10.56 orders of magnitude, with geometric mean $|R_{\text{total}}| \approx 8 \times 10^{-3}$; (ii) the mass-induced contribution $\Delta S = S_{\text{ent}} - S_{\text{vac}}$ is uniformly negative (localized masses *reduce* entanglement); (iii) results are converged with respect to lattice size to four decimal places for $m \geq 1$, so the scatter is not a finite-size artifact.

L	m	$N = 100$	$N = 300$	$N = 1000$	$N = 3000$
4	0.01	2.6388	3.3549	4.1245	4.7797
4	0.1	0.2350	0.2678	0.2827	0.2864
4	1.0	0.00737	0.00737	0.00737	0.00737
4	10	0.000495	0.000495	0.000495	0.000495
10	1.0	0.00488	0.00488	0.00488	0.00488
40	1.0	0.00206	0.00206	0.00206	0.00206
160	1.0	—	0.000739	0.000739	0.000739

Table III. R_{total} at fixed (L, m) vs. N in 1+1D. For $m \geq 1$ the ratio is N -independent to four decimals already at $N = 100$: the result is converged.

$L \backslash m$	0.001	0.01	0.1	1	10	100
2	83.3	8.25	0.563	0.00881	1.07×10^{-5}	2.27×10^{-9}
4	41.6	4.12	0.283	0.00737	4.95×10^{-4}	4.92×10^{-5}
10	16.7	1.65	0.115	0.00488	4.43×10^{-4}	4.43×10^{-5}
40	4.16	0.413	0.0301	0.00206	2.03×10^{-4}	2.03×10^{-5}
160	1.04	0.103	0.00844	0.000739	7.37×10^{-5}	7.37×10^{-6}
640	0.26	0.026	0.00249	0.000242	2.42×10^{-5}	2.42×10^{-6}

Table IV. R_{total} across the (L, m) grid at $N = 1000$, spanning 10.56 orders of magnitude; no rescaling recovers order-unity agreement.

4.2 Why the literal postulate fails: the structural mismatch

The escrow prediction (2) grows linearly with both m and L without bound. The free-scalar entanglement entropy behaves very differently: the vacuum part S_{vac} scales as $(1/6) \ln(2N/\pi)$ (bounded for fixed N), and the mass-induced part ΔS is bounded above by $O(1)$ nats—localized masses entangle or disentangle modes only within a correlation length $\xi \sim 1/m$, giving an $O(1)$ contribution that saturates as $L \rightarrow \infty$. Heavy masses act as effective Dirichlet walls, decreasing the bipartition entanglement and giving $\Delta S < 0$. The mismatch is structural, not a missing prefactor: S_{esc} is extensive ($\propto mL$) while ΔS is intensive and bounded. No constant rescaling or modified temperature can reconcile a bounded quantity with an unbounded one.

4.3 Robustness against UV artifacts

The result is invariant under the mass implementation. Single-site delta masses and Gaussian-smear profiles with widths $\sigma \in \{1, 2, 5\}$ all give the same qualitative behavior: ΔS uniformly negative, R_{total} varying across many orders of magnitude. For $m \leq 0.1$, ΔS is invariant to smearing to better than 1%; for $m \gtrsim 1$, smearing makes ΔS more negative (a broader “wall” removes more entanglement). Single-site UV artifacts are not driving the result.

5 Independent confirmations

5.1 Mutual information $I(M_1:M_2)$

A natural alternative measure is the mutual information between regions surrounding the two masses,

$$I(M_1:M_2) = S(M_1) + S(M_2) - S(M_1 \cup M_2). \quad (8)$$

For free massless scalars in 1+1D the leading contribution at large separation scales as [10] $I(M_1:M_2) \sim (w/L)^4$ with w the region width. We measured it for $w = 4$ and $L \in \{10, 20, 40, 80, 160, 320\}$: the mutual information decays as L^{-4} , while the postulate predicts growth $\propto 2\pi mL$. The ratio R_{MI} thus spans more than 12 orders of magnitude in the opposite direction to what the postulate requires—an independent confirmation.

5.2 3+1D continuum-limit study

The 3+1D bipartition entanglement entropy was computed across $N \in \{10, \dots, 20\}$; the vacuum area-law validation (Table II) confirmed the implementation. The diagnostic ratio R_{total} has no finite continuum limit—it grows as N^2 across the entire (L, m) grid, with

$$R_{\text{total}} \cdot S_{\text{esc}} = S_{\text{vac}} \quad [\text{to better than 1\% at every point; residual is the } R_{\Delta} \lesssim 10^{-3} \text{ mass effect}]. \quad (9)$$

The reason is transparent: the 3+1D bipartition entropy is dominated by the area-law vacuum contribution $S_{\text{vac}} \propto N^2$, unrelated to the mass binding, while $S_{\text{esc}} = 2\pi mL$ is N -independent.

L	m	$N = 10$	$N = 12$	$N = 14$	$N = 16$	$N = 18$	$N = 20$ (R_{Δ})
2	0.1	1.71	2.50	3.45	4.55	5.81	7.22 (-1×10^{-4})
2	1.0	0.171	0.249	0.345	0.455	0.581	0.722 (-8×10^{-4})
4	1.0	0.0855	0.125	0.173	0.228	0.291	0.361 (-2×10^{-4})

Table V. R_{total} in 3+1D grows as N^2 (area-law driven), no finite continuum limit. The mass-induced ratio R_{Δ} is bounded by 10^{-3} , three orders below the postulate’s prediction $R_{\Delta} = 1$.

The 3+1D failure is structurally distinct from the 1+1D one: in 1+1D R_{total} is bounded but spans 10.56 orders of magnitude because S_{ent} saturates while S_{esc} grows; in 3+1D R_{total} has no continuum limit at all because the bipartition entropy is dominated by the cut-area vacuum term.

6 Modular Hamiltonian content: a suggestive window in 1+1D

6.1 Bisognano–Wichmann motivation

The Bisognano–Wichmann theorem [6] states that the modular Hamiltonian K_A of a Lorentz-invariant QFT restricted to a half-space is, in vacuum,

$$K_{\text{BW}} = 2\pi \int x_{\perp} T_{00}(x) d^d x, \quad (10)$$

with $x_{\perp}$ the perpendicular distance from the entangling surface. For a localized energy m at distance L , the linear-response prediction is $\Delta K_{\text{BW}} \simeq 2\pi mL$, exactly the form of the escrow prediction (2). This structural match raises the possibility that the postulate’s correct identification is with modular Hamiltonian content rather than entanglement entropy.

6.2 Lattice implementation

We computed $\Delta K = \langle K_A^{\text{vac}} \rangle_{\text{pert}} - \langle K_A^{\text{vac}} \rangle_{\text{vac}}$ across the same 1+1D grid. Using the Williamson normal form of the vacuum covariance, with modular eigenvalues $\varepsilon_k = \ln[(\nu_k + \frac{1}{2})/(\nu_k - \frac{1}{2})]$, the quadratic modular kernels are $Q = M_{\phi}^{-1/2} V \text{diag}(\varepsilon_k \nu_k) V^{\text{T}} M_{\phi}^{-1/2}$ and $P = M_{\phi}^{1/2} V \text{diag}(\varepsilon_k / \nu_k) V^{\text{T}} M_{\phi}^{1/2}$, so that $\langle K_A \rangle = \frac{1}{2} \text{tr}(QM'_{\phi}) + \frac{1}{2} \text{tr}(PM'_{\pi})$. The implementation was validated by three checks: relative-entropy positivity $S_{\text{rel}} = \Delta K + S_{\text{vac}} - S_{\text{pert}} \geq 0$ for all states; a modal-vs-original-basis cross-check agreeing to six decimals; and constant-offset consistency with $\langle K_A \rangle_{\text{vac}} = S_{\text{vac}}$.

6.3 Result: right form, wrong magnitude

In the $m \approx 1$ window the lattice ΔK reproduces the BW structural shape (a non-perturbative feature, opposite to the negative ΔK of small- m linear response): it is positive, grows with L , and is approximately linear in m for moderate values. Two discrepancies emerge. First, the L -scaling is not a single power law. A sliding-window analysis over d_1 (the perpendicular distance of the mass in A to the cut, $= L/2$ in the symmetric placement) shows the local exponent evolving continuously from $\Delta K \propto d_1^{1.02 \pm 0.05}$ at small $d_1 \in [2, 6]$ (indistinguishable from the BW linear asymptote) down to $\Delta K \propto d_1^{0.5 \pm 0.04}$ in the decay tail $d_1 \in [16, 64]$. The previously-published single exponent “0.7” was an artifact of fitting one power law across a smooth crossover, and is retracted as a universal description.

Second, the magnitude is smaller than the literal BW prediction—though part of the gap is a normalization convention: the $\frac{1}{2}m^2\phi_i^2$ site perturbation inserts ground-state energy $\Delta E \approx 0.256$ at $m = 1$, and normalizing ΔK to ΔE rather than to m raises the prefactor from $\approx 1/30$ to $\approx 1/7$. In the small- d_1 BW regime ($d_1 \in [2, 6]$, $m = 1$, $N = 1000$), the prefactor $\Delta K/(2\pi m d_1)$ is ≈ 0.030 – 0.037 , a suppression of roughly $1/30$ relative to the literal BW value. At heavier mass the apparent ratio drops further, as ΔK saturates rather than continuing to grow as $2\pi m d_1$; the positive, approximately linear-in- d_1 window we report sits at $m \lesssim 1$ and $d_1 \in [2, 8]$. This window is an empirical non-perturbative feature, not the small-mass linear-response regime—in genuine linear response ($m \ll 1$), ΔK is negative and d_1 -flat.

6.4 Interpretation

The 1+1D modular result is consistent with the BW asymptote at small d_1 with a suppressed prefactor. The smooth crossover into a sublinear tail at $d_1 \gtrsim 8$ reflects geometric corrections to the BW half-space assumption (Dirichlet rather than free boundary conditions; finite subsystem- A depth), consistent with a finite-size deformation of BW rather than a fundamental failure. A substantial part of the prefactor is a normalization convention (energy-normalizing gives $\approx 1/7$); whether the residual suppression has a calculable origin is left for future work. We emphasize that this is a suggestive-but-inconclusive result: it is an empirical non-perturbative feature at $m = 1$, *not* a controlled linear-response recovery of BW—in the linear-response regime ΔK is negative and d_1 -flat.

A candidate explanation worth flagging is a UV/IR mismatch: the BW formula is an infrared prediction valid in the continuum, whereas a lattice mass with $m \approx 1$ has correlation length $\xi \sim 1/m \sim 1$ lattice spacing—a UV-scale perturbation. The $1/30$ suppression may reflect this mismatch and could in principle be computed by evaluating the lattice modular Hamiltonian in the continuum scaling limit. A second candidate is that the Dirichlet boundary conditions used on the lattice differ from the free boundary conditions assumed in the standard BW derivation, which may contribute a calculable boundary-correction factor to the residual prefactor. The dimensional dependence is investigated in a companion paper [11], which finds the 3+1D BW asymptote is *not* recovered within the resolvable d_1 range: ΔK peaks at $d_1 \approx 2$ and decays as d_1^{-2} to -3 , with a matched-mass peak ratio ≈ 0.0026 , about $14\times$ smaller than the 1+1D peak ≈ 0.037 . The companion paper notes that the rate of approach to the continuum BW asymptote may genuinely be slower in 3+1D—where the conformal-symmetry group of the massless free scalar is finite-dimensional—than in 1+1D, where it is the infinite-dimensional Virasoro algebra; 3+1D BW recovery at much larger d_1 inaccessible to the present

lattice therefore cannot be strictly ruled out. We do not adjudicate among these readings; the result is reported as the most promising direction for future modification of the postulate and as a sharp constraint on what such modifications must satisfy.

7 Falsifiability check

A central concern is that any apparent agreement could be a tautology fixed by the choice of dimensional combinations. The Unruh temperature varies as $T_{\text{Unruh}} \propto m/L^2$, spanning five orders of magnitude across the scan. We verified that no choice of T produces a uniform $R \simeq 1$ across the scan; T_{Unruh} is not privileged in this sense. The 10.56 orders of magnitude variation in R is therefore a real measurement of the disagreement, not an artifact of dimensional rescaling. The Section IV result is a genuine quantitative measurement of the postulate’s failure as bipartition entanglement entropy.

8 What this settles, and what it does not

What this settles. The literal static escrow postulate (1), as a quantitative identification of S_{esc} with QFT bipartition entanglement entropy in flat space, does not survive in either 1+1D or 3+1D. In 1+1D the qualitative behavior is wrong (sign, scaling, boundedness in N), the magnitude is wrong (10.56 orders of scatter), and lattice-size convergence rules out finite-size explanations. In 3+1D the bipartition entropy is dominated by the area-law vacuum term and has no finite continuum limit; the mass-induced ratio R_Δ is bounded by 10^{-3} . Two code paths agree to five decimals, and mutual information confirms the failure from a third direction.

What this does not settle. The horizon limit is unaffected: the recovery of Bekenstein–Hawking entropy [1, §5] goes through surface gravity and the Unruh temperature at the horizon directly, not through volume-bipartition entanglement. Curved-spacetime regimes are outside scope. The framework’s empirical content—the constant- a_0 SPARC reanalysis, the deep-MOND asymptote, and the baryonic Tully–Fisher relation—does not depend on the literal bipartition-entanglement reading. Modular-content variants remain partially open: the 1+1D window at $m = 1$ (not a linear-response recovery; ΔK is negative and flat in linear response) with prefactor $\approx 1/30$, and the companion paper’s 3+1D non-recovery, show the modular-content interpretation has dimension-dependent validity within currently accessible lattice sizes.

9 Methodological notes

The calculations were performed via autonomous runs on local hardware (NVIDIA RTX 5090, Intel Core Ultra 9, 256 GB DDR5) with full code execution. Three features are worth highlighting. First, validations against published results (Calabrese–Cardy, Bombelli–Srednicki), not internal consistency checks alone, are the binding constraint—agreement to better than 0.1% (1+1D) and a few percent (3+1D) confirms the implementations. Second, the falsifiability check (Section VII) is load-bearing: without verifying that T_{Unruh} is privileged, the result could be dismissed as a tautology. Third, modular-content code requires special validation: the ΔK computation passed relative-entropy positivity, modal-vs-original-basis agreement to six decimals, and constant-offset consistency; an initial implementation produced order- 10^4 relative-entropy violations that immediately flagged an error in the Williamson decomposition. The 3+1D study was run on two independent paths (CPU NumPy, ~ 48 min; GPU

PyTorch, <3 min) agreeing on ΔS to better than 2×10^{-5} , ruling out implementation and round-off artifacts.

10 Conclusion

The literal static escrow postulate of [1], interpreted as a quantitative identification of $S_{\text{esc}} = |U_{\text{grav}}|/T_{\text{Unruh}}$ with QFT bipartition entanglement entropy, fails in both 1+1D and 3+1D. In 1+1D the failure is qualitative (wrong sign, Dirichlet-wall behavior), quantitative (10.56 orders of magnitude in R), and robust; in 3+1D it is structurally sharper— R_{total} has no finite continuum limit because the bipartition entropy is dominated by the Bombelli–Srednicki area law, and R_{Δ} is bounded by 10^{-3} . Two code paths agree to five decimals, and mutual information confirms the failure independently.

One result is positive within a finite regime: the modular Hamiltonian content ΔK , under the Bisognano–Wichmann conjecture, shows a positive, approximately linear window ($\Delta K \propto d_1$) at $m = 1$ —resembling but not controllably recovering the BW asymptote (in linear response ΔK is negative and d_1 -flat)—in a small- d_1 window, with a suppressed prefactor $\approx 1/30$ of the literal BW value ($\approx 1/7$ energy-normalized). The published “sublinear scaling at exponent 0.7” framing is corrected to a regime-dependent characterization. The companion paper [11] reports that the analogous 3+1D test does not recover BW within the resolvable d_1 range. The published framework’s interpretive content and empirical results are not refuted; what is ruled out is the literal bipartition-entanglement reading of its central postulate. Future tests should target the modular-content reinterpretation, the horizon limit directly, or curved-spacetime regimes—including 2+1D as an intermediate case and $N \gtrsim 20$ GPU runs in 3+1D to test whether BW recovery is reached at larger d_1 .

Acknowledgments

LLM-assisted code review and cross-comparison of the lattice run output were performed using Anthropic Claude Opus 4.7, xAI Grok, and Perplexity Deep Research; these tools were used for code review and consistency checks, not as substitutes for peer review. The substantive cross-validation reported here is the agreement to better than five decimal places between two independent code paths (CPU NumPy/OpenBLAS and GPU PyTorch/CUDA) on different hardware (§IX). The author is solely responsible for the conclusions and any remaining errors.

Data and code availability

Lattice scripts, validation code, and parameter-scan results were produced for this study. The Zenodo deposit (record 20057538) currently contains the manuscript, a deposit README, a consolidated-findings document, and the self-contained 1+1D script `lattice_1d_modular.py`, which reproduces the §VI.C modular-Hamiltonian claims; the 3+1D code, mutual-information code, full parameter-scan results, and diagnostic plots are not yet deposited and will be added in a follow-up archive. A clean rerun of the 1+1D pipeline takes ≈ 7 min on the reference hardware.

Data, Code & Resources

Manuscript source and archived record: <https://github.com/Windstorm-Institute/lattice-qft-test> (Zenodo: 10.5281/zenodo.20057537). Experiments, datasets, and analysis code:

<https://github.com/Windstorm-Labs/lattice-qft-test>. Windstorm Institute: <https://windstorminstitute.org>. Windstorm Labs: <https://windstormlabs.com>.

References

- [1] G. L. Whitmer, *Gravitational Entropy Escrow: An Interpretive Synthesis*, Zenodo (2026). DOI:10.5281/zenodo.20031931.
- [2] E. P. Verlinde, *On the origin of gravity and the laws of Newton*, JHEP **04**, 029 (2011). arXiv:1001.0785.
- [3] G. L. Whitmer, *On the Status of Local Entropy-Current Extensions of the Gravitational Entropy Escrow Framework*, Zenodo (2026). DOI:10.5281/zenodo.20041991.
- [4] J. D. Bekenstein, *Universal upper bound on the entropy-to-energy ratio for bounded systems*, Phys. Rev. D **23**, 287 (1981).
- [5] I. Peschel, *Calculation of reduced density matrices from correlation functions*, J. Phys. A **36**, L205 (2003).
- [6] J. J. Bisognano and E. H. Wichmann, *On the duality condition for quantum fields*, J. Math. Phys. **17**, 303 (1976).
- [7] P. Calabrese and J. Cardy, *Entanglement entropy and quantum field theory*, J. Stat. Mech. P06002 (2004). hep-th/0405152.
- [8] L. Bombelli, R. K. Koul, J. Lee, and R. D. Sorkin, *Quantum source of entropy for black holes*, Phys. Rev. D **34**, 373 (1986).
- [9] M. Srednicki, *Entropy and area*, Phys. Rev. Lett. **71**, 666 (1993).
- [10] P. Calabrese, J. Cardy, and E. Tonni, *Entanglement entropy of two disjoint intervals in conformal field theory*, J. Stat. Mech. P11001 (2009). arXiv:0905.2069.
- [11] G. L. Whitmer, *Dimensional Dependence of Modular Hamiltonian Content in the Gravitational Entropy Escrow Framework: 3+1D Cubic-Lattice Test (companion paper)*, Zenodo (2026).