

Outline Draft

Loveless, Nix, Stokes, Abernathy's

June 10, 2026

1 Introduction

Alzheimer's Disease (AD) is characterized as a complex neurodegenerative disease affecting more than 55 million people globally [10]. Symptoms are caused by the degeneration of the central nervous system, and are manifested by progressive memory loss, difficulty with language, drastic mood swings, among other effects. It is a disease that causes an extensive burden to those suffering from the disease through disorientation, to relatives in the form of mental taxation, as well as economically, projecting an estimated global economic burden of 16.9 trillion dollars by 2050 [12]. These factors are just a few reasons as to the importance of research in this area, where many scientists have worked to understand the underlying mechanisms of AD. However, much is still unknown in how AD is caused, as well as what biological processes occur in the brain, making it difficult to come to a decisive conclusion regarding its cause.

Many studies have been conducted in regard to understanding Alzheimer's Disease. With this, there are a few main hypotheses that are generally considered to be factors in the causation and progression of AD [16]. The so-called *Amyloid Cascade Hypothesis* is among the most extensively studied in the literature, where it is proposed that a misfolding of β -amyloid (A_β) protein forms senile plaques. These plaques are hypothesized to be the main driving force in the intracellular deposition of misfolded τ proteins in neurofibrillary tangles (NFTs) and memory loss in the progression of Alzheimer's Disease [7]. Growing evidence, however, shows that this hypothesis is suggested to be but a small factor in the larger picture of the underlying operative mechanisms of the disease [8]. This has led researchers to branch into other pathways concerning AD, looking at the interactions between A_β and τ in an attempt to further understand the pathogenesis of these two factors [15]. Some studies have also observed the role of calcium in this process, further suggesting a more complex process that is occurring biologically. This occurs through a feedback interaction between an upregulation of Ca^{2+} and the amyloidogenic pathway, remodeling the Ca^{2+} signaling pathway [4].

The various complex interactions in AD have shown to be an obstacle to yet overcome when modeling the behavior of the disease. Numerous different studies have in turn given an attempt at creating mathematical models observing different aspects of the disease in order to observe the underlying dynamics. This has led to many models being introduced, each aiming to understand various components in Alzheimer's Disease through different methodologies. Compartments concerning the interaction between A_β and τ are studied, observing the suspected effects on neurodegeneration and cognitive decline through a mechanistic causal model [14, 6]. The feedback loop between A_β and Ca^{2+} has also been analyzed using a

deterministic system, adding a noise-inducing element to the system [13, 17]. Models that observe lower-level processes have also been studied, lending to valuable insight as to what is suggested to be seen on the cellular level [11, 9].

It should be noted that in these models, while each has its own way of observing some of the underlying mechanisms, there are still potential shortfalls when it comes to finding effective treatments. Evidence suggests that focusing solely on anti- A_β treatments or τ protein treatments, rather than attempting to determine treatments for many different aspects of the disease [1, 15]. This provides difficulty in understanding exactly the importance of which component of the disease should be the focus of further study in order to provide evidence for treatments having higher efficacy. In this study, a mechanistic model is proposed, analyzing the interaction between A_β , Ca^{2+} , and τ proteins; observing the effects of this multifaceted interaction on neurodegeneration, and cognitive decline. Sensitivity analysis is performed in order to better understand the importance of each aspect of the model, and optimal control is used in order to determine the suggested avenues of effective treatment.

2 Proposed Model

To understand the role calcium may have in the interaction of AD, a model is proposed using five biomarkers found to have interactions within the disease: hallmark biomarkers through amyloid beta and tau, an interactional biomarker through calcium, suggested to have a role in amyloid beta and tau protein accumulation. Also observed are neuronal loss biomarkers, and cognitive decline. The pathology of AD is suggested to start through a feedback loop between amyloid beta and calcium through a dysregulation of calcium; this exacerbates the aggregation of amyloid beta, forming plaques. These plaques then promotes the hyperphosphorylation of tau proteins, leading to neurodegeneration and subsequently cognitive decline.

2.1 Amyloid Beta and Calcium Equations

Because amyloid beta and calcium are suggested to interact in such a way that a dysregulation of calcium causes an acceleration of amyloid beta accumulation [13], this is modeled mathematically as

$$\frac{dA_\beta}{dt} = a_1 + a_2 \frac{Ca}{Ca + \sigma} - k_1 A_\beta - u_1 A_\beta \quad \text{with} \quad A_\beta(T_0) = A_0, \quad (1)$$

where A_0 is the initial condition of A_β at age T_0 , and can vary on a per patient basis [6]. The rate of A_β growth increases at a linear rate a_1 and decreases proportionally to the population of A_β , as described by $k_1 A_\beta$. Here, $a_2 \frac{Ca}{Ca + \sigma}$ acts as the interaction of calcium on A_β using a Hill equation of $n = 1$. The removal rate can be modified by choosing different values for u_1 which represents the effect of treatment targeting A_β . The term u_1 can either be constant or be a function of time.

Similarly, an A_β interaction is observed to have a role in the further dysregulation of calcium, and the equation for calcium is defined as

$$\frac{dCa}{dt} = b_1 + b_2 A_\beta - k_2 Ca - u_2 Ca \quad \text{with} \quad Ca(T_0) = Ca_0. \quad (2)$$

Here, calcium grows at a linear rate b_1 , and the accumulation of A_β is positive at a rate b_2 . There is also a natural removal rate of calcium, k_2 . Notice that there is also the option to administer treatment targeting calcium through the term u_2 .

2.2 Tau Equation

Many studies have suggested that the pathology of tau is initiated by amyloid beta accumulation, and have modeled this process mathematically [18, 6]. However, it has also been suggested that calcium dysregulation also has an effect inducing the production of tau proteins [5]. This leads to a definition of the tau equation to be given by

$$\frac{d\tau}{dt} = c_1 + c_2 A_\beta + c_3 Ca - k_3 \tau - u_3 \tau \quad \text{with} \quad \tau(T_0) = \tau_0. \quad (3)$$

It is noted here that only pathological tau is observed, where other studies have used compartments relating to non-amyloid dependent tau accumulation, or suspected non-Alzheimer pathology (SNAP) [14, 6]. This function is similar to Equation (1) and Equation (2) in that $\frac{d\tau}{dt}$ grows at a linear rate d_1 and decreases proportional to the population of τ through $k_3 \tau$. There is also the treatment term $u_3 \tau$.

2.3 Neurodegeneration Equation

When tau is in a pathological state, it no longer binds to microtubules as intended; instead, the formation of NFTs occur in the neurons, leading to neurodegeneration [2]. Hence, this equation is defined as

$$\frac{dN}{dt} = d_1 + d_2 \tau - k_4 N.$$

This equation follows the established structure and also includes the interaction term $d_2 \tau$.

2.4 Cognitive Decline Equation

Both tau and neurodegeneration are evidenced to directly affect cognitive decline [3, 6]. Because of this, the equation for cognitive decline is defined as

$$\frac{dC}{dt} = e_1 + e_2 NR + e_3 \tau - k_5 C.$$

The parameter R represents a patient having the APOE $\epsilon 4$ gene, which has been shown to be a risk factor in Alzheimer's disease (need to find citation of this).

This model is summarized as a system of ordinary system of differential equations with

$$\begin{cases} \frac{dA_\beta}{dt} = a_1 + a_2 \frac{Ca}{Ca+\sigma} - k_1 A_\beta - u_1 A_\beta \\ \frac{dCa}{dt} = b_1 + b_2 A_\beta - k_2 Ca - u_2 Ca \\ \frac{d\tau}{dt} = c_1 + c_2 A_\beta + c_3 Ca - k_3 \tau - u_3 \tau \\ \frac{dN}{dt} = d_1 + d_2 \tau - k_4 N \\ \frac{dC}{dt} = e_1 + e_2 NR + e_3 \tau - k_5 C \end{cases} \quad \text{with} \quad \begin{cases} A_\beta(T_0) = A_0 \\ Ca(T_0) = Ca_0 \\ \tau(T_0) = \tau_0 \\ N(T_0) = N_0 \\ C(T_0) = C_0. \end{cases} \quad (4)$$

3 Analysis

3.1 Preliminary Results

Assuming (4) is subject to initial conditions that are non-negative, it is noted that the system is invariant in the non-negative orthant. Let $r_1 = k_1 + u_1$ and $r_2 = k_2 + u_2$ for the simplicity of calculations. Considering the components uncoupled from the rest of the proposed model, $dA_\beta/dt, dCa/dt$ can be shown to be bounded. This is done by first noticing

$$a_1 + a_2 \frac{Ca}{Ca + \sigma} - r_1 A_\beta \leq a_1 + a_2 - r_1 A_\beta$$

since $Ca/(Ca + \sigma) \in [0, 1)$. So, $a_1 + a_2 - r_1 A_\beta < 0$ when $A_\beta > \frac{a_1 + a_2}{r_1}$. Similarly, using that A_β is bounded, it follows that

$$\frac{dCa}{dt} = b_1 + b_2 A_\beta - r_2 Ca \leq b_1 + b_2 M - r_2 Ca$$

as $M > A_\beta(t)$ for all $t \geq 0$. So, $b_1 + b_2 M - r_2 Ca < 0$ when $Ca > \frac{b_1 + b_2 M}{r_2}$, thus Ca is also bounded. The rest of the system follows from these two variables, so the rest of the system is bounded as well.

Lemma 1 (Establishment of Unique Equilibria of Uncoupled System). *There exists a unique equilibria (A_β^*, Ca^*) for the system*

$$\begin{cases} \frac{dA_\beta}{dt} = a_1 + a_2 \frac{Ca}{Ca + \sigma} - r_1 A_\beta \\ \frac{dCa}{dt} = b_1 + b_2 A_\beta - r_2 Ca. \end{cases} \quad (5)$$

Proof. To establish the existence of equilibria for Equation (5), a solution to the following must be obtained:

$$\begin{aligned} a_1 + a_2 \frac{Ca}{Ca + \sigma} - r_1 A_\beta &= 0 \\ b_1 + b_2 A_\beta - r_2 Ca &= 0. \end{aligned}$$

By solving $\frac{dA_\beta}{dt} = a_1 + \frac{a_2 Ca}{Ca + \sigma} - r_1 A_\beta$ for A_β , we get

$$A_\beta = \frac{a_1 Ca + a_2 Ca + a_1 \sigma}{r_1 (Ca + \sigma)}.$$

We then substitute this value into $\frac{dCa}{dt} = b_1 + b_2 A_\beta - r_2 Ca$. The resulting expression is,

$$b_1 - r_2 Ca + \frac{b_2 (a_1 Ca + a_2 Ca + a_1 \sigma)}{r_1 (Ca + \sigma)}. \quad (6)$$

Setting (6) equal to 0 and doing further simplification yields the quadratic,

$$-(r_1 r_2) Ca^2 + (a_1 b_2 + a_2 b_2 + b_1 r_1 - r_1 r_2 \sigma) Ca + a_1 b_2 \sigma + b_1 r_1 \sigma = 0. \quad (7)$$

Notice that in (7) the quadratic term is always negative and the constant term is always positive. The linear term can either be positive or negative depending on the choice of parameters. Regardless of the parity of the linear term, there is only one sign change in the expression, so by Descartes' Rule of Signs, there is one positive real root to this quadratic. Thus, there exists a unique positive equilibrium (A_β^*, Ca^*) for the system in Equation (5). $\square$

Lemma 2 (Global Stability of Equilibrium). *The equilibrium (A_β^*, Ca^*) is globally asymptotically stable.*

Proof. Let $\mathbf{F}(A_\beta, Ca) = \langle a_1 + \frac{a_2 Ca}{Ca + \sigma} - r_1 A_\beta, b_1 + b_2 A_\beta - r_2 Ca \rangle$. Then, $\nabla \cdot \mathbf{F} = -r_1 - r_2$. Choose the function $\alpha(A_\beta, Ca) = 1$. So, $\nabla \cdot (\alpha \mathbf{F}) < 0$, therefore by the Bendixson-Dulac Theorem, there is no limit cycle in the system of ODEs. So, the equilibrium point must be globally stable in the non-negative orthant. $\square$

Theorem 1 (Global Stability of Equilibrium of System). *There exists a unique globally asymptotically stable equilibrium $(A_\beta^*, Ca^*, \tau^*, N^*, C^*)$.*

Proof. By Lemma 2, there exists a unique globally asymptotic stable equilibrium (A_β^*, Ca^*) . By solving (3) for τ we get the following equilibrium value:

$$\tau^* = \frac{c_1 + c_2 A_\beta^* + c_3 Ca^*}{k_3 + u_3}.$$

By similar methods we find

$$N^* = \frac{d_1 + d_2 \tau^*}{k_4}$$

and

$$C^* = \frac{e_1 + e_2 N^* R + e_3 \tau^*}{k_5}.$$

Thus there is a unique globally asymptotic stable equilibrium solution for the system in Equation (4). $\square$

4 Parameters

params in Table 1. need elaboration and characterizations

5 Sensitivity Analysis

Sobol sensitivity analysis is used to provide a quantitative assessment of the impact model parameters have on the causal dynamics of the proposed model. The first, second, and total-order indices (S_T) were computed using different sample sizes, ranging from $N = 2^4$ to $N = 2^{20}$. Using the parameter ranges from Table (blank), the first-order and total-order sensitivity indices are highlighted in Figure (blank); initial conditions were set for $Ca(0) = 200$, and zero for the rest of the state variables. In order to validate these indices, Figure (blank) shows the six most sensitive parameters for each sample size. The total-order indices are quantified along with the confidence interval for each parameter at $N = 2^{20}$ in Table (blank).

Var	Value	Unit	Source
A_0	0	μM	
Ca_0	100	nM	
τ_0	0	μM	
N_0	0	billions neurons lost	
C_0	0	MMSE point decline	
a_1	6.5641	$\mu M/yr$	
a_2	1.5778	$\mu M/yr$	
b_1	315569260	nM/yr	
b_2	6311385.2	yr^{-1} (100)	
c_1	52.2958	$\mu M/yr$	
c_2	1.78367	yr^{-1}	
c_3	0.1	yr^{-1} (1/100)	
d_1	0.0716	billions neurons lost/ yr	
d_2	0.398406	billions neurons lost/ $(yr \cdot \mu M)$	
e_1	0.01463	MMSE decline/ yr	
e_2	0.008684	MMSE decline/ $(yr \cdot \text{billions neurons lost})$	
e_3	0.01992	MMSE decline/ $(yr \cdot \mu M)$	
k_1	0.303598	yr^{-1}	
k_2	3155692.6	yr^{-1}	
k_3	1.8333	yr^{-1}	
k_4	0.3588	yr^{-1}	
k_5	0.08684	yr^{-1}	
u_1	varies	yr^{-1}	
u_2	varies	yr^{-1}	
u_3	varies	yr^{-1}	
σ	50 – 150	nM	

Table 1: Parameters used for simulation

6 Results

Using the derived biological parameters in Table 1, we run the simulation forward for 50 years with the initial conditions $(A_0, Ca_0, \tau_0, N_0, C_0) = (0, 100, 0, 0, 0)$. This simulation uses `solve_ivp` from the SciPy package in Python. The results from the simulation are shown in Figure 1. Initially, A_β and Ca increase and drive each other's growth. This leads to a large growth of τ proteins, thus leading to neuron loss over time, and cognitive decline over an even longer time scale. The biomarker populations reach their equilibrium values relatively early, but it takes cognitive decline many years to reach its equilibrium value.

A phase portrait (Figure 2) shows how different trajectories from varying initial conditions traverse the $A_\beta - Ca$ space. Trajectories will primarily move toward the Ca nullcline with slight movement along the horizontal axis. This is because the time-scale of Ca is much smaller than that of A_β . Once trajectories are near this nullcline, they travel directly toward the equilibrium solution.




Figure 1: ODE system simulation




Figure 2: Phase portrait




Figure 3: Treatment effect on C^*

To determine which proteins are most effective to treat, we graphed the equilibrium cognitive decline value against varying treatments against each protein, as seen in Figure 3. Each u_n term increases such that the behavior of the function can be observed. We see that the variable that has the largest impact on C^* is u_3 , the treatment that reduces the number of τ proteins. The next most effective treatment is u_1 , which reduces the concentration of A_β plaques in the brain. We observe that treating the average level of Ca does not have a large effect on the equilibrium cognitive decline.

7 Optimal Control

Optimal control is a method we can use to determine the optimal values of u_n while tracking some cost function. In optimal control, we will let each u_n be a function of time, $u_n(t)$, as it is not realistic to maintain a constant treatment over a long time-frame. Tracking a cost function is useful for our application as some drugs used to treat Alzheimer's Disease have been shown to induce strong side effects such as brain swelling and brain bleeding (citation needed). So, minimizing this cost function while maximizing treatments will let the medication have its strongest impact without crossing some side-effect threshold.

7.1 Objective Function

The following equation is our objective function which tracks our cost over the time horizon of the treatment.

$$J(u_1, u_2, u_3) = \int_0^K A_1 N + A_2 C + A_3 u_1^2 + A_4 u_2^2 + A_5 u_3^2 dt. \quad (8)$$

The goal of treatment is to minimize the equilibrium values N^* and C^* while at the same time minimizing the side effects from the drugs. The terms A_n are the weights which determine how important each associated variable is in the optimization problem. The variables u_n are the treatment inputs, subject to the following constraints:

$$u = \{(u_1, u_2, u_3) \in L^\infty(0, K)^3 \mid (u_1, u_2, u_3) \in [0, u_1^{\max}], [0, u_2^{\max}], [0, u_3^{\max}] \forall t \in [0, K]\}$$

Each term u_n must be non-negative during the duration of the treatment.

7.2 Hamiltonian

$$\begin{aligned}\mathcal{H} = & A_1 N + A_2 C + A_3 u_1^2 + A_4 u_2^2 + A_5 u_3^2 + \lambda_1 \left(a_1 + a_2 \frac{Ca}{Ca + \sigma} - k_1 A_\beta - u_1 A_\beta \right) \\ & + \lambda_2 (b_1 + b_2 A_\beta - k_2 Ca - u_2 Ca) + \lambda_3 (c_1 + c_2 A_\beta + c_3 Ca - k_3 \tau - u_3 \tau) \\ & + \lambda_4 (d_1 + d_2 \tau - k_4 N) + \lambda_5 (e_1 + e_2 NR + e_3 \tau - k_5 C).\end{aligned}$$

The Hamiltonian includes the cost function as described in Equation (8) as well as the co-state vectors for each variable present in our system. Thus, we seek an optimal control set (u_1^*, u_2^*, u_3^*) such that

$$J(u_1^*, u_2^*, u_3^*) = \min \{ J(u_1, u_2, u_3) \mid (u_1, u_2, u_3) \in \mathcal{U} \}.$$

more stuff to add here

Theorem 2 (Optimal Control). *There exists an optimal control set (u_1^*, u_2^*, u_3^*) that minimizes $J(u_1, u_2, u_3)$ over $\mathcal{U}$. Furthermore, there exists adjoint variables $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ and λ_5 that satisfy the following system of differential equations:*

$$\begin{cases} \frac{d\lambda_1}{dt} = \frac{-\partial \mathcal{H}}{\partial A_\beta} = k_1 \lambda_1 + u_1 \lambda_1 - b_2 \lambda_2 - c_2 \lambda_3 \\ \frac{d\lambda_2}{dt} = \frac{-\partial \mathcal{H}}{\partial Ca} = -\lambda_1 \frac{a_2 \sigma}{(Ca + \sigma)^2} + k_2 \lambda_2 + u_2 \lambda_2 - c_3 \lambda_3 \\ \frac{d\lambda_3}{dt} = \frac{-\partial \mathcal{H}}{\partial \tau} = k_3 \lambda_3 + u_3 \lambda_3 - d_2 \lambda_4 - e_3 \lambda_5 \\ \frac{d\lambda_4}{dt} = \frac{-\partial \mathcal{H}}{\partial N} = -A_1 + k_4 \lambda_4 - e_2 R \lambda_5 \\ \frac{d\lambda_5}{dt} = \frac{-\partial \mathcal{H}}{\partial C} = -A_2 + k_5 \lambda_5 \end{cases}$$

with transversality conditions $\lambda_1(K) = \lambda_2(K) = \lambda_3(K) = \lambda_4(K) = \lambda_5(K) = 0$. Also, the optimal control set has the characterization:

$$\begin{cases} \frac{\partial \mathcal{H}}{\partial u_1} = 2A_3 u_1 - \lambda_1 A_\beta = 0, & u_1^* = \min \left\{ u_1^{\max}, \max \left\{ 0, \frac{\lambda_1 A_\beta}{2A_3} \right\} \right\} \\ \frac{\partial \mathcal{H}}{\partial u_2} = 2A_4 u_2 - \lambda_2 Ca = 0, & u_2^* = \min \left\{ u_2^{\max}, \max \left\{ 0, \frac{\lambda_2 Ca}{2A_4} \right\} \right\} \\ \frac{\partial \mathcal{H}}{\partial u_3} = 2A_5 u_3 - \lambda_3 \tau = 0, & u_3^* = \min \left\{ u_3^{\max}, \max \left\{ 0, \frac{\lambda_3 \tau}{2A_5} \right\} \right\}. \end{cases}$$

Proof. The

□

References

- [1] Altaf A Abdulkhaliq et al. “Amyloid- β and Tau in Alzheimer’s disease: pathogenesis, mechanisms, and interplay”. In: *Cell Death & Disease* 17.1 (2026), p. 21.
- [2] Carlo Ballatore, Virginia M-Y Lee, and John Q Trojanowski. “Tau-mediated neurodegeneration in Alzheimer’s disease and related disorders”. In: *Nature reviews neuroscience* 8.9 (2007), pp. 663–672.
- [3] Alexandre Bejanin et al. “Tau pathology and neurodegeneration contribute to cognitive impairment in Alzheimer’s disease”. In: *Brain* 140.12 (2017), pp. 3286–3300.
- [4] Michael J Berridge. “Calcium hypothesis of Alzheimer’s disease”. In: *Pflügers archiv-european journal of physiology* 459.3 (2010), pp. 441–449.
- [5] Pei-Pei Guan, Long-Long Cao, and Pu Wang. “Elevating the levels of calcium ions exacerbate Alzheimer’s disease via inducing the production and aggregation of β -amyloid protein and phosphorylated tau”. In: *International journal of molecular sciences* 22.11 (2021), p. 5900.
- [6] Wenrui Hao, Suzanne Lenhart, and Jeffrey R Petrella. “Optimal anti-amyloid-beta therapy for Alzheimer’s disease via a personalized mathematical model”. In: *PLoS computational biology* 18.9 (2022), e1010481.
- [7] John A Hardy and Gerald A Higgins. “Alzheimer’s disease: the amyloid cascade hypothesis”. In: *Science* 256.5054 (1992), pp. 184–185.
- [8] Kasper P Kepp et al. “The amyloid cascade hypothesis: an updated critical review”. In: *Brain* 146.10 (2023), pp. 3969–3990.
- [9] Ruoyun Lang, Yuanshun Tan, and Yu Mu. “Stationary distribution and extinction of a stochastic Alzheimer’s disease model”. In: *AIMS Mathematics* 8.10 (2023), pp. 23313–23335.
- [10] Mitali Maji and Subhas Khajanchi. “Mathematical models on Alzheimer’s disease and its treatment: A review”. In: *Physics of Life Reviews* 52 (2025), pp. 207–244.
- [11] Mitali Maji, Laurent Pujo-Menjouet, and Subhas Khajanchi. “Quantifying uncertainty and sensitivity in an Alzheimer’s disease model: A mathematical approach”. In: *Acta Biotheoretica* 74.1 (2026), p. 5.
- [12] Arindam Nandi et al. “Global and regional projections of the economic burden of Alzheimer’s disease and related dementias from 2019 to 2050: a value of statistical life approach”. In: *EClinicalMedicine* 51 (2022).
- [13] Swadesh Pal and Roderick Melnik. “Nonequilibrium landscape of amyloid-beta and calcium ions in application to Alzheimer’s disease”. In: *Physical Review E* 111.1 (2025), p. 014418.
- [14] Jeffrey R Petrella et al. “Computational causal modeling of the dynamic biomarker cascade in Alzheimer’s disease”. In: *Computational and mathematical methods in medicine* 2019.1 (2019), p. 6216530.
- [15] Huiqin Zhang et al. “Interaction between A β and tau in the pathogenesis of Alzheimer’s disease”. In: *International journal of biological sciences* 17.9 (2021), p. 2181.

- [16] Jifa Zhang et al. “Recent advances in Alzheimer’s disease: Mechanisms, clinical trials and new drug development strategies”. In: *Signal transduction and targeted therapy* 9.1 (2024), p. 211.
- [17] Yongxin Zhang and Wendi Wang. “Mathematical analysis for stochastic model of Alzheimer’s disease”. In: *Communications in Nonlinear Science and Numerical Simulation* 89 (2020), p. 105347.
- [18] W-H Zheng et al. “Amyloid β peptide induces tau phosphorylation and loss of cholinergic neurons in rat primary septal cultures”. In: *Neuroscience* 115.1 (2002), pp. 201–211.