

Palmer amaranth (*Amaranthus palmeri*) adaptation to agroecosystems

Maxwel C Oliveira¹, Amit J Jhala², Mark Bernards³, Chris Proctor², Strahinja Stepanovic², Rodrigo Werle^{1*}

¹ Department of Agronomy, University of Wisconsin-Madison, Madison, Wisconsin, United States

² Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, Nebraska, United States

³ Department of Agronomy, Western Illinois University, Macomb, Illinois, United States

Correspondence*:

Rodrigo Werle

rwerle@uwisc.edu

2 ABSTRACT

3 Abstract length and content varies depending on article type. Refer to [http://](http://www.frontiersin.org/about/AuthorGuidelines)
4 www.frontiersin.org/about/AuthorGuidelines for abstract requirement and length
5 according to article type.

6 **Keywords:** Evolution Flowering Management Pigweed Weed

INTRODUCTION

7 Palmer amaranth (*Amaranthus palmeri* S. Watson) is currently considered one of the most economically
8 damaged weed species to cropping systems in the United States (Ward et al., 2013). The species has showed
9 a remarkable capacity to evolve resistance to herbicides. Palmer amaranth has evolved resistance to eight
10 herbicide sites of action (Heap, 2021), increasing the weed management complexity (Lindsay et al., 2017).
11 Uncontrolled Palmer amaranth in competition for water, light and nutrients can drastically reduce crop
12 yields (Berger et al., 2015). Palmer amaranth is documented with potential to reduce 91%, 68%, and 54%
13 of corn (Massinga et al., 2001), soybean (Klingaman and Oliver, 1994), and cotton (Morgan et al., 2001)
14 yields, respectively.

15 Palmer amaranth is a fast growing summer annual forb indigenous to Sonoran Desert (Sauer, 1957). The
16 species would eventually emerge as a threat to US agriculture in the 1990s. Palmer amaranth weediness
17 is likely a result of human-assisted selection in combination with species biology. Farm mechanization,
18 conservation agriculture (e.g., no-till), and reliance on herbicides for weed management are the main human
19 mediated selection of Palmer amaranth to cropping systems. On the other hand, Palmer amaranth is a
20 prolific seed producer with a C4 photosynthetic apparatus (Ward et al., 2013). With a dioecy nature, Palmer
21 amaranth male and female plants are obligate outcrosser species, increasing the chances of exchanging
22 herbicide resistant alleles among plants (Oliveira et al., 2018). Also, Palmer amaranth small seed size (e.g.,
23 1 mm) tend to thrive in no-tillage systems (Price et al., 2011), and spread across locations through farm
24 equipment (Sauer, 1972), manure (Hartzler and Anderson, 2016), animals (Farmer et al., 2017), and plant

propagules (Yu et al., 2021). Therefore, Palmer amaranth dispersal capacity make the species one of the most successful cases of weed adaption to cropping systems.

Light and temperature are likely the main environment requirements for Palmer amaranth successful adaptation. Palmer amaranth is reported with an extended germination period (Jha et al., 2010). Germination of Palmer amaranth is triggered by 18 C soil temperature (Keeley et al., 1987), and optimal germination and biomass production occur at 35/30 C day and night temperatures (Guo and Al-Khatib, 2003). Water has not shown to limit Palmer amaranth fitness. Under continuous water stress, Palmer amaranth survived and produced at least 14000 seeds plant⁻¹ (Chahal et al., 2018). Also, seeds from Palmer amaranth growing under water stress conditions were heavier, less dormant, and prompt for germination (Matzrafi et al., 2021). The continuous global temperature warming can impact agriculture and promote niches for Palmer amaranth invasion/adaptation into new environments. Currently, it is estimated that the greatest climatic risk of Palmer amaranth establishment are agronomic crops in Australia and Sub-Sahara Africa (Kistner and Hatfield, 2018). Temperature is a key factor limiting Palmer amaranth expansion to cooler geographies (Briscoe Runquist et al., 2019); however, under future climate change Palmer amaranth is likely to expand northward into Canada and Northern Europe (Kistner and Hatfield, 2018; Briscoe Runquist et al., 2019).

Palmer amaranth is already documented in agronomic crops of South America (Larran et al., 2017; Küpper et al., 2017) and Southern Europe (Milani et al., 2021). In the US, Palmer amaranth is established at crop (Garetson et al., 2019) and non-crop land (Bagavathiannan and Norsworthy, 2016) in the warm southern United States but its range is expanding to cool temperatures northward. For example, herbicide resistant Palmer amaranth is widespread in Nebraska (Oliveira et al., 2021), Michigan (Kohrt et al., 2017), and Connecticut (Aulakh et al., 2021). Successful cases of Palmer amaranth invasion and near to eradication is reported in Minnesota (Yu et al., 2021). No Palmer amaranth actively growing was found in Canada; however, Palmer amaranth seeds was detected in sweet potato slips (Page et al., 2021). Nonetheless, it seems fated the need to manage Palmer amaranth in agronomic crops throughout multiple environments in the near future. Therefore, strategies on Palmer amaranth management should encompass the agroecosystem level but not only attempts to eradicate the weed. Most tactics to manage Palmer amaranth are based on technology fixes (Scott, 2011), which are short-term (e.g., herbicide and/or tillage) rather than long-term weed management.

The continuous Palmer amaranth dispersal and potential establishment into northern United States warrant investigations on species morphology growing in such environments. Understanding Palmer amaranth biology and growing strategies under different agroecosystems can enhance our knowledge on species adaptation. It can also aid on designing proactive and ecological tactics to limit the species range expansion, reduce its negative impact, and design resilient and sustainable farming systems (MacLaren et al., 2020). Therefore, the objective of this study was to investigate the flowering pattern, biomass production, and height of Palmer amaranth growing under in corn, soybean and fallow at two timings across five locations in the mid/upper United States Midwest.

MATERIAL AND METHODS

Plant material and growing conditions

The study was performed with a *A. palmeri* accession (Per1) from Perkins County, Nebraska. Per1 accession collection is documented with no reported herbicide resistance (Oliveira et al., 2021). Three weeks prior to the field experiment, seeds were planted in plastic trays containing potting-mix. Emerged seedlings (1 cm) were transplanted into 200 cm³ plastic pots (a plant pot⁻¹). Palmer amaranth seedlings were supplied with adequate water and kept under greenhouse conditions at Arlington, Clay Center, Lincoln,

and Macomb; and kept outdoors in Grant. Palmer amaranth seedlings were kept under greenhouse/outdoors until the onset of the experiment (7 to 10 cm height).

Field study

The experiment was conducted in 2018 and 2019 under field conditions at five locations: Arlington (Washington County, Wisconsin), Clay Center (Clay County, Nebraska), Grant (Perkins County, Nebraska), Lincoln (Lancaster County, Nebraska), and Macomb (McDonough County, Illinois).

The field experimental unit were six adjacent 9.1 m wide (12 rows at 72.2 cm row spacing) by 10.7 m long. Each experimental unit was planted with corn or soybean, or under fallow condition. Palmer amaranth seedlings were transplanted to the field experiment by making a whole in the soil (6 cm deep and 8 cm wide); and gently transferring to the ground (potting mix + two seedlings). After a week, one was eliminated and one was kept. There were two transplant timing: first (June 1st) and second (July 1st). There were 24 Palmer amaranth plants in each experimental unit, with a total of 144 plants. The study was repeated twice.

After transplanting, Palmer amaranth flowering was monitored until the end of the study. When a plant started flowering, the day was recorded, plant sex was identified as male or female, and plant height was measured from soil surface to the plant top. Then, aboveground plant biomass was harvest near soil surface and oven dried at 65 C until reaching constant weight before the weight of biomass (g plant⁻¹) was recorded.

Statistical analyses

The statistical analyses were performed using R statistical software version 4.0.1. Data across locations and year were combined.

The cumulative Palmer amaranth flowering estimation was determined using a asymmetrical three parameter log logistic Weibull model of the drc package (Ritz et al., 2015).

$$Y(x) = 0 + (d - 0)\exp(-\exp(b(\log(x) - e)))$$

In this model, Y is the Palmer amaranth cumulative flowering, d is the upper limit (set to 100), and e is the XXX, and x day of year (doy).

The doy for 10, 50, and 90% Palmer amaranth cumulative flowering were determined using the *ED* function of drc package. Also, the 10, 50, and 90% Palmer amaranth cumulative flowering were compared among crop/fallow and timings using the *EDcomp* function of drc package. The *EDcomp* function compares the ratio of cumulative flowering using t-statistics, where P-value < 0.05 indicates that we fail to reject the null hypothesis.

Palmer amaranth height and biomass were performed with a linear mixed model using *lmer* function from “lme4” package (Bates et al., 2015). Plant height and biomass were transformed to meet model assumption of normality. In the model, agroecosystem (crop, soybean, fallow) was the fixed effect and year nested with location the random effects. Analysis of variance was performed with *anova* function from “car” package (Fox and Weisberg, 2018). Marginal means and compact letter display were estimated with *emmeans* and *cld* from packages “emmeans” and multcomp (Hothorn et al., 2008)

RESULTS

Cumulative flowering

You can use R chunks directly to plot graphs.

Palmer amaranth height and biomass

Frontiers requires figures to be submitted individually, in the same order as they are referred to in the manuscript. Figures will then be automatically embedded at the bottom of the submitted manuscript. Kindly ensure that each table and figure is mentioned in the text and in numerical order. Permission must be obtained for use of copyrighted material from other sources (including the web). Please note that it is compulsory to follow figure instructions. Figures which are not according to the guidelines will cause substantial delay during the production process.

1 DISCUSSION

DISCLOSURE/CONFLICT-OF-INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

RW: designed the experiments; AJ, CP, MB, MO, and SS: conducted the experiments; MO: analyzed the data and wrote the manuscript; AJ, CP, MB, MO, SS, and RW: conceptualized the research. All authors reviewed the manuscript.

ACKNOWLEDGMENTS

Funding: This work received no specific grant from any funding agency, commercial, or not-for-profit sectors

2 SUPPLEMENTAL DATA

Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures, please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be found in the Frontiers LaTeX folder

3 REFERENCES

A reference list should be automatically created here. However it won't. Pandoc will place the list of references at the end of the document instead. There are no convenient solution for now to force Pandoc to do otherwise. The easiest way to get around this problem is to edit the LaTeX file created by Pandoc before compiling it again using the traditional LaTeX commands.

FIGURES

Aulakh, J. S., Chahal, P. S., Kumar, V., Price, A. J., and Guillard, K. (2021). Multiple herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) in Connecticut: Confirmation and response to POST herbicides. *Weed Technology* 35, 457–463. doi:10.1017/wet.2021.6.

Bagavathiannan, M. V., and Norsworthy, J. K. (2016). Multiple-Herbicide Resistance Is Widespread in Roadside Palmer Amaranth Populations. *PLOS ONE* 11, e0148748. doi:10.1371/journal.pone.0148748.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using Lme4. *Journal of Statistical Software* 67, 1–48. doi:10.18637/jss.v067.i01.

Berger, S. T., Ferrell, J. A., Rowland, D. L., and Webster, T. M. (2015). Palmer Amaranth (*Amaranthus palmeri*) Competition for Water in Cotton. *Weed Science* 63, 928–935. doi:10.1614/WS-D-15-00062.1.




Figure 1. Mean average temperature (C) and monthly sum precipitation (mm) at Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL

- 134 Briscoe Runquist, R. D., Lake, T., Tiffin, P., and Moeller, D. A. (2019). Species distribution models
135 throughout the invasion history of Palmer amaranth predict regions at risk of future invasion and reveal
136 challenges with modeling rapidly shifting geographic ranges. *Sci Rep* 9, 2426. doi:10.1038/s41598-
137 018-38054-9.
- 138 Chahal, P. S., Irmak, S., Jugulam, M., and Jhala, A. J. (2018). Evaluating Effect of Degree of Water Stress
139 on Growth and Fecundity of Palmer amaranth (*Amaranthus palmeri*) Using Soil Moisture Sensors.
140 *Weed Science* 66, 738–745. doi:10.1017/wsc.2018.47.
- 141 Farmer, J. A., Webb, E. B., Pierce, R. A., and Bradley, K. W. (2017). Evaluating the potential for weed seed
142 dispersal based on waterfowl consumption and seed viability. *Pest Management Science* 73, 2592–2603.
143 doi:10.1002/ps.4710.
- 144 Fox, J., and Weisberg, S. (2018). *An R Companion to Applied Regression*. SAGE Publications Available at:
145 <http://books.google.com?id=uPNrDwAAQBAJ>.
- 146 Garetson, R., Singh, V., Singh, S., Dotray, P., and Bagavathiannan, M. (2019). Distribution of herbicide-
147 resistant Palmer amaranth (*Amaranthus palmeri*) in row crop production systems in Texas. *Weed*
148 *Technology* 33, 355–365. doi:10.1017/wet.2019.14.
- 149 Guo, P., and Al-Khatib, K. (2003). Temperature effects on germination and growth of redroot pigweed
150 (*Amaranthus retroflexus*), Palmer amaranth (*A. Palmeri*), and common waterhemp (*A. rudis*). *Weed*
151 *Science* 51, 869–875. doi:10.1614/P2002-127.
- 152 Hartzler, B., and Anderson, M. (2016). Palmer amaranth: It's here, now what? 10.




Figure 2. Figure caption

153 Heap, I. (2021). International Herbicide-Resistant Weed Database. Available at: [http://www.](http://www.weedscience.org/Home.aspx)
 154 [weedscience.org/Home.aspx](http://www.weedscience.org/Home.aspx) [Accessed July 26, 2021].

155 Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous Inference in General Parametric Models.
 156 *Biometrical Journal* 50, 346–363. doi:10.1002/bimj.200810425.

- Jha, P., Norsworthy, J. K., Riley, M. B., and Bridges, W. (2010). Annual Changes in Temperature and Light Requirements for Germination of Palmer Amaranth (*Amaranthus palmeri*) Seeds Retrieved from Soil. *Weed Science* 58, 426–432. doi:10.1614/WS-D-09-00038.1.
- Keeley, P. E., Carter, C. H., and Thullen, R. J. (1987). Influence of Planting Date on Growth of Palmer Amaranth (*Amaranthus palmeri*). *Weed Science* 35, 199–204. doi:10.1017/S0043174500079054.
- Kistner, E. J., and Hatfield, J. L. (2018). Potential Geographic Distribution of Palmer Amaranth under Current and Future Climates. *Agricultural & Environmental Letters* 3, 170044. doi:10.2134/acl2017.12.0044.
- Klingaman, T. E., and Oliver, L. R. (1994). Palmer Amaranth (*Amaranthus palmeri*) Interference in Soybeans (*Glycine max*). *Weed Science* 42, 523–527. doi:10.1017/S0043174500076888.
- Kohrt, J. R., Sprague, C. L., Nadakuduti, S. S., and Douches, D. (2017). Confirmation of a Three-Way (Glyphosate, ALS, and Atrazine) Herbicide-Resistant Population of Palmer Amaranth (*Amaranthus palmeri*) in Michigan. *Weed Science* 65, 327–338. doi:10.1017/wsc.2017.2.
- Küpper, A., Borgato, E. A., Patterson, E. L., Netto, A. G., Nicolai, M., Carvalho, S. J. P. de, Nissen, S. J., Gaines, T. A., and Christoffoleti, P. J. (2017). Multiple Resistance to Glyphosate and Acetolactate Synthase Inhibitors in Palmer Amaranth (*Amaranthus palmeri*) Identified in Brazil. *Weed Science* 65, 317–326. doi:10.1017/wsc.2017.1.
- Larran, A. S., Palmieri, V. E., Perotti, V. E., Lieber, L., Tuesca, D., and Permingeat, H. R. (2017). Target-site resistance to acetolactate synthase (ALS)-inhibiting herbicides in *Amaranthus palmeri* from Argentina. *Pest Management Science* 73, 2578–2584. doi:10.1002/ps.4662.
- Lindsay, K., Popp, M., Norsworthy, J., Bagavathiannan, M., Powles, S., and Lacoste, M. (2017). PAM: Decision Support for Long-Term Palmer Amaranth (*Amaranthus palmeri*) Control. *Weed Technology* 31, 915–927. doi:10.1017/wet.2017.69.
- MacLaren, C., Storkey, J., Menegat, A., Metcalfe, H., and Dehnen-Schmutz, K. (2020). An ecological future for weed science to sustain crop production and the environment. A review. *Agron. Sustain. Dev.* 40, 24. doi:10.1007/s13593-020-00631-6.
- Massinga, R. A., Currie, R. S., Horak, M. J., and Boyer, J. (2001). Interference of Palmer amaranth in corn. *Weed Science* 49, 202–208. doi:10.1614/0043-1745(2001)049[0202:IOPAIC]2.0.CO;2.
- Matzrafi, M., Osipitan, O. A., Ohadi, S., and Mesgaran, M. B. (2021). Under pressure: Maternal effects promote drought tolerance in progeny seed of Palmer amaranth (*Amaranthus palmeri*). *Weed Science* 69, 31–38. doi:10.1017/wsc.2020.75.
- Milani, A., Panozzo, S., Farinati, S., Iamonico, D., Sattin, M., Loddo, D., and Scarabel, L. (2021). Recent Discovery of *Amaranthus palmeri* S. Watson in Italy: Characterization of ALS-Resistant Populations and Sensitivity to Alternative Herbicides. *Sustainability* 13, 7003. doi:10.3390/su13137003.
- Morgan, G. D., Baumann, P. A., and Chandler, J. M. (2001). Competitive Impact of Palmer Amaranth (*Amaranthus palmeri*) on Cotton (*Gossypium hirsutum*) Development and Yield. *Weed Technology* 15, 408–412. doi:10.1614/0890-037X(2001)015[0408:CIOPAA]2.0.CO;2.
- Oliveira, M. C., Gaines, T. A., Patterson, E. L., Jhala, A. J., Irmak, S., Amundsen, K., and Knezevic, S. Z. (2018). Interspecific and intraspecific transference of metabolism-based mesotrione resistance in dioecious weedy *Amaranthus*. *The Plant Journal* 96, 1051–1063. doi:10.1111/tj.14089.

- 197 Oliveira, M. C., Giacomini, D. A., Arsenijevic, N., Vieira, G., Tranel, P. J., and Werle, R. (2021).
198 Distribution and validation of genotypic and phenotypic glyphosate and PPO-inhibitor resistance in
199 Palmer amaranth (*Amaranthus palmeri*) from southwestern Nebraska. *Weed Technology* 35, 65–76.
200 doi:10.1017/wet.2020.74.
- 201 Page, E. R., Nurse, R. E., Meloche, S., Bosveld, K., Grainger, C., Obeid, K., Filotas, M., Simard, M.-J., and
202 Laforest, M. (2021). Import of Palmer amaranth (*Amaranthus palmeri* S. Wats.) Seed with sweet potato
203 (*Ipomea batatas* (L.) Lam) slips. *Can. J. Plant Sci.*, CJPS-2020-0321. doi:10.1139/CJPS-2020-0321.
- 204 Price, A. J., Balkcom, K. S., Culpepper, S. A., Kelton, J. A., Nichols, R. L., and Schomberg, H. (2011).
205 Glyphosate-resistant Palmer amaranth: A threat to conservation tillage. *Journal of Soil and Water*
206 *Conservation* 66, 265–275. doi:10.2489/jswc.66.4.265.
- 207 Ritz, C., Baty, F., Streibig, J. C., and Gerhard, D. (2015). Dose-Response Analysis Using R. *PLOS ONE*
208 10, e0146021. doi:10.1371/journal.pone.0146021.
- 209 Sauer, J. (1957). Recent Migration and Evolution of the Dioecious Amaranths. *Evolution* 11, 11–31.
210 doi:10.2307/2405808.
- 211 Sauer, J. D. (1972). The dioecious amaranths: A new species name and major range extensions. *Madroño*
212 21, 426–434. Available at: <http://www.jstor.org/stable/41423815>.
- 213 Scott, D. (2011). The Technological Fix Criticisms and the Agricultural Biotechnology Debate. *J Agric*
214 *Environ Ethics* 24, 207–226. doi:10.1007/s10806-010-9253-7.
- 215 Ward, S. M., Webster, T. M., and Steckel, L. E. (2013). Palmer Amaranth (*Amaranthus palmeri*): A Review.
216 *Weed Technology* 27, 12–27. doi:10.1614/WT-D-12-00113.1.
- 217 Yu, E., Blair, S., Hardel, M., Chandler, M., Thiede, D., Cortilet, A., Gunsolus, J., and Becker, R. (2021).
218 Timeline of Palmer amaranth (*Amaranthus palmeri*) invasion and eradication in Minnesota. *Weed*
219 *Technology*, 1–31. doi:10.1017/wet.2021.32.