

# Palmer amaranth (*Amaranthus palmeri*) adaptation to agroecosystems

Maxwel C Oliveira<sup>1</sup>, Amit J Jhala<sup>2</sup>, Mark Bernards<sup>3</sup>, Chris Proctor<sup>2</sup>, Strahinja Stepanovic<sup>2</sup>, Rodrigo Werle<sup>1\*</sup>

<sup>1</sup> Department of Agronomy, University of Wisconsin-Madison, Madison, Wisconsin, United States

<sup>2</sup> Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, Nebraska, United States

<sup>3</sup> Department of Agronomy, Western Illinois University, Macomb, Illinois, United States

Correspondence\*:

Rodrigo Werle

rwerle@uwisc.edu

## 2 ABSTRACT

Abstract length and content varies depending on article type. Refer to <http://www.frontiersin.org/about/AuthorGuidelines> for abstract requirement and length according to article type.

Keywords: Evolution Flowering Management Pigweed Weed

## INTRODUCTION

Palmer amaranth (*Amaranthus palmeri* S. Watson) is currently considered one of the most economically damaged weed species to cropping systems in the United States. Unmanaged Palmer amaranth in competition for water, light and nutrients can drastically impact on crop yields (Berger et al., 2015). For example, Palmer amaranth is documented with potential to reduce 91%, 68%, and 54% of corn (Massinga et al., 2001), soybean (Klingaman and Oliver, 1994), and cotton (Morgan et al., 2001) yields, respectively. Moreover, Palmer amaranth has showed a remarkable capacity to evolve resistance to herbicides. To date, Palmer amaranth has evolved resistance to eight herbicide sites of action (Heap, 2021), increasing the weed management complexity (Lindsay et al., 2017). Thus, Palmer amaranth poses an economical risk to sustainable agriculture.

Palmer amaranth is a fast growing summer annual forb indigenous to Sonoran Desert (Sauer, 1957). The species would eventually emerge as a threat to US agriculture in the 1990s. Palmer amaranth weediness is likely a result of human-assisted selection in combination with species biology. Farm mechanization, conservation agriculture (e.g., no-till), and reliance on herbicides for weed management are the main human-mediated selection of Palmer amaranth into cropping systems (Ward et al., 2013). On the other hand, Palmer amaranth is a prolific seed producer with a C4 photosynthetic apparatus (Wang et al., 1992). With a dioecy nature, Palmer amaranth male and female plants are obligate outcrosser species, increasing the chances of exchanging adaptive traits among plants (Oliveira et al., 2018). Also, Palmer amaranth small seed size (e.g, 1 mm) tend to thrive in no-tillage systems (Price et al., 2011), and spread across locations through farm equipment (Sauer, 1972), manure (Hartzler and Anderson, 2016), animals (Farmer et al.,

26 2017), and plant propagules (Yu et al., 2021). The dispersal capacity of Palmer amaranth make the species  
27 one of the most successful cases of weed adaption to cropping systems.

28 Light and temperature are likely the main environment requirements for Palmer amaranth successful  
29 grow and development. Palmer amaranth is reported with an extended germination period (Jha et al., 2010).  
30 Germination of Palmer amaranth is triggered by 18 C soil temperature (Keeley et al., 1987), and optimal  
31 germination and biomass production occur at 35/30 C day and night temperatures (Guo and Al-Khatib,  
32 2003). Also, Palmer amaranth germination are human-mediated by tillage timings and preemergence-  
33 applied herbicides (Chahal et al., 2021). Water has not shown to limit Palmer amaranth fitness. Under  
34 continuous water stress, Palmer amaranth survived and produced at least 14000 seeds plant-1 (Chahal et  
35 al., 2018). Also, seeds from Palmer amaranth growing with limited water conditions were heavier, less  
36 dormant, and prompt for germination (Matzrafi et al., 2021). The continuous global temperature warming  
37 can impact agriculture and promote niches for Palmer amaranth invasion/adaptation into new environments.  
38 Currently, it is estimated that the greatest climatic risk of Palmer amaranth establishment are agronomic  
39 crops in Australia and Sub-Saharan Africa (Kistner and Hatfield, 2018). Temperature is a key factor limiting  
40 Palmer amaranth expansion to cooler geographies (Briscoe Runquist et al., 2019); however, under future  
41 climate change Palmer amaranth is likely to expand northward into Canada and Northern Europe (Kistner  
42 and Hatfield, 2018; Briscoe Runquist et al., 2019).

43 Palmer amaranth is already found in agronomic crops of South America (Larran et al., 2017; Küpper et al.,  
44 2017) and Southern Europe (Milani et al., 2021). In the US, Palmer amaranth is well established at cotton  
45 fields (Garetson et al., 2019) and non-crop areas (Bagavathiannan and Norsworthy, 2016) in the southern  
46 United States but its range is expanding northward. For example, herbicide resistant Palmer amaranth is  
47 widespread in Nebraska (Oliveira et al., 2021), Michigan (Kohrt et al., 2017), Connecticut (Aulakh et al.,  
48 2021), and potential damaged to soybean fields in Illinois is reported (Davis et al., 2015). Successful cases  
49 of Palmer amaranth invasion and near to eradication is documented in Minnesota (Yu et al., 2021). No  
50 Palmer amaranth actively growing was found in Canada; however, Palmer amaranth seeds was detected in  
51 sweet potato slips (Page et al., 2021). Nonetheless, it seems fated the need to manage Palmer amaranth  
52 in agronomic crops throughout multiple environments in the near future. Strategies on Palmer amaranth  
53 management should encompass the agroecosystem level but not only attempts to eradicate the weed. Most  
54 tactics to manage Palmer amaranth are based on technology fixes (Scott, 2011), which are short-term (e.g.,  
55 herbicide and/or tillage) rather than long-term weed management. Palmer amaranth management should be  
56 built on minimizing the species ability to adapt, grow and develop into agroecossystems.

57 In the southeastern US, early growing Palmer amaranth is well known to have higher impact on cotton  
58 yields than late established plants (MacRae et al., 2013). In the northern states, Palmer amaranth impact on  
59 the agroecosystem is recent. Studies investigating Palmer amaranth in those locations is limited due to  
60 the plant classification as noxious weed (Yu et al., 2021). Nonetheless, the continuous Palmer amaranth  
61 dispersal and potential establishment across northern United States is concerning and warrant investigations  
62 on species morphology in such environments. Understanding Palmer amaranth biology and growing  
63 strategies under different agroecosystems can enhance our knowledge on species adaptation. It can also  
64 aid on designing proactive and ecological tactics to limit the species range expansion, reduce its negative  
65 impact, and design resilient and sustainable farming systems (MacLaren et al., 2020). Therefore, the  
66 objective of this study was to investigate the flowering pattern, biomass production, and height of Palmer  
67 amaranth growing under corn, soybean and fallow at two timings across five locations in the United States  
68 Midwest.

## MATERIAL AND METHODS

### 69 Plant material and growing conditions

70 The study was performed with a *A. palmeri* accession (Per1) from Perkins County, Nebraska. Per1  
71 accession collection is documented with no reported herbicide resistance (Oliveira et al., 2021). Three  
72 weeks prior to the field experiment, seeds were planted in plastic trays containing potting-mix. Emerged  
73 seedlings (1 cm) were transplanted into 200 cm<sup>3</sup> plastic pots (a plant pot-1). Palmer amaranth seedlings  
74 were supplied with adequate water and kept under greenhouse conditions at Arlington, Clay Center, Lincoln,  
75 and Macomb; and kept outdoors in Grant. Palmer amaranth seedlings were kept under greenhouse/outdoors  
76 until the onset of the experiment (2-3 leaf stage/5 to 8 cm height).

### 77 Field study

78 The experiment was conducted in 2018 and 2019 under field conditions at five locations: Arlington  
79 (Washington County, Wisconsin), Clay Center (Clay County, Nebraska), Grant (Perkins County, Nebraska),  
80 Lincoln (Lancaster County, Nebraska), and Macomb (McDonough County, Illinois).

81 A glyphosate-resistant soybean cultivar (DSR-1950 R2Y at 296,400 seeds ha<sup>-1</sup>), and a corn hybrid were  
82 planted at

83 Monthly mean air temperature and sum precipitation were obtained using Daymet weather data from  
84 June through September across the five locations in 2018 and 2019 (Correndo et al., 2021) (Figure 1)

85 The field experimental unit were six adjacent 9.1 m wide (12 rows at 72.2 cm row spacing) by 10.7  
86 m long. Each experimental unit was planted with corn or soybean (DSR-1950 R2Y at 296,400 seeds ha<sup>-1</sup>), or under fallow condition. Palmer amaranth seedlings (potting mix + two seedlings) were and gently  
87 transferring to the ground (6 cm deep and 8 cm wide). Twenty-four plants were equidistantly placed (0.76  
88 m apart) between rows within each agroecosystems. After a week, one was eliminated and one was kept.  
90 There were two transplant timing: first (June 1) and second (July 1). There were 24 Palmer amaranth plants  
91 in each experimental unit, with a total of 144 plants for each location. The study was repeated twice.

92 After transplanting, Palmer amaranth flowering was monitored until the end of the study. When a plant  
93 started flowering, the day was recorded, plant sex was identified as male or female, and plant height  
94 was measured from soil surface to the plant top. Then, aboveground plant biomass was harvest near soil  
95 surface and oven dried at 65 C until reaching constant weight before the weight of biomass (g plant<sup>-1</sup>) was  
96 recorded.

### 97 Statistical analyses

98 The statistical analyses were performed using R statistical software version 4.0.1. Data across locations  
99 and year were combined.

100 The cumulative Palmer amaranth flowering estimation was determined using a asymmetrical three  
101 parameter log logistic Weibull model of the drc package (Ritz et al., 2015).

$$Y(x) = 0 + (d - 0)\exp(-\exp(b(\log(x) - e)))$$

102 In this model, *Y* is the Palmer amaranth cumulative flowering, *d* is the upper limit (set to 100), and *e* is the  
103 XXX, and *x* day of year (doy).

104 The doy for 10, 50, and 90% Palmer amaranth cumulative flowering were determined using the *ED*  
105 function of drc package. Also, the 10, 50, and 90% Palmer amaranth cumulative flowering were compared  
106 among agroecosystems and timings using the *EDcomp* function of drc package. The EDcomp function

107 compares the ratio of cumulative flowering using t-statistics, where P-value < 0.05 indicates that we fail to  
108 reject the null hypothesis.

109 Palmer amaranth height and biomass were performed with a linear mixed model using *lmer* function from  
110 “lme4” package (Bates et al., 2015). Plant height and biomass were transformed to meet model assumption  
111 of normality. In the model, agroecosystem (crop, soybean, fallow) was the fixed effect and year nested with  
112 location the random effects. Analysis of variance was performed with *anova* function from “car” package  
113 (Fox and Weisberg, 2018). Marginal means and compact letter display were estimated with *emmeans* and  
114 *cld* from packages “emmeans” and *multcomp* (Hothorn et al., 2008).

## RESULTS

### 115 Cumulative flowering

116 Palmer amaranth growing in corn resulted in a longer flowering pattern compared to fallow and soybean  
117 at first cohort (Figure 2A). The 10% cumulative Palmer amaranth flowering in soybean, fallow and corn  
118 occurred at the end of June. Palmer amaranth reached 10% flowering in soybean at doy 180, which was  
119 slightly different from fallow (doy 180.9;  $P = 0.01$ ) and corn (doy 181.7;  $P = 0.00$ ). The 50% Palmer  
120 amaranth cumulative flowering occurred in July. For example, Palmer amaranth reached 50% in fallow at  
121 doy 193.4, followed by soybean (doy 194.8), corn (doy 206.6). Similar trend was observed at 90% Palmer  
122 amaranth cumulative flowering. Palmer amaranth growing in corn reached 90% flowering at doy 252.6  
123 (early September), which was 37.8 and 32.2 days after Palmer amaranth 90% flowering in fallow and  
124 soybean, respectively.

125 Palmer amaranth cumulative flowering at second cohort ranged from mid July to mid September (Figure  
126 2). All cumulative flowering comparisons among agroecosystems were significant different ( $P < 0.00$ )  
127 Palmer amaranth growing in fallow resulted in earlier flowering time compared to soybean and corn. Palmer  
128 amaranth growing in fallow reached 10%, 50%, and 90% flowering time at day 203.8, 214.4, and 232.2,  
129 respectively. Palmer amaranth growing in soybean reached 10% flowering at doy 210.9, which was 6 days  
130 prior to corn ( $P$ -value = 0.00). Similar trend was observed at 50% flowering, whereas Palmer amaranth  
131 reached 50% flowering in corn (doy 233.0) 4 days after soybeans (doy 228.9;  $P = 0.00$ ). The 90% Palmer  
132 amaranth cumulative flowering occurred at same day in corn (260.9) and soybean (260.5;  $P = 0.66$ ).

### 133 Height and biomass

134 Palmer amaranth accumulated more biomass when growing in fallow compared to Palmer amaranth  
135 growing in soybean and corn (Figure 3A). At first cohort time, Palmer amaranth biomass was 78.3 g plant-1,  
136 26.4 g plant-1 and 14.9 g plant-1 in fallow, soybean and corn, respectively. Moreover, Palmer amaranth  
137 produce less biomass at cohort time. Palmer amaranth growing in fallow resulted in 49.2 g plant-1, 3.4 g  
138 plant-1 in soybean and 1.4 g plant when growing in corn.

139 The Palmer amaranth achieve 92.3 cm tall when growing at first cohort timing in corn (Figure 3B).

## DISCUSSION

140 Our study showed that Palmer amaranth flowering time, biomass and height varied within agroecosystems  
141 and cohort timing. In general, Palmer amaranth produced more biomass and taller plants when growing  
142 at first cohort rather than second cohort. At first cohort, resources (e.g., soil nutrients) and conditions  
143 (e.g., light) were more timely available for both species, crop and weed. High biomass and taller Palmer  
144 amaranth plants is likely a weed strategy to compete for light in between crop rows in absence of canopy.  
145 In such condition, Palmer amaranth showed an extraordinary plasticity to adapt upon the agroecosystem.  
146 This is evident when comparing Palmer amaranth shape, and its extended flowering pattern when growing

147 into corn compared to soybean. In corn, Palmer amaranth was taller and thinner when growing in corn.  
148 The Palmer amaranth competition strategy was likely to mimic the crop grow and development (Figure 4).  
149 These results are concerning as breeding more competitive crop varieties is likely to select more competitive  
150 Palmer amaranth biotypes (Bravo et al., 2017).


151 Palmer amaranth grow and development at second cohort was limited due to the crop competitive ability  
152 at advanced development stages. Palmer amaranth was transplanted when corn canopy was nearly closed,  
153 which reduced Palmer amaranth competitiveness. As a result, Palmer amaranth height and biomass was  
154 lower compared to its first cohort. Under crop canopy, Palmer amaranth flowering pattern was near to  
155 similar into corn and soybean. Palmer amaranth growing without crop competition produced highest  
156 amounts of biomass and lowest extended flowering pattern. The Palmer amaranth strategy in bareground  
157 was to invest biomass in growing plant width and height. Nonetheless, Palmer amaranth produced 31% less  
158 biomass in second cohort compared to first cohort timing. In a bareground study, early emerged Palmer  
159 amaranth without competition was 50% taller than late emerged plants (Webster and Grey, 2015). These  
160 results suggests that crop competition is not the only factor limiting late Palmer amaranth establishment.  
161 The limited growth of Palmer amaranth at second cohort is likely a reduced plant response to thermal  
162 units (e.g, growing degree days). It is hypothesize that reduced day length contributed to smaller plants at  
163 second cohort as well as shorter flowering period. A study in North Carolina and Illinois predicted that less  
164 than 10% Palmer amaranth seedlings emergence occurred after June (Piskackova et al., 2021). In addition,  
165 Palmer amaranth negative impact on soybean (Korres et al., 2020) and cotton (Webster and Grey, 2015)  
166 yields was higher when plants were established near to crop planting.

167 Seed production was not evaluated due to Palmer amaranth harvest at flowering time. Nonetheless, it  
168 is well documented a strong positive correlation between Palmer amaranth biomass and seed production  
169 (Schwartz et al., 2016; Spaunhorst et al., 2018). In our study, Palmer amaranth growing at first cohort  
170 accumulated an overall 50% more biomass when compared to second cohort. Therefore, Palmer amaranth  
171 plants growing in the second cohort is likely to produce less seeds regardless the agroecosystem. Our  
172 observation is consistent with the findings that first Palmer amaranth cohort produced 50% more seeds per  
173 plant than Palmer amaranth plants established six weeks later in bareground (Webster and Grey, 2015).  
174 Nonetheless, seed production at second cohort is likely to replenish the soil seedbank. Seed production and  
175 deposition in the seedbank is a key factor for species perpetuation (Menges, 1987). Palmer amaranth can  
176 produce hundred thousands seeds per plant (Schwartz et al., 2016; Keeley et al., 1987), and stay viable  
177 buried in the seedbank for at least 36 months (Sosnoskie et al., 2013).

178 Our study demonstrated the short-term Palmer amaranth plasticity to grow and develop into cropping-  
179 systems. Palmer amaranth management priority should focus on minimizing Palmer amaranth dispersal  
180 into new geographies. Early-season management programs would have a large negative effect in Palmer  
181 amaranth growth and development. Tactics that promote crop advantage against Palmer amaranth, including  
182 early crop planting, crop rotation, plant width, preemergence applied herbicide (Sanctis et al., 2021), and  
183 crop residue (e.g. cover crops) would minimize the negative impact of Palmer amaranth in agroecosystems.

## DISCLOSURE/CONFLICT-OF-INTEREST STATEMENT

184 The authors declare that the research was conducted in the absence of any commercial or financial  
185 relationships that could be construed as a potential conflict of interest.



**Figure 1.** Mean average temperature (C) and monthly sum precipitation (mm) at Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL

## AUTHOR CONTRIBUTIONS

186 RW: designed the experiments; AJ, CP, MB, MO, and SS: conducted the experiments; MO: analyzed the  
 187 data and wrote the manuscript; AJ, CP, MB, MO, SS, and RW: conceptualized the research. All authors  
 188 reviewed the manuscript.

## ACKNOWLEDGMENTS


189 Funding: This work received no specific grant from any funding agency, commercial, or not-for-profit  
 190 sectors

## 1 SUPPLEMENTAL DATA


191 Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures,  
 192 please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be  
 193 found in the Frontiers LaTeX folder

## 2 REFERENCES

194 A reference list should be automatically created here. However it won't. Pandoc will place the list of  
 195 references at the end of the document instead. There are no convenient solution for now to force Pandoc to  
 196 do otherwise. The easiest way to get around this problem is to edit the LaTeX file created by Pandoc before  
 197 compiling it again using the traditional LaTeX commands.



**Figure 2.** Cumulative flowering of Palmer amaranth at first and second transplant timing (A) and day of year of 10, 50, and 90 cumulative flowering at first and second transplant timing (B)



**Figure 3.** Palmer amaranth biomass (A) and height (B) growing in corn, fallow, and soybean across Arlington, Clay Center, Grant, Lincoln, and Macomb



**Figure 4.** Harvest Palmer amaranth plants at 40 days after first transplant timing. From left to right, Palmer amaranth growing in fallow, soybean and corn in Arlington, Wisconsin

## FIGURES

- 198 Aulakh, J. S., Chahal, P. S., Kumar, V., Price, A. J., and Guillard, K. (2021). Multiple herbicide-resistant  
199 Palmer amaranth (*Amaranthus palmeri*) in Connecticut: Confirmation and response to POST herbicides.  
200 *Weed Technology* 35, 457–463. doi:10.1017/wet.2021.6.
- 201 Bagavathiannan, M. V., and Norsworthy, J. K. (2016). Multiple-Herbicide Resistance Is Widespread in  
202 Roadside Palmer Amaranth Populations. *PLOS ONE* 11, e0148748. doi:10.1371/journal.pone.0148748.
- 203 Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using  
204 *Lme4*. *Journal of Statistical Software* 67, 1–48. doi:10.18637/jss.v067.i01.
- 205 Berger, S. T., Ferrell, J. A., Rowland, D. L., and Webster, T. M. (2015). Palmer Amaranth (*Amaranthus*  
206 *palmeri*) Competition for Water in Cotton. *Weed Science* 63, 928–935. doi:10.1614/WS-D-15-00062.1.
- 207 Bravo, W., Leon, R. G., Ferrell, J. A., Mulvaney, M. J., and Wood, C. W. (2017). Differentiation of  
208 Life-History Traits among Palmer Amaranth Populations (*Amaranthus palmeri*) and Its Relation to  
209 Cropping Systems and Glyphosate Sensitivity. *Weed Science* 65, 339–349. doi:10.1017/wsc.2017.14.
- 210 Briscoe Runquist, R. D., Lake, T., Tiffin, P., and Moeller, D. A. (2019). Species distribution models  
211 throughout the invasion history of Palmer amaranth predict regions at risk of future invasion and reveal  
212 challenges with modeling rapidly shifting geographic ranges. *Sci Rep* 9, 2426. doi:10.1038/s41598-  
213 018-38054-9.
- 214 Chahal, P. S., Barnes, E. R., and Jhala, A. J. (2021). Emergence pattern of Palmer amaranth (*Amaranthus*  
215 *palmeri*) influenced by tillage timings and residual herbicides. *Weed Technology* 35, 433–439.  
216 doi:10.1017/wet.2020.136.
- 217 Chahal, P. S., Irmak, S., Jugulam, M., and Jhala, A. J. (2018). Evaluating Effect of Degree of Water Stress  
218 on Growth and Fecundity of Palmer amaranth (*Amaranthus palmeri*) Using Soil Moisture Sensors.  
219 *Weed Science* 66, 738–745. doi:10.1017/wsc.2018.47.

- 220 Correndo, A. A., Moro Rosso, L. H., and Ciampitti, I. A. (2021). Retrieving and processing agro-  
221 meteorological data from API-client sources using R software. *BMC Research Notes* 14, 205.  
222 doi:10.1186/s13104-021-05622-8.
- 223 Davis, A. S., Schutte, B. J., Hager, A. G., and Young, B. G. (2015). Palmer Amaranth (*Amaranthus palmeri*)  
224 Damage Niche in Illinois Soybean Is Seed Limited. *Weed Science* 63, 658–668. doi:10.1614/WS-D-14-  
225 00177.1.
- 226 Farmer, J. A., Webb, E. B., Pierce, R. A., and Bradley, K. W. (2017). Evaluating the potential for weed seed  
227 dispersal based on waterfowl consumption and seed viability. *Pest Management Science* 73, 2592–2603.  
228 doi:10.1002/ps.4710.
- 229 Fox, J., and Weisberg, S. (2018). *An R Companion to Applied Regression*. SAGE Publications Available at:  
230 <http://books.google.com?id=uPNrDwAAQBAJ>.
- 231 Garetson, R., Singh, V., Singh, S., Dotray, P., and Bagavathiannan, M. (2019). Distribution of herbicide-  
232 resistant Palmer amaranth (*Amaranthus palmeri*) in row crop production systems in Texas. *Weed  
233 Technology* 33, 355–365. doi:10.1017/wet.2019.14.
- 234 Guo, P., and Al-Khatib, K. (2003). Temperature effects on germination and growth of redroot pigweed  
235 (*Amaranthus retroflexus*), Palmer amaranth (*A. Palmeri*), and common waterhemp (*A. rudis*). *Weed  
236 Science* 51, 869–875. doi:10.1614/P2002-127.
- 237 Hartzler, B., and Anderson, M. (2016). Palmer amaranth: It's here, now what? 10.
- 238 Heap, I. (2021). Internation Herbicide-Resistant Weed Database. Available at: <http://www.weedscience.org/Home.aspx> [Accessed July 26, 2021].
- 239
- 240 Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous Inference in General Parametric Models.  
241 *Biometrical Journal* 50, 346–363. doi:10.1002/bimj.200810425.
- 242 Jha, P., Norsworthy, J. K., Riley, M. B., and Bridges, W. (2010). Annual Changes in Temperature and Light  
243 Requirements for Germination of Palmer Amaranth (*Amaranthus palmeri*) Seeds Retrieved from Soil.  
244 *Weed Science* 58, 426–432. doi:10.1614/WS-D-09-00038.1.
- 245 Keeley, P. E., Carter, C. H., and Thullen, R. J. (1987). Influence of Planting Date on Growth of Palmer  
246 Amaranth (*Amaranthus palmeri*). *Weed Science* 35, 199–204. doi:10.1017/S0043174500079054.
- 247 Kistner, E. J., and Hatfield, J. L. (2018). Potential Geographic Distribution of Palmer  
248 Amaranth under Current and Future Climates. *Agricultural & Environmental Letters* 3, 170044.  
249 doi:10.2134/ael2017.12.0044.
- 250 Klingaman, T. E., and Oliver, L. R. (1994). Palmer Amaranth (*Amaranthus palmeri*) Interference in  
251 Soybeans (*Glycine max*). *Weed Science* 42, 523–527. doi:10.1017/S0043174500076888.
- 252 Kohrt, J. R., Sprague, C. L., Nadakuduti, S. S., and Douches, D. (2017). Confirmation of a Three-Way  
253 (Glyphosate, ALS, and Atrazine) Herbicide-Resistant Population of Palmer Amaranth (*Amaranthus  
254 palmeri*) in Michigan. *Weed Science* 65, 327–338. doi:10.1017/wsc.2017.2.
- 255 Korres, N. E., Norsworthy, J. K., Mauromoustakos, A., and Williams, M. M. (2020). Soybean density and  
256 Palmer amaranth (*Amaranthus palmeri*) establishment time: Effects on weed biology, crop yield, and  
257 economic returns. *Weed Science* 68, 467–475. doi:10.1017/wsc.2020.41.

- 258 Küpper, A., Borgato, E. A., Patterson, E. L., Netto, A. G., Nicolai, M., Carvalho, S. J. P. de, Nissen, S.  
259 J., Gaines, T. A., and Christoffoleti, P. J. (2017). Multiple Resistance to Glyphosate and Acetolactate  
260 Synthase Inhibitors in Palmer Amaranth (*Amaranthus palmeri*) Identified in Brazil. *Weed Science* 65,  
261 317–326. doi:10.1017/wsc.2017.1.
- 262 Larran, A. S., Palmieri, V. E., Perotti, V. E., Lieber, L., Tuesca, D., and Permingeat, H. R. (2017). Target-site  
263 resistance to acetolactate synthase (ALS)-inhibiting herbicides in *Amaranthus palmeri* from Argentina.  
264 *Pest Management Science* 73, 2578–2584. doi:10.1002/ps.4662.
- 265 Lindsay, K., Popp, M., Norsworthy, J., Bagavathiannan, M., Powles, S., and Lacoste, M. (2017). PAM:  
266 Decision Support for Long-Term Palmer Amaranth (*Amaranthus palmeri*) Control. *Weed Technology*  
267 31, 915–927. doi:10.1017/wet.2017.69.
- 268 MacLaren, C., Storkey, J., Menegat, A., Metcalfe, H., and Dehnen-Schmutz, K. (2020). An ecological  
269 future for weed science to sustain crop production and the environment. A review. *Agron. Sustain. Dev.*  
270 40, 24. doi:10.1007/s13593-020-00631-6.
- 271 MacRae, A. W., Webster, T. M., Sosnoskie, L. M., Culpepper, A. S., and Kichler, J. M. (2013). Cotton  
272 Yield Loss Potential in Response to Length of Palmer Amaranth (*Amaranthus palmeri*) Interference.  
273 17, 6.
- 274 Massinga, R. A., Currie, R. S., Horak, M. J., and Boyer, J. (2001). Interference of Palmer amaranth in corn.  
275 *Weed Science* 49, 202–208. doi:10.1614/0043-1745(2001)049[0202:IOPAIC]2.0.CO;2.
- 276 Matzrafi, M., Osipitan, O. A., Ohadi, S., and Mesgaran, M. B. (2021). Under pressure: Maternal effects  
277 promote drought tolerance in progeny seed of Palmer amaranth (*Amaranthus palmeri*). *Weed Science*  
278 69, 31–38. doi:10.1017/wsc.2020.75.
- 279 Menges, R. M. (1987). Weed Seed Population Dynamics during Six Years of Weed Management Systems  
280 in Crop Rotations on Irrigated Soil. *Weed Science* 35, 328–332. Available at: <http://www.jstor.org/stable/4044593>.
- 282 Milani, A., Panozzo, S., Farinati, S., Iamonico, D., Sattin, M., Loddo, D., and Scarabel, L. (2021). Recent  
283 Discovery of *Amaranthus palmeri* S. Watson in Italy: Characterization of ALS-Resistant Populations  
284 and Sensitivity to Alternative Herbicides. *Sustainability* 13, 7003. doi:10.3390/su13137003.
- 285 Morgan, G. D., Baumann, P. A., and Chandler, J. M. (2001). Competitive Impact of Palmer Amaranth  
286 (*Amaranthus palmeri*) on Cotton (*Gossypium hirsutum*) Development and Yield. *Weed Technology* 15,  
287 408–412. doi:10.1614/0890-037X(2001)015[0408:CIOPAA]2.0.CO;2.
- 288 Oliveira, M. C., Gaines, T. A., Patterson, E. L., Jhala, A. J., Irmak, S., Amundsen, K., and Knezevic, S.  
289 Z. (2018). Interspecific and intraspecific transference of metabolism-based mesotrione resistance in  
290 dioecious weedy *Amaranthus*. *The Plant Journal* 96, 1051–1063. doi:10.1111/tpj.14089.
- 291 Oliveira, M. C., Giacomini, D. A., Arsenijevic, N., Vieira, G., Tranel, P. J., and Werle, R. (2021).  
292 Distribution and validation of genotypic and phenotypic glyphosate and PPO-inhibitor resistance in  
293 Palmer amaranth (*Amaranthus palmeri*) from southwestern Nebraska. *Weed Technology* 35, 65–76.  
294 doi:10.1017/wet.2020.74.
- 295 Page, E. R., Nurse, R. E., Meloche, S., Bosveld, K., Grainger, C., Obeid, K., Filotas, M., Simard, M.-J., and  
296 Laforest, M. (2021). Import of Palmer amaranth (*Amaranthus palmeri* S. Wats.) Seed with sweet potato  
297 (*Ipomea batatas* (L.) Lam) slips. *Can. J. Plant Sci.*, CJPS-2020-0321. doi:10.1139/CJPS-2020-0321.

- 298 Piskackova, T. A. R., Reberg-Horton, S. C., Richardson, R. J., Jennings, K. M., Franca, L., Young, B. G.,  
299 and Leon, R. G. (2021). Windows of action for controlling palmer amaranth (*Amaranthus palmeri*)  
300 using emergence and phenology models. *Weed Research* 61, 188–198. doi:10.1111/wre.12470.
- 301 Price, A. J., Balkcom, K. S., Culpepper, S. A., Kelton, J. A., Nichols, R. L., and Schomberg, H. (2011).  
302 Glyphosate-resistant Palmer amaranth: A threat to conservation tillage. *Journal of Soil and Water  
303 Conservation* 66, 265–275. doi:10.2489/jswc.66.4.265.
- 304 Ritz, C., Baty, F., Streibig, J. C., and Gerhard, D. (2015). Dose-Response Analysis Using R. *PLOS ONE*  
305 10, e0146021. doi:10.1371/journal.pone.0146021.
- 306 Sanctis, J. H. S. de, Barnes, E. R., Knezevic, S. Z., Kumar, V., and Jhala, A. J. (2021). Residual herbicides  
307 affect critical time of Palmer amaranth removal in soybean. *Agronomy Journal* 113, 1920–1933.  
308 doi:10.1002/agj2.20615.
- 309 Sauer, J. (1957). Recent Migration and Evolution of the Dioecious Amaranths. *Evolution* 11, 11–31.  
310 doi:10.2307/2405808.
- 311 Sauer, J. D. (1972). The dioecious amaranths: A new species name and major range extensions. *Madroño*  
312 21, 426–434. Available at: <http://www.jstor.org/stable/41423815>.
- 313 Schwartz, L. M., Norsworthy, J. K., Young, B. G., Bradley, K. W., Kruger, G. R., Davis, V. M., Steckel,  
314 L. E., and Walsh, M. J. (2016). Tall Waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth  
315 (*Amaranthus palmeri*) Seed Production and Retention at Soybean Maturity. *Weed Technology* 30,  
316 284–290. doi:10.1614/WT-D-15-00130.1.
- 317 Scott, D. (2011). The Technological Fix Criticisms and the Agricultural Biotechnology Debate. *J Agric  
318 Environ Ethics* 24, 207–226. doi:10.1007/s10806-010-9253-7.
- 319 Sosnoskie, L. M., Webster, T. M., and Culpepper, A. S. (2013). Glyphosate Resistance Does Not  
320 Affect Palmer Amaranth (*Amaranthus palmeri*) Seedbank Longevity. *Weed Science* 61, 283–288.  
321 doi:10.1614/WS-D-12-00111.1.
- 322 Spaunhorst, D. J., Devkota, P., Johnson, W. G., Smeda, R. J., Meyer, C. J., and Norsworthy, J. K. (2018).  
323 Phenology of Five Palmer amaranth (*Amaranthus palmeri*) Populations Grown in Northern Indiana and  
324 Arkansas. *Weed Science* 66, 457–469. doi:10.1017/wsc.2018.12.
- 325 Wang, J. L., Klessig, D. F., and Berry, J. O. (1992). Regulation of C4 Gene Expression in Developing  
326 Amaranth Leaves. *The Plant Cell* 4, 173–184. doi:10.1105/tpc.4.2.173.
- 327 Ward, S. M., Webster, T. M., and Steckel, L. E. (2013). Palmer Amaranth (*Amaranthus palmeri*): A Review.  
328 *Weed Technology* 27, 12–27. doi:10.1614/WT-D-12-00113.1.
- 329 Webster, T. M., and Grey, T. L. (2015). Glyphosate-Resistant Palmer Amaranth (*Amaranthus palmeri*)  
330 Morphology, Growth, and Seed Production in Georgia. *Weed Science* 63, 264–272. doi:10.1614/WS-D-  
331 14-00051.1.
- 332 Yu, E., Blair, S., Hardel, M., Chandler, M., Thiede, D., Cortilet, A., Gunsolus, J., and Becker, R. (2021).  
333 Timeline of Palmer amaranth (*Amaranthus palmeri*) invasion and eradication in Minnesota. *Weed  
334 Technology*, 1–31. doi:10.1017/wet.2021.32.