

The background of the slide is a light gray gradient. It is decorated with numerous realistic water droplets of various sizes. Some droplets are at the top left, some are along the bottom edge, and others are scattered on the right side. The droplets have highlights and shadows, giving them a three-dimensional appearance.

HERBICIDE MODE-OF-ACTION

SYMPTOMOLOGY, SELECTIVITY, & PHYSIOLOGY

WEED CONTEST PRACTICE

UNIVERSITY OF WISCONSIN-MADISON

2018

The background of the slide is a light gray gradient. It is decorated with several realistic water droplets of various sizes, primarily located in the top-left, top-right, and bottom-right corners. The droplets have highlights and shadows, giving them a three-dimensional appearance.

WHAT'S IN A NAME *AND* WHY SHOULD I CARE?

WHAT'S IN A NAME?

- WHAT DO MAD DOG, BUCCANEER, CORNERSTONE, DURANGO, RANGER, HONCHO, ETC..... HAVE IN COMMON?
- WHAT IF I ADDED ROUNDUP WEATHERMAX AND ROUNDUP POWERMAX TO THE LIST?

ALL ARE EPSPS INHIBITOR HERBICIDES, GROUP 9

AI & PRODUCT EXAMPLES

PRODUCT (TRADE NAME)	ACTIVE INGREDIENT (COMMON NAME)
ROUNDUP + MORE	GLYPHOSATE
AATREX, + MORE	ATRAZINE
RAPTOR	IMAZAMOX
PURSUIT	IMAZETHAPYR
LIBERTY 280	GLUFOSINATE

WAYS TO CATEGORIZE HERBICIDES

- SITE OF UPTAKE
- SOIL OR FOLIAR
- CONTACT VS. SYSTEMIC
- XYLEM VS. PHLOEM MOBILE
- NONSELECTIVE VS. SELECTIVE
- (BROADLEAF OR GRASS)
- MOA
- SOA
- CHEMICAL FAMILY

WAYS HERBICIDES MOVE IN PLANTS

❖CONTACT HERBICIDES


- NO MOVEMENT IN PLANT (GRAMOXONE, COBRA)

❖SYSTEMIC HERBICIDES

- WITHIN THE PLANT (PHLOEM MOBILE)
- FROM SOIL TO TOP OF PLANT (XYLEM MOBILE)

CONTACT HERBICIDES

- ACTIVITY ONLY WHERE CONTACT PLANT TISSUE
- FOLIARLY APPLIED
- COVERAGE CRITICAL!
- EXAMPLES:
 - ✓ PARAQUAT
 - ✓ LACTOFEN



SYSTEMIC HERBICIDES




Figure 8.—Glyphosate: Uptake through leaves and transport through both phloem and xylem.




Figure 7.—Imidazolinones and sulfonyleureas: Uptake through leaves and roots and transport through both phloem and xylem.

HERBICIDE TYPES – SELECTIVITY

❖ NONSELECTIVE

- WILL INJURE ANY PLANT IT CONTACTS **IF**
ABSORBED

❖ SELECTIVE

- WON'T HARM GRASSES
OR
- WON'T HARM
BROADLEAF SPECIES
OR
- RANDOM SELECTIVITY

HERBICIDE MODE AND SITE OF ACTION

- MODE OF ACTION

- HOW THE HERBICIDE CONTROLS THE PLANT

- SITE OF ACTION

- WHERE THE HERBICIDE ACTS WITHIN THE PLANT



07/18/2013 10:28

SYMPTOMOLOGY

MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **SYNTHETIC AUXIN**
2,4-D, DICAMBA, QUINCLORAC, CLOPYRALID



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

ALS INHIBITORS

**NICOSULFURON, HALOSULFURON,
IMAZETHAPYR, CHLORIMURON-ETHYL**



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **PHOTOSYSTEM II INHIBITOR SITE A OR B**
ATRAZINE (A), METRIBUZIN (A), BROMOXYNIL (B)



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **MICROTUBULE ASSEMBLY INHIBITORS**
PENDIMETHALIN;
- **VLCFA INHIBITORS**
ACETOCHLOR, PYROXASULFONE





MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **GLUTAMINE SYNTHETASE INHIBITOR**
GLUFOSINATE







MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **PHOTOSYSTEM I ELECTRON DIVERTER**
PARAQUAT







MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **EPSPS INHIBITOR**
GLYPHOSATE



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **ACCASE INHIBITORS**
CLETHODIM, QUIZALOFOP-P



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **PPO INHIBITORS**
**FOMESAFEN, LACTOFEN, CARFENTRAZONE-ETHYL,
FLUMIOXAZIN, SULFENTRAZONE**



MODE OF ACTION OF HERBICIDES

SYMPTOMOLOGY

- **HPPD INHIBITORS**
ISOXAFLUTOLE, MESOTRIONE, TEMBOTRIONE
- **DOXP SYNTHASE INHIBITOR**
CLOMAZONE



NARROWING DOWN YOUR CHOICES

QUESTIONS TO ASK?

❖ ARE PLANT CARCASSES PRESENT?

- PRE VS. POST

❖ WHAT IS THE ACTIVITY OF THE HERBICIDE?

- CONTACT
- SYSTEMIC



NARROWING DOWN YOUR CHOICES

❖ WHAT IS THE ACTIVITY OF THE HERBICIDE?

- CONTACT
- SYSTEMIC

