16/11/2017 HackerRank

















Points: 25 Rank: 183204



Dashboard > Data Structures > Heap > Find the Running Median

# Find the Running Median



Problem	Submissions	Leaderboard	Discussions	Editorial 🖀

The median of a set of integers is the midpoint value of the data set for which an equal number of integers are less than and greater than the value. To find the median, you must first sort your set of integers in non-decreasing order, then:

- If your set contains an odd number of elements, the median is the middle element of the sorted sample. In the sorted set {1, 2, 3}, 2 is the median.
- If your set contains an even number of elements, the median is the average of the two middle elements of the sorted sample. In the sorted set  $\{1, 2, 3, 4\}$ ,  $\frac{2+3}{2} = 2.5$  is the median.

Given an input stream of n integers, you must perform the following task for each  $i^{th}$  integer:

- 1. Add the  $\emph{i}^{th}$  integer to a running list of integers.
- 2. Find the median of the updated list (i.e., for the first element through the  $i^{th}$  element).
- 3. Print the list's updated median on a new line. The printed value must be a double-precision number scaled to 1 decimal place (i.e., 12.3 format).

#### **Input Format**

The first line contains a single integer, n, denoting the number of integers in the data stream. Each line i of the n subsequent lines contains an integer,  $a_i$ , to be added to your list.

## **Constraints**

- $1 \le n \le 10^5$
- $0 \le a_i \le 10^5$

# **Output Format**

After each new integer is added to the list, print the list's updated median on a new line as a single double-precision number scaled to 1 decimal place (i.e., 12.3 format).

## Sample Input

- 6
- 12
- -
- 3
- 7

#### **Sample Output**

- 12.0
- 8.0 5.0
- 4.5
- 5.0
- 6.0

16/11/2017 HackerRank

#### **Explanation**

There are n=6 integers, so we must print the new median on a new line as each integer is added to the list:

```
1. list = \{12\}, median = 12.0
2. list = \{12,4\} \rightarrow \{4,12\}, median = \frac{12+4}{2} = 8.0
3. list = \{12,4,5\} \rightarrow \{4,5,12\}, median = 5.0
4. list = \{12,4,5,3\} \rightarrow \{3,4,5,12\}, median = \frac{4+5}{2} = 4.5
5. list = \{12,4,5,3,8\} \rightarrow \{3,4,5,8,12\}, median = 5.0
6. list = \{12,4,5,3,8,7\} \rightarrow \{3,4,5,7,8,12\}, median = \frac{5+7}{2} = 6.0
```

f in Submissions:<u>9432</u>
Max Score:50
Difficulty: Hard
Rate This Challenge:
☆☆☆☆☆☆



Join us on IRC at #hackerrank on freenode for hugs or bugs.

Contest Calendar | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy | Request a Feature