



Tower Breakers - The Final Battle

by allllekssssa

Problem

Submissions

Leaderboard

Discussions

Editorial

Our unsung tower-breaking heroes (players P_1 and P_2) only have one tower left, and they've decided to break it for a special game commemorating the end of 5 days of Game Theory! The rules are as follows:

- P_1 always moves first, and both players always move optimally.
- Initially there is 1 tower of height N .
- The players move in alternating turns. The moves performed by each player are different:
 - At each turn, P_1 divides the current tower into some number of smaller towers. If the turn starts with a tower of height H and P_1 breaks it into $x \geq 2$ smaller towers, the following condition must apply: $H = h_1 + h_2 + \dots + h_x$, where h_i denotes the height of the i^{th} new tower.
 - At each turn, P_2 chooses some tower k of the x new towers made by P_1 (where $1 \leq k \leq x$). Then P_1 must pay k^2 coins to P_2 . After that, P_1 gets another turn with tower h_k and the game continues.
- The game is over when no valid move can be made by P_1 , meaning that $H = 1$.
- P_1 's goal is to pay as few coins as possible, and P_2 's goal is to earn as many coins as possible.

Can you predict the number of coins that P_2 will earn?

Input Format

The first line contains a single integer, T , denoting the number of test cases.

Each of the T subsequent lines contains a single integer, N , defining the initial tower height for a test case.

Constraints

- $1 \leq T \leq 100$
- $2 \leq N \leq 10^{18}$

Output Format

For each test case, print a single integer denoting the number of coins earned by P_2 on a new line.

Sample Input

```
3
4
2
7
```

Sample Output

```
6
4
8
```

Explanation

Test Case 0:

Our players make the following moves:

1. $H = N = 4$

1. P_1 splits the initial tower into 2 smaller towers of sizes 3 and 1.

2. P_2 chooses the first tower and earns $1^2 = 1$ coin.

2. $H = 3$

1. P_1 splits the tower into 2 smaller towers of sizes 2 and 1.

2. P_2 chooses the first tower and earns $1^2 = 1$ coin.

3. $H = 2$

1. P_1 splits the tower into 2 smaller towers of size 1.

2. P_2 chooses the second tower and earns $2^2 = 4$ coins.

The total number of coins earned by P_2 is $1 + 1 + 4 = 6$, so we print 6 on a new line.

f t in

Submissions: 195



Max Score: 50

Difficulty: Medium

Rate This Challenge:

☆☆☆☆☆

More

Current Buffer (saved locally, editable)  

Java 7



```
1 import java.io.*;
2 import java.util.*;
3 import java.text.*;
4 import java.math.*;
5 import java.util.regex.*;
6
7 public class Solution {
8
9     public static void main(String[] args) {
10         /* Enter your code here. Read input from STDIN. Print output to STDOUT. Your class should be named Solution. */
11     }
12 }
```

Line: 1 Col: 1

 Upload Code as File

☐ Test against custom input

Run Code

Submit Code

Join us on IRC at [#hackerrank](#) on freenode for hugs or bugs.

[Contest Calendar](#) | [Blog](#) | [Scoring](#) | [Environment](#) | [FAQ](#) | [About Us](#) | [Support](#) | [Careers](#) | [Terms Of Service](#) | [Privacy Policy](#) | [Request a Feature](#)