

Calculation of diatom-atom collisions in a magnetic field

Marcin Welter

Winter semester 2024/2025

1 Alkali like atom + atom collision

For the problem of collision of two alkali like atoms $A + B$ we consider the following Hamiltonian

$$\hat{H} = -\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} + \frac{\hat{L}^2}{2\mu R^2} + \hat{H}_{\text{int}}^A + \hat{H}_{\text{int}}^B + \hat{V}(R), \quad (1)$$

where R is the distance between atoms, μ is the reduced mass of two atoms, \hat{L}^2 is the angular momentum of the two atoms, \hat{H}_{int}^k is the internal structure of a k atom and $\hat{V}(R)$ consists of triplet and singlet potential curves.

1.1 Hyperfine and Zeeman structure

Internal structure of alkali like atoms in the magnetic field consists of Zeeman splitting and hyperfine structure. The internal hamiltonian for such an atom k is written as

$$\hat{H}_{\text{int}}^k = \hat{H}_{\text{Zee}}^k + \hat{H}_{\text{hf}}^k, \quad (2)$$

with $\hat{H}_{\text{Zee}}^k = -\gamma_e B \hat{s}_z^k - \gamma_n^k B \hat{i}_z^k$ and $\hat{H}_{\text{hf}}^k = a_{\text{hf}}^k \hat{\mathbf{s}}^k \cdot \hat{\mathbf{i}}^k$.

In the basis $|sm_s im_i\rangle \equiv |m_s m_i\rangle$ the matrix elements of the internal Hamiltonian are

$$\langle m_s m_i | H_{\text{Zee}}^k | m'_s m'_i \rangle = -B(\gamma_e m_s + \gamma_n^k m_i) \delta_{m_s m'_s}^{m'_i m_i}, \quad (3)$$

for Zeeman part, and the hyperfine part is

$$\langle m_s m_i | H_{\text{hf}}^k | m'_s m'_i \rangle = a_{\text{hf}}^k \left(m_s m_i \delta_{m_s m'_s}^{m'_i m_i} + \frac{1}{2} \sum_{p=\pm 1} C(s, m_s, m'_s) C(i, m_i, m'_i) \delta_{m_s m'_s}^{m'_i m_i + p, m'_i - p} \right), \quad (4)$$

where $C(s, m_s, m'_s) = \sqrt{s(s+1) - m_s m'_s}$. We see that the total projection $m_f = m_i + m_s$ is preserved.

1.2 Singlet and triplet potential curves

Potential term consists of singlet and triplet potential curves, we can write it as

$$\hat{V}(R) = \hat{\mathcal{P}}_1 V_1(R) + \hat{\mathcal{P}}_0 V_0(R), \quad (5)$$

where \mathcal{P}_S is a projection to the subspace with combined electron spin of two atoms equal to S and $V_S(R)$ is a corresponding potential curve.

In the electron spin basis $|m_s^A, m_s^B\rangle$ of each atom, the matrix elements of the projection operator are

$$\langle m_s^A m_s^B | \mathcal{P}_S | m_s'^A m_s'^B \rangle = \sum_{M_S} \langle m_s^A m_s^B | S M_S \rangle \langle S M_S | m_s'^A m_s'^B \rangle. \quad (6)$$

1.3 Full basis

Full basis is then $|m_s^A m_s^B m_i^A m_i^B\rangle$ with total spin projection M_F being conserved. Additionally, angular momentum and its projection are good quantum numbers. For homo-nuclear we can transform the basis to the combined electron and nuclear spin for which the symmetry is evident and we restrict ourselves to only the needed symmetry of the system.

2 Spinless rigid rotor + atom collision

Here we consider a diatom + atom collision without spin structure in a rigid rotor approximation. The Hamiltonian for such a system is

$$\hat{H} = -\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} + \frac{\hat{L}^2}{2\mu R^2} + \hat{H}_{\text{rot}} + V(R, \theta) \quad (7)$$

where R, θ are the Jacobi coordinates, μ is the reduced mass of the atom and diatom, \hat{L}^2 is the orbital angular momentum of the system, \hat{H}_{rot} is the Hamiltonian of the rigid-rotor and $\hat{V}(R, \theta)$ is the potential surface.

Rigid rotor Hamiltonian is

$$\hat{H}_{\text{rot}} = B_e \hat{N}^2, \quad (8)$$

where B_e is a rotational constant of the rotor.

We work in $|(lN)JM_J\rangle$ basis being the combined angular momentum basis J of orbital angular momentum of the system l and rotation of the molecule N . In this basis the matrix elements of the potential terms are

$$\langle (lN)JM_J | V(R, \theta) | (l'N')J'M_J' \rangle = \sum_{\lambda} V_{\lambda}(R) f_{\lambda}(lN, l'N'; J) \delta_{J'M_J'}^{JM_J}, \quad (9)$$

where $V_{\lambda}(R)$ is the λ -th term of the Legendre decomposition of the potential $V(R, \theta)$ and

$$f_{\lambda}(lN, l'N'; J) = (-1)^{l+l'+J} \sqrt{(2N+1)(2N'+1)(2l+1)(2l'+1)} \\ \times \begin{pmatrix} l & \lambda & l' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} N & \lambda & N' \\ 0 & 0 & 0 \end{pmatrix} \left\{ \begin{matrix} l & \lambda & l' \\ N' & J & N \end{matrix} \right\}, \quad (10)$$

are the Percival-Seaton coefficients. Other terms in the Hamiltonian are straightforward and total angular momentum J and its projection is conserved M_J . Additionally if the potential only have even numbered terms in the Legendre decomposition, divisibility of l, N by two is conserved.

3 Alkali like rigid rotor + atom collision

We now consider a alkali like diatom + atom collision, by that we mean that the diatom consists of alkali like atom and a structureless atom that collide with the alkali like atom. The Hamiltonian for such a system, following Ref. [1] is

$$\hat{H} = -\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} + \frac{\hat{L}^2}{2\mu R^2} + \hat{H}_{\text{int}}^{\text{rot}} + \hat{H}_{\text{int}}^{\text{A}} + \hat{\mathcal{P}}_1 V_1(R, \theta) + \hat{\mathcal{P}}_0 V_0(R, \theta), \quad (11)$$

where internal atom Hamiltonian and projection operators are the same as in Sec. 1 and the internal rotor hamiltonian can be further written as

$$\hat{H}_{\text{int}}^{\text{rot}} = \hat{H}_{\text{rot}} + \hat{H}_{\text{spin-rot}} + \hat{H}_{\text{ahf}} + \hat{H}_{\text{spin}}. \quad (12)$$

The first term is the same as without the spin that is $\hat{H}_{\text{rot}} = B_e \hat{N}^2$, second term is the coupling between rotor rotation and electron spin of the rotor

$$\hat{H}_{\text{spin-rot}} = \gamma_{\text{sr}} \hat{\mathbf{N}} \cdot \hat{\mathbf{s}}^{\text{r}}, \quad (13)$$

where γ_{sr} is a spin-rot coupling constant,

$$\hat{H}_{\text{ahf}} = \frac{c\sqrt{6}}{3} \left(\frac{4\pi}{5} \right)^{1/2} \sum_{q=-2}^2 (-1)^q Y_{2-q}(\theta_r, \phi_r) [\hat{\mathbf{r}} \otimes \hat{\mathbf{s}}^r]_q^{(2)}, \quad (14)$$

is an anisotropic hyperfine contribution with anisotropic hyperfine constant being c . Last term is the hyperfine and Zeeman structure which is the same as in the atom case in Sec. 1.

The basis used is $|(lN)JM_J\rangle |(s^r)m_s^r\rangle |(i^r)m_i^r\rangle |(s^a)m_s^a\rangle |(i^a)m_i^a\rangle$. In this basis, matrix elements of the spin-rot contribution are

$$\begin{aligned} \langle (lN)JM_J | \langle m_s^r | \hat{H}_{\text{spin-rot}} | (l'N')JM_J \rangle | m_s'^r \rangle &= \delta_{lN}^{l'N'} \gamma_{\text{sr}} p_3(N) p_3(s^r) [(2J+1)(2J'+1)]^{1/2} \\ &\times \sum_p (-1)^p (-1)^{J-M_J} (-1)^{N+l+J'+1} (-1)^{s^r-m_s^r} \\ &\times \left\{ \begin{matrix} N & J & l \\ J' & N & 1 \end{matrix} \right\} \begin{pmatrix} J & 1 & J' \\ -M_J & p & M_J' \end{pmatrix} \begin{pmatrix} s^r & 1 & s^r \\ -m_s^r & -p & m_s'^r \end{pmatrix} \end{aligned} \quad (15)$$

where $p_3(X) = \sqrt{X(2X+1)(X+1)}$ and matrix elements of the anisotropic hyperfine contribution are

$$\begin{aligned} \langle (lN)JM_J | \langle m_s^r m_i^r | \hat{H}_{\text{ahf}} | (l'N')J'M_J' \rangle | m_s'^r m_i'^r \rangle &= \delta_{lN}^{l'N'} c \frac{\sqrt{30}}{3} (-1)^{J-M_J+l+J'} p_3(i^r) p_3(s^r) [(2J+1)(2J'+1)]^{1/2} [(2N+1)(2N'+1)]^{1/2} \\ &\times \left\{ \begin{matrix} N & J & l \\ J' & N' & 2 \end{matrix} \right\} \begin{pmatrix} 1 & 1 & 2 \\ m_i^r - m_i'^r & m_s^r - m_s'^r & M_J - M_J' \end{pmatrix} \begin{pmatrix} N & 2 & N' \\ 0 & 0 & 0 \end{pmatrix} \\ &\times \begin{pmatrix} J & 2 & J' \\ -M_J & M_J - M_J' & M_J' \end{pmatrix} \begin{pmatrix} i^r & 1 & i^r \\ -m_i^r & m_i^r - m_i'^r & m_i'^r \end{pmatrix} \begin{pmatrix} s^r & 1 & s^r \\ -m_s^r & m_s - m_s'^r & m_s'^r \end{pmatrix}. \end{aligned} \quad (16)$$

4 Reconstructing results

4.1 Two Li^6 feshbach resonances




Figure 1: Magnetic Feshbach resonances between two Li_6 in the lowest channel, reconstruction of [2].

4.2 Spinless AlF + Rb cross sections




Figure 2: Reconstruction of elastic cross section of AlF + Rb vs maximum value of N .

4.3 SrF + Rb Feshbach resonances




Figure 3: Magnetic Feshbach resonances of SrF + Rb, reconstruction of [3] in supplementary materials.

References

- [1] Timur V Tscherbul and Jose P D’Incao. Ultracold molecular collisions in magnetic fields: Efficient incorporation of hyperfine structure in the total rotational angular momentum representation. *Physical Review A*, 108(5):053317, 2023.
- [2] Paul S Julienne and Jeremy M Hutson. Contrasting the wide feshbach resonances in Li^6 and Li^7 . *Physical Review A*, 89(5):052715, 2014.
- [3] Masato Morita, Maciej B Kosicki, Piotr S Żuchowski, Paul Brumer, and Timur V Tscherbul. Magnetic feshbach resonances in ultracold atom-molecule collisions. *Physical Review A*, 110(2):L021301, 2024.